

Name:

CID:

Tutorial 7

Any marks received for the tutorial are only indicative and may be subject to moderation and scaling.

Exercise 1 (Absolute stability)	% of CW mark: 1.0
--	--------------------------

Find the interval of absolute stability of the Trapezoidal rule.

Exercise 2 (Absolute stability)	% of CW mark: 1.0
--	--------------------------

Find the restriction on the time step for which Euler's method is absolutely stable when applied to the equation

$$x' = \lambda x, \quad \lambda = -4 + 3i.$$

Exercise 3 (Absolute stability for systems)	% of CW mark: 1.5
--	--------------------------

Find the restriction on the time step for which Euler's method is absolutely stable when applied to the system of equations

$$\begin{pmatrix} u_1' \\ u_2' \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}.$$

Exercise 4 (PECE)	% of CW mark: 2.0
--------------------------	--------------------------

Mastery Component

Combine the Euler method and Trapezoidal rule into a predictor-corrector method, and calculate the local truncation error and interval of absolute stability of the method when applied to the equation

$$x' = \lambda x, \quad \operatorname{Re}(\lambda) < 0.$$