

DEPARTMENT OF ELECTRICAL AND ELECTRONIC ENGINEERING
EXAMINATIONS 2016

MSc and EEE/EIE PART IV: MEng and ACGI

Corrected copy

SYSTEMS IDENTIFICATION

Tuesday, 3 May 10:00 am

Time allowed: 3:00 hours

There are FOUR questions on this paper.

Answer FOUR questions.

All questions carry equal marks

Any special instructions for invigilators and information for candidates are on page 1.

Examiners responsible First Marker(s) : T. Parisini
Second Marker(s) : S.A. Evangelou

1. Consider a discrete-time stochastic system described by the block-scheme shown in Fig. 1.1.




Figure 1.1 Block-scheme of the stochastic dynamic system.

where $u(t)$ is a known deterministic input and $e(\cdot) \sim WN(0, 1)$ is a white stochastic process.

- a) Show that the stochastic system depicted in Fig. 1.1 can be described by a model belonging to the family of $ARMAX(n, n)$ stochastic models in canonical form, that is

$$A(z)y(t) = B(z)u(t) + C(z)e(t)$$

where $A(z) = 1 + a_1z^{-1} + \dots + a_nz^{-n}$, $B(z) = b_1z^{-1} + \dots + b_nz^{-n}$, and $C(z) = 1 + c_1z^{-1} + \dots + c_nz^{-n}$.

[6 Marks]

- b) Determine the difference equation by which the optimal one-step ahead prediction $\hat{y}(t+1|t)$ can be computed using the knowledge of the sequences $y(t)$ and $u(t)$ till time-instant t .

[3 Marks]

- c) Set $u(t) = 0, \forall t \geq 0$.

- i) Sketch approximately the spectrum $\Gamma_y(\omega)$ of the process $y(\cdot)$ described by the model obtained in your answer to Question 1a) in the interval $\omega \in [-\pi, \pi]$.

[4 Marks]

- ii) Determine the difference equation by which the optimal two-step ahead prediction $\hat{y}(t+2|t)$ can be computed using the knowledge of the sequence $y(t)$ till time-instant t .

[4 Marks]

- iii) Compute the variance of the prediction error $\varepsilon_1(t) = y(t+1) - \hat{y}(t+1|t)$ associated with the optimal one-step ahead predictor determined in your answer to Question 1b) and the variance of the prediction error $\varepsilon_2(t) = y(t+2) - \hat{y}(t+2|t)$ associated with the optimal two-step ahead predictor determined in your answer to Question 1c)ii). Compare the two variances and discuss your findings.

[3 Marks]

2. Consider a stochastic process $v(\cdot)$ and an arbitrarily large number N of measurements $\{v(1), v(2), \dots, v(N)\}$. Moreover, consider the following two families of AR models:

$$\begin{aligned}\mathcal{M}_1(\theta_1): \quad & v(t) = av(t-1) + \xi(t), \quad \theta_1 = a \\ \mathcal{M}_2(\theta_2): \quad & v(t) = a_1v(t-1) + a_2v(t-2) + \xi(t), \quad \theta_2 = [a_1, a_2]^T.\end{aligned}$$

where $\xi(\cdot)$ is a generic white process uncorrelated with $v(\cdot)$. Denote with $\hat{\theta}_1(N) = \hat{a}(N)$ the least squares estimate of θ_1 based on N measurements $\{v(1), v(2), \dots, v(N)\}$. Likewise denote with $\hat{\theta}_2(N) = [\hat{a}_1(N), \hat{a}_2(N)]^T$ the least squares estimate of θ_2 based on N measurements $\{v(1), v(2), \dots, v(N)\}$.

- a) **Case 1.** Suppose that the process is generated as

$$v(t) = e(t) + \frac{1}{2}e(t-1), \quad e(\cdot) \sim WN(0, 1)$$

- i) Determine the value $\bar{\theta}_1^{(1)}$ the estimate $\hat{\theta}_1(N)$ approaches for large values of N (that is, $\bar{\theta}_1^{(1)} = \lim_{N \rightarrow \infty} \hat{\theta}_1(N)$, a.s.) when the family \mathcal{M}_1 of models is used.

[3 Marks]

- ii) Determine the value $\bar{\theta}_2^{(1)}$ the estimate $\hat{\theta}_2(N)$ approaches for large values of N (that is, $\bar{\theta}_2^{(1)} = \lim_{N \rightarrow \infty} \hat{\theta}_2(N)$, a.s.) when the family \mathcal{M}_2 of models is used.

[3 Marks]

- iii) Compute the variances of the prediction errors when using the models $\mathcal{M}_1(\bar{\theta}_1^{(1)})$ and $\mathcal{M}_2(\bar{\theta}_2^{(1)})$ and compare them. Discuss your findings.

[3 Marks]

- iv) Determine the analytical expressions of the correlation function $\gamma_e(\tau)$, $\forall \tau \geq 0$ and of the spectrum $\Gamma_e(\omega)$, $\omega \in [-\pi, \pi]$ of the prediction error $\varepsilon(\cdot)$ when the model $\mathcal{M}_1(\bar{\theta}_1^{(1)})$ is used.

[3 Marks]

- b) **Case 2.** Suppose that the process is generated as

$$v(t) = e(t) + \frac{1}{2}e(t-1) + \frac{1}{4}e(t-2), \quad e(\cdot) \sim WN(0, 1)$$

- i) Determine the value $\bar{\theta}_1^{(2)}$ the estimate $\hat{\theta}_1(N)$ approaches for large values of N (that is, $\bar{\theta}_1^{(2)} = \lim_{N \rightarrow \infty} \hat{\theta}_1(N)$, a.s.) when the family \mathcal{M}_1 of models is used.

[4 Marks]

- ii) Determine the value $\bar{\theta}_2^{(2)}$ the estimate $\hat{\theta}_2(N)$ approaches for large values of N (that is, $\bar{\theta}_2^{(2)} = \lim_{N \rightarrow \infty} \hat{\theta}_2(N)$, a.s.) when the family \mathcal{M}_2 of models is used.

[4 Marks]

3. Consider the discrete-time stochastic dynamic system described by the block-scheme shown in Fig. 3.1.




Figure 3.1 Block-scheme of the stochastic dynamic system.

where $\eta(\cdot) \sim WN(1, 1)$ and $\xi(\cdot) \sim WN(3, 1)$ are white non-zero mean uncorrelated stochastic processes.

- Show that the steady-state stochastic process $y(\cdot)$ is stationary.
[3 Marks]
- Compute the expected value $m_v = \mathbb{E}(v)$ of the steady-state stochastic process $v(\cdot)$.
[3 Marks]
- Compute the expected value $m_w = \mathbb{E}(w)$ of the steady-state stochastic process $w(\cdot)$.
[3 Marks]
- Compute the covariance matrix of the stochastic processes $v(\cdot)$ and $w(\cdot)$, given by

$$\Lambda = \begin{bmatrix} \lambda_{vv} & \lambda_{vw} \\ \lambda_{wv} & \lambda_{ww} \end{bmatrix}$$

where

$$\lambda_{vv} = \text{var}[v(t)] = \mathbb{E}[(v(t) - m_v)^2]$$

$$\lambda_{ww} = \text{var}[w(t)] = \mathbb{E}[(w(t) - m_w)^2]$$

$$\lambda_{vw} = \text{cov}[v(t), w(t)] = \mathbb{E}[(v(t) - m_v)(w(t) - m_w)]$$

- Determine the transfer function $G_{\eta y}(z)$ from the input $\eta(t)$ to the output $y(t)$ and the transfer function $G_{\xi y}(z)$ from the input $\xi(t)$ to the output $y(t)$.
[7 Marks]

[4 Marks]

4. Consider three visual sensors VS_1 , VS_2 , and VS_3 respectively providing three samples x_i , $i = 1, 2, 3$ of the position x of a given object O as depicted in Fig. 4.1.




Figure 4.1 Scheme of the position measurements provided by three visual sensors VS_1 , VS_2 , and VS_3 .

x_i , $i = 1, 2, 3$ are realisations of *Gaussian and independent* random variables:

$$x_1 \sim \mathcal{G}(\bar{x}, 1/3), \quad x_2 \sim \mathcal{G}(0.1 + \bar{x}, 2), \quad x_3 \sim \mathcal{G}(\bar{x}, 1).$$

where the second measurement is affected by a bias equal to 0.1. The objective is to design an estimator of the mean position \bar{x} as a function of the samples x_i , $i = 1, 2, 3$.

- a) Consider the empirical mean estimator \tilde{x} of \bar{x} given by:

$$\tilde{x} = \frac{1}{3}(x_1 + x_2 + x_3)$$

Compute $\mathbb{E}(\tilde{x})$ and $\text{var}(\tilde{x})$.

[3 Marks]

- b) Consider an estimator of the form

$$\hat{x}(a, b, c, d) = ax_1 + bx_2 + cx_3 + d,$$

where $a, b, c, d \in \mathbb{R}$ are four parameters to be determined. Determine, a set of constraints on a, b, c, d guaranteeing that the estimator $\hat{x}(a, b, c, d)$ is unbiased.

[4 Marks]

- c) Consider an estimator of the form

$$\tilde{x}(\alpha, \beta, \gamma) = \alpha x_1 + \beta x_2 + \gamma x_3,$$

where $\alpha, \beta, \gamma \in \mathbb{R}$ are three parameters to be determined. Determine, a set of constraints on α, β, γ guaranteeing that the estimator $\tilde{x}(\alpha, \beta, \gamma)$ is unbiased.

[4 Marks]

- d) Among the values of a, b, c, d determined in your answer to Question 4b) and among the values of α, β, γ determined in your answer to Question 4c), find:

$$(a^\circ, b^\circ, c^\circ, d^\circ) = \arg \min_{a, b, c, d} \text{var}[\hat{x}(a, b, c, d)]; \quad (\alpha^\circ, \beta^\circ, \gamma^\circ) = \arg \min_{\alpha, \beta, \gamma} \text{var}[\tilde{x}(\alpha, \beta, \gamma)]$$

that is, find $(a^\circ, b^\circ, c^\circ, d^\circ)$ and $(\alpha^\circ, \beta^\circ, \gamma^\circ)$ such that $\text{var}(\hat{x})$ and $\text{var}(\tilde{x})$, respectively, take on the smallest value possible.

[6 Marks]

- e) Compute $\text{var}[\hat{x}(a^\circ, b^\circ, c^\circ, d^\circ)]$ and $\text{var}[\tilde{x}(\alpha^\circ, \beta^\circ, \gamma^\circ)]$. Compare these variances among themselves and also with $\text{var}(\tilde{x})$ obtained in the answer to Question 4a). Discuss your findings.

[3 Marks]

