

IMPERIAL COLLEGE LONDON

DEPARTMENT OF ELECTRICAL AND ELECTRONIC ENGINEERING
EXAMINATIONS 2016

EEE PART I: MEng, BEng and ACGI

Corrected Copy

MATHEMATICS 1B (E-STREAM AND I-STREAM)

Tuesday, 24 May 10:00 am

Time allowed: 2:00 hours

There are FOUR questions on this paper.

Answer ALL questions. All questions carry equal marks (25% each)

NO CALCULATORS ALLOWED

Any special instructions for invigilators and information for candidates are on page 1.

Examiners responsible First Marker(s) : D. Nucinkis, D. Nucinkis
Second Marker(s) : I.M. Jaimoukha, I.M. Jaimoukha

EE1-10B MATHEMATICS II

1. a) Obtain the Fourier transform of the rectangular pulse function

$$f(t) = \begin{cases} 1, & |t| < T/2, \\ 0, & |t| \geq T/2, \end{cases}$$

where T is a positive constant. [5]

- b) Given that a function $g(t)$ has Fourier transform $G(\omega)$, prove the frequency shift property, that is,

$$\mathcal{F} [e^{i\omega_0 t} g(t)] = G(\omega - \omega_0),$$

where \mathcal{F} denotes the Fourier transform. [5]

- c) Two planes, Π_1 and Π_2 , have cartesian equations

$$-2x + y + 2z = 1, \text{ and } x - 2y + z = 3.$$

Find the intersection of Π_1 and Π_2 and give a geometric interpretation. [5]

- d) Let

$$A = \begin{pmatrix} \alpha & -1 \\ 1 & 0 \end{pmatrix}.$$

- i) Find the values of α such that $A^n = I$, where I is the identity matrix, for $n = 3$ and $n = 4$. [5]
- ii) Prove that A^{-1} exists for all α , but that there is no value of α such that $A = A^{-1}$. [5]

2. a) The vector $\underline{y} \in \mathbb{R}^3$ satisfies

$$\underline{y} = \underline{a} + \underline{b} \times \underline{y},$$

where \underline{a} and \underline{b} are given vectors. By taking the scalar and vector products of this equation with \underline{b} , or otherwise, solve for \underline{y} in terms of \underline{a} and \underline{b} . [6]

- b) Given a matrix and two vectors:

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ -1 & 1 & \alpha \end{pmatrix} \quad \text{and} \quad \underline{b} = \begin{pmatrix} 1 \\ \beta \\ 2 \end{pmatrix} \quad \text{and} \quad \underline{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

- i) Find $\det(A)$ and hence, or otherwise, give conditions on α and β so that the linear system of equation with matrix

$$A\underline{x} = \underline{b}$$

has (I) no solutions; (II) a unique solution; (III) infinitely many solutions; [5]

- ii) For $\alpha = 2$, use Gaussian elimination to obtain the inverse of A . [4]

- iii) Solve the system of equations

$$\begin{aligned} -x + y + 2z &= 1 \\ x + y + z &= 2 \\ x + 2y + 3z &= 5. \end{aligned} \quad [2]$$

- c) Given the symmetric matrix

$$B = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

- i) Obtain the eigenvalues and normalized eigenvectors of B . [4]
 ii) Find an orthogonal diagonalizing matrix P and a diagonal matrix D satisfying

$$B = PDP^T,$$

and without further calculation, write down P^{-1} . [4]

3. a) Find the general solution of the differential equation

$$\left(2xt e^{x^2} + t^2 \sin x + \sec^2 x\right) \frac{dx}{dt} = 2t \cos x + \cos t - e^{x^2}. \quad [6]$$

- b) Consider the following second order differential equation:

$$\frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + 2y = 85 \cos(3x). \quad (1)$$

- i) Obtain the complementary function for (1). [2]
- ii) Obtain a particular integral for (1). [4]
- iii) Hence obtain the general solution of the ODE in (1). [1]

- c) Find the solution of the first-order ordinary differential equation

$$t^2 \frac{dx}{dt} = x^2 + xt + t^2,$$

which satisfies the initial condition $x = 1$ when $t = 1$. [6]

- d) Consider the following first order differential equation:

$$\cos y \frac{dy}{dx} = 3x + \sin y. \quad (2)$$

- i) Use the transformation $u = 3x + \sin y$ to rewrite the differential equation in terms of x and u . [2]
- ii) Hence, or otherwise, obtain the solution u to the differential equation in part (i). [2]
- iii) Hence find the solution y of the differential equation (2) which satisfies the initial condition $y = 0$ when $x = 0$. [2]

4. a) The function $f(x, t)$ satisfies the wave equation:

$$\frac{\partial^2 f}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 f}{\partial t^2}.$$

- i) By considering the change of coordinates

$$\rho = x - ct, \quad \phi = x + ct,$$

and using the chain rule, obtain

$$\frac{\partial f}{\partial x}, \frac{\partial f}{\partial t}, \frac{\partial^2 f}{\partial x^2}, \frac{\partial^2 f}{\partial t^2}$$

in terms of derivatives with respect to ρ and ϕ . [6]

- ii) Use the results of (i) to show that the wave equation in the new variables is

$$\frac{\partial^2 f}{\partial \rho \partial \phi} = 0.$$

[2]

- iii) Hence find the general solution of the wave equation for f in terms of x and t . [4]

- b) Let $g(x, y) = (x - y)(xy - 1)$.

- i) Evaluate the gradient of g and derive its stationary points. [4]
- ii) Evaluate the Hessian of g and use it to classify the stationary points of g . Justify your classification. [4]
- iii) Evaluate the solutions of $g(x, y) = 0$ and hence, or otherwise, sketch the contours of this function. [5]

