

UNIVERSITY OF LONDON

[II(3)E 2004]

B.ENG. AND M.ENG. EXAMINATIONS 2004

For Internal Students of Imperial College London

This paper is also taken for the relevant examination for the Associateship of the City & Guilds of London Institute

PART II : MATHEMATICS 3 (ELECTRICAL ENGINEERING)

Wednesday 2nd June 2004 2.00 - 5.00 pm

Answer EIGHT questions.

Answers to Section A questions must be written in a different answer book from answers to Section B questions.

Corrected Copy

[Before starting, please make sure that the paper is complete; there should be 7 pages, with a total of 12 questions. Ask the invigilator for a replacement if your copy is faulty.]

Copyright of the University of London 2004

1. Consider the complex mapping

$$w = \frac{z}{z-1}$$

from the z -plane where $z = x + iy$ to the w -plane where $w = u(x, y) + iv(x, y)$.

- (i) Show that circles $(x-1)^2 + y^2 = a^2$ in the z -plane map to circles

$$(u-1)^2 + v^2 = a^{-2}$$

in the w -plane. For some value of $a \neq 1$, make separate sketches of each circle. Show that for any value of $a \neq 1$, if both fixed points of the mapping (that is, points that satisfy $w = z$) lie inside one of the circles then they must lie outside the other and vice-versa.

- (ii) Show also that the y -axis in the z -plane maps to the circle centred at $(\frac{1}{2}, 0)$ and radius $\frac{1}{2}$ in the w -plane.
- (iii) Given the straight lines in the z -plane of the form $y = m(x-1)$, show that for arbitrary finite values of m , these map to the lines

$$mu + v = m$$

in the w -plane.

2. By choosing a suitable closed contour C in the upper half of the complex plane for the complex integral

$$\oint_C \frac{e^{2iz} dz}{(z^2 + 4)(z^2 + 9)},$$

use the Residue Theorem to show that

$$\int_{-\infty}^{\infty} \frac{\cos 2x dx}{(x^2 + 4)(x^2 + 9)} = \frac{\pi}{5} \left(\frac{e^{-4}}{2} - \frac{e^{-6}}{3} \right).$$

PLEASE TURN OVER

3. The contour integral

$$\oint_C \frac{e^{iz}}{z} dz$$

is taken around a closed contour C that contains no poles. C is comprised of

- (i) A semi-circle in the upper half-plane of radius R ;
- (ii) A small semi-circular indentation around the pole at $z = 0$ that has radius r and which lies in the upper half-plane ;
- (iii) Those two parts of the x -axis, namely $(-R, 0) \rightarrow (-r, 0)$ and $(r, 0) \rightarrow (R, 0)$, that connect the two semi-circles .

By considering the integral I in the limits $r \rightarrow 0$ and $R \rightarrow \infty$, show that

$$\int_{-\infty}^{\infty} \frac{\sin x}{x} dx = \pi .$$

4. The square-wave $\Pi(t)$, the tent function $\Lambda(t)$ and the sinc-function $\text{sinc } t$, are defined respectively by

$$\Pi(t) = \begin{cases} 1, & -1/2 \leq t \leq 1/2, \\ 0, & \text{otherwise.} \end{cases}$$

$$\Lambda(t) = \begin{cases} 1+t, & -1 \leq t \leq 0, \\ 1-t, & 0 \leq t \leq 1, \\ 0, & \text{otherwise,} \end{cases}$$

$$\text{sinc } t = \frac{\sin(t/2)}{(t/2)} .$$

- (i) Show that the Fourier transform of $\Pi(t)$ is given by

$$\overline{\Pi}(\omega) = \text{sinc } \omega .$$

- (ii) Show that the Fourier transform of $\Lambda(t)$ is given by

$$\overline{\Lambda}(\omega) = \text{sinc}^2 \omega .$$

- (iii) Given that

$$\int_{-\infty}^{\infty} \frac{e^{ipt}}{t} dt = \begin{cases} +i\pi & p > 0 \\ -i\pi & p < 0 \end{cases}$$

where p is an arbitrary real number, show directly that the Fourier transform of $\text{sinc } t$ is $2\pi\Pi(\omega)$.

5. Show that the Dirac delta-function has an integral representation of the form

$$\int_{-\infty}^{\infty} e^{\pm i \tau \omega} d\omega = 2\pi \delta(\tau)$$

or with τ and ω reversed.

Hence, prove Plancherel's integral relation between the two functions $f(t)$ and $g(t)$ and their Fourier transforms $\bar{f}(\omega)$ and $\bar{g}(\omega)$

$$\int_{-\infty}^{\infty} f(t)g^*(t) dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} \bar{f}(\omega)\bar{g}^*(\omega) d\omega,$$

where $*$ represents the complex conjugate.

If $f(t) = e^{-|t|}$ and $g(t) = \cos \Omega t$, where Ω is a constant frequency, show that

$$\int_{-\infty}^{\infty} e^{-|t|} \cos \Omega t dt = \frac{2}{1 + \Omega^2}.$$

6. A function $y(t)$ satisfies the differential equation

$$\ddot{y} + 5\dot{y} + 6y = f(t),$$

subject to the initial conditions $y(0) = y_0$; $\dot{y}(0) = -2y_0$, where y_0 is a constant. $f(t)$ is a given function of t . Using a Laplace transform and the Laplace Convolution Theorem, obtain the solution of this differential equation in the form

$$y(t) = y_0 e^{-2t} + \int_0^t \left\{ e^{-2(t-u)} - e^{-3(t-u)} \right\} f(u) du.$$

PLEASE TURN OVER

7. P and Q are continuous functions of x and y with continuous first partial derivatives in a simply connected region R with a piecewise smooth boundary C . Use Green's Theorem in a plane to find a two-dimensional vector \mathbf{u} , defined in terms of P and Q , to show that Green's theorem can be re-expressed as the two-dimensional version of the Divergence Theorem

$$\oint_C \mathbf{u} \cdot \mathbf{n} \, ds = \iint_R \operatorname{div} \mathbf{u} \, dx dy$$

where \mathbf{n} is the unit normal to the curve C .

If $\mathbf{u} = i x^2 + j y^2$ and R is the region between the pair of parabolae $2y = x^2$ and $2x = y^2$ in the first quadrant, evaluate the double integral directly to show that

$$\iint_R \operatorname{div} \mathbf{u} \, dx dy = 24/5$$

Green's Theorem in a plane says that

$$\oint_C (P \, dx + Q \, dy) = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \, dx dy.$$

8. The double integral I is given by

$$I = \iint_R (x+y)^n f(x^2 - y^2) \, dx \, dy ,$$

where $n > 0$ is an integer and f is an arbitrary function. The domain of integration R is the finite region in the x - y plane enclosed by the lines $x = 0$, $y = 0$ and $y = 1 - x$.

- (i) Show that, after the variable transformation,

$$u = x^2 - y^2, \quad v = x + y ,$$

the integral can be written as

$$I = \frac{1}{2} \int_0^1 v^{n-1} \left(\int_{-v^2}^{v^2} f(u) \, du \right) dv .$$

- (ii) Evaluate the integral for the special case $n = 2$, $f(u) \equiv e^u$.
 (iii) Evaluate the integral for the special case $n = 0$, and $f(u) \equiv 1$.

Give a simple interpretation of your result.

9. A vector field \mathbf{F} is defined as

$$\mathbf{F}(x, y, z) = 2x \sin z \mathbf{i} + ze^y \mathbf{j} + (x^2 \cos z + ae^y) \mathbf{k},$$

where a is a constant.

- (i) Find $\text{div } \mathbf{F}$ and $\text{curl } \mathbf{F}$.
- (ii) Find the value of a for which there exists a scalar function $\phi(x, y, z)$ such that $\mathbf{F} = \nabla \phi$ and find $\phi(x, y, z)$.
- (iii) Find $(\mathbf{F} \cdot \nabla)\phi$ and $\nabla^2 \phi$ for the ϕ obtained in (ii).

10. (i) The two-dimensional vector field \mathbf{F} is defined by $\mathbf{F} = (y \cos x, 3y + \sin x, 0)$.

Show from Green's theorem in the plane that the line integral

$$I = \int_C (F_1 dx + F_2 dy)$$

depends only on the end points A and B of the path C and is otherwise independent of C .

Find a potential function $\phi(x, y)$ such that $\mathbf{F} = \nabla \phi$ and hence evaluate the integral I when A is the point $(0, 0)$ and B is the point $(\pi/2, 1)$.

- (ii) Let R be the region in the first quadrant of the xy -plane bounded by the ellipse $(x/2)^2 + y^2 = 1$ and the lines $x = 0, y = 0$. Let C be the boundary of R taken in the counter-clockwise direction.

Using Green's theorem in the plane, evaluate the line integral

$$\oint_C (y^2 dx - x^2 dy).$$

Green's theorem in the plane states that

$$\oint_C (f dx + g dy) = \iint_R \left(\frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dx dy,$$


where C is the counter-clockwise boundary of the region R .

PLEASE TURN OVER

SECTION B

[II(3)E 2004]

11. In a certain electrical subsystem three components are arranged as shown below: c1 and c2 are in series, and c3 is in parallel with them, and the components function independently. The probability of failure of component c1 is p_1 , that of c2 is p_2 , and that of c3 is p_3 . Calculate the probability of failure of the system.



The cost of the system with just c1 and c2 is k , the additional cost of installing c3 is l , and the cost of a system failure is m . Show that installation of c3 is justified in terms of expected cost if l/m is less than a certain function of p_1 , p_2 and p_3 .

12. The table below shows the bivariate probability distribution of two random variables, X_1 and X_2 .

	$X_1 = 1$	2	3
$X_2 = 1$	0.12	0.06	0.22
2	0.05	0.02	0.13
3	0.13	0.02	0.25

- Calculate the marginal distributions of X_1 and X_2 .
- Calculate the conditional distribution of X_1 given $X_2 = 3$.
- Compute $E(X_1)$, $E(X_2)$, $\text{var}(X_1)$, $\text{var}(X_2)$, $E(X_1X_2)$, and $\text{cov}(X_1, X_2)$.
- Are X_1 and X_2 correlated? Are they independent? Give your reasoning.

END OF PAPER

MATHEMATICAL FORMULAE

1. VECTOR ALGEBRA

$$\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k} = (a_1, a_2, a_3)$$

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$

Scalar (dot) product:

Vector (cross) product:

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

Scalar triple product:

$$[\mathbf{a}, \mathbf{b}, \mathbf{c}] = \mathbf{a} \cdot \mathbf{b} \times \mathbf{c} = \mathbf{b} \cdot \mathbf{c} \times \mathbf{a} = \mathbf{c} \cdot \mathbf{a} \times \mathbf{b} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$\text{Vector triple product:} \quad \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{c} \cdot \mathbf{a})\mathbf{b} - (\mathbf{b} \cdot \mathbf{a})\mathbf{c}$$

2. SERIES

$$(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!}x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}x^3 + \dots \quad (\alpha \text{ arbitrary, } |x| < 1)$$

$$e^x = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} + \dots,$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^n \frac{x^{2n}}{(2n)!} + \dots,$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \dots,$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^n \frac{x^{n+1}}{(n+1)} + \dots \quad (-1 < x \leq 1)$$

3. TRIGONOMETRIC IDENTITIES AND HYPERBOLIC FUNCTIONS

$$\sin(a+b) = \sin a \cos b + \cos a \sin b;$$

$$\cos(a+b) = \cos a \cos b - \sin a \sin b.$$

$$\cos iz = \cosh z; \quad \cosh iz = \cos z; \quad \sin iz = i \sinh z; \quad \sinh iz = i \sin z.$$

4. DIFFERENTIAL CALCULUS

(a) Leibniz's formula:

$$D^n(fg) = f D^n g + \binom{n}{1} Df D^{n-1} g + \dots + \binom{n}{r} D^r f D^{n-r} g + \dots + D^n f g.$$

(b) Taylor's expansion of $f(x)$ about $x = a$:

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!}f''(a) + \dots + \frac{h^n}{n!}f^{(n)}(a) + \epsilon_n(h),$$

$$\text{where } \epsilon_n(h) = h^{n+1}f^{(n+1)}(a + \theta h)/(n+1)!, \quad 0 < \theta < 1.$$

(c) Taylor's expansion of $f(x, y)$ about (a, b) :

$$f(a+h, b+k) = f(a, b) + [hf_x + kf_y]_{a,b} + \frac{1}{2!} [h^2 f_{xx} + 2hk f_{xy} + k^2 f_{yy}]_{a,b} + \dots$$

(d) Partial differentiation of $f(x, y)$:

$$\text{i. If } y = y(x), \text{ then } f = F(x), \text{ and } \frac{dF}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx}.$$

$$\text{ii. If } x = x(t), y = y(t), \text{ then } f = F(t), \text{ and } \frac{dF}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

$$\text{iii. If } x = x(u, v), y = y(u, v), \text{ then } f = F(u, v), \text{ and}$$

$$\frac{\partial F}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}, \quad \frac{\partial F}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}.$$

(e) Stationary points of $f(x, y)$ occur where $f_x = 0, f_y = 0$ simultaneously.

Let (a, b) be a stationary point: examine $D = [f_{xx}f_{yy} - (f_{xy})^2]_{a,b}$.

If $D > 0$ and $f_{xx}(a, b) < 0$, then (a, b) is a maximum;

If $D > 0$ and $f_{xx}(a, b) > 0$, then (a, b) is a minimum;

If $D < 0$ then (a, b) is a saddle-point.

(f) Differential equations:

i. The first order linear equation $dy/dx + P(x)y = Q(x)$ has an integrating

$$\text{factor } I(x) = \exp\left[\int P(x)dx\right], \text{ so that } \frac{d}{dx}(Iy) = IQ.$$

ii. $P(x, y)dx + Q(x, y)dy = 0$ is exact if $\partial Q/\partial x = \partial P/\partial y$.

5. INTEGRAL CALCULUS

(a) An important substitution: $\tan(\theta/2) = t$:
 $\sin \theta = 2t/(1+t^2)$, $\cos \theta = (1-t^2)/(1+t^2)$, $d\theta = 2dt/(1+t^2)$.

(b) Some indefinite integrals:

$$\int (a^2 - x^2)^{-1/2} dx = \sin^{-1} \left(\frac{x}{a} \right), \quad |x| < a.$$

$$\int (a^2 + x^2)^{-1/2} dx = \sinh^{-1} \left(\frac{x}{a} \right) = \ln \left\{ \frac{x}{a} + \left(1 + \frac{x^2}{a^2} \right)^{1/2} \right\}.$$

$$\int (x^2 - a^2)^{-1/2} dx = \cosh^{-1} \left(\frac{x}{a} \right) = \ln \left| \frac{x}{a} + \left(\frac{x^2}{a^2} - 1 \right)^{1/2} \right|.$$

$$\int (a^2 + x^2)^{-1} dx = \left(\frac{1}{a} \right) \tan^{-1} \left(\frac{x}{a} \right).$$

6. NUMERICAL METHODS

(a) Approximate solution of an algebraic equation:

If a root of $f(x) = 0$ occurs near $x = a$, take $x_0 = a$ and
 $x_{n+1} = x_n - [f(x_n)/f'(x_n)]$, $n = 0, 1, 2 \dots$

(Newton Raphson method).

(b) Formulae for numerical integration: Write $x_n = x_0 + nh$, $y_n = y(x_n)$.

- i. Trapezium rule (1-strip): $\int_{x_0}^{x_1} y(x) dx \approx (h/2) [y_0 + y_1]$.
- ii. Simpson's rule (2-strip): $\int_{x_0}^{x_2} y(x) dx \approx (h/3) [y_0 + 4y_1 + y_2]$.

(c) Richardson's extrapolation method: Let $I = \int_a^b f(x) dx$ and let I_1, I_2 be two estimates of I obtained by using Simpson's rule with intervals h and $h/2$.

Then, provided h is small enough,

$$I_2 + (I_2 - I_1)/15,$$

is a better estimate of I .

7. LAPLACE TRANSFORMS

Function	Transform	Function	Transform
$f(t)$	$F(s) = \int_0^\infty e^{-st} f(t) dt$	$af(t) + bg(t)$	$aF(s) + bG(s)$
df/dt	$sF(s) - f(0)$	$d^2 f/dt^2$	$s^2 F(s) - sf(0) - f'(0)$
$e^{at} f(t)$	$F(s-a)$	$tf(t)$	$-dF(s)/ds$
$(\partial/\partial \alpha) f(t, \alpha)$	$(\partial/\partial \alpha) F(s, \alpha)$	$\int_0^t f(t) dt$	$F(s)/s$
$\int_0^t f(u) g(t-u) du$	$F(s)G(s)$		
1	$1/s$	$t^n (n = 1, 2 \dots)$	$n!/s^{n+1}, (s > 0)$
e^{at}	$1/(s-a), (s > a)$	$\sin \omega t$	$\omega/(s^2 + \omega^2), (s > 0)$
$\cos \omega t$	$s/(s^2 + \omega^2), (s > 0)$	$H(t-T) = \begin{cases} 0, & t < T \\ 1, & t > T \end{cases}$	$e^{-sT}/s, (s, T > 0)$

8. FOURIER SERIES

If $f(x)$ is periodic of period $2L$, then $f(x+2L) = f(x)$, and

$$f(x) = \frac{1}{2} a_0 + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{L} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}, \text{ where}$$

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx, \quad n = 0, 1, 2, \dots, \text{ and}$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx, \quad n = 1, 2, 3, \dots$$

Parseval's theorem

$$\frac{1}{L} \int_{-L}^L |f(x)|^2 dx = \frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2).$$

EE
2nd yr
Paper 3
2004

MATHEMATICS FOR ENGINEERING STUDENTS

EXAMINATION QUESTION / SOLUTION

2003 - 2004

Please write on this side only, legibly and neatly, between the margins

PAPER

II (3)

QUESTION

SOLUTION
/E1



$$w = u + iv = \frac{z}{z-1} \quad \text{or} \quad w-1 = \frac{1}{z-1}$$

$$\therefore u-1 + iv = \frac{1}{(x-1) + iy} = \frac{x-1-iy}{(x-1)^2 + y^2}$$

$$\therefore (u-1) = \frac{x-1}{(x-1)^2 + y^2} \quad v = -\frac{y}{(x-1)^2 + y^2} \quad \text{--- } (*)$$

$$1) \quad (u-1)^2 + v^2 = \frac{1}{(x-1)^2 + y^2}$$

Circles $(x-1)^2 + y^2 = a^2$ maps to $(u-1)^2 + v^2 = \frac{1}{a^2}$.



Fixed p/s $w=2$ are $w=2 \Rightarrow 0, 2$. When $a > 1$ the pictures are reversed.

$$2) \quad x=0 \text{ is the } y\text{-axis: } w \quad (*)$$

$$u-1 = -\frac{1}{1+y^2} \quad v = -\frac{y}{1+y^2}$$

$$v^2 + (u-1)^2 = \frac{1}{1+y^2} = -(u-1)$$

$$\therefore (u-1 + \frac{1}{2})^2 + v^2 = \frac{1}{4} \quad \text{Completing the square.}$$

$$\text{or } (u - \frac{1}{2})^2 + v^2 = (\frac{1}{2})^2 \quad \text{Circle centred at } (\frac{1}{2}, 0) \text{ radius } \frac{1}{2}.$$

$$3) \quad y = m(x-1): \quad w \quad (*)$$

$$u-1 = \frac{1}{m^2+1} \cdot \frac{1}{x-1}, \quad v = -\frac{m}{m^2+1} \cdot \frac{1}{x-1}$$

$$\therefore m(u-1) + v = 0$$


Setter: J.D. GIBSON

Checker: F.L. J

Setter's signature: J.D. Gibson

Checker's signature:

Please write on this side only, legibly and neatly, between the margins



$$H_R = e^{i\theta} z \quad z = R e^{i\theta}$$

$$0: 0 \rightarrow \pi$$

$$f(z) = \frac{e^{2iz}}{(z^2+4)(z^2+9)} \quad \text{has 4 poles at} \quad \begin{matrix} z = \pm 2i \\ z = \pm 3i \end{matrix}$$

Choose only those in upper $\frac{1}{2}$ -plane.

$$\begin{aligned} \text{Residue of } f(z) \text{ at } z = 2i & \quad \frac{e^{-4}}{5 \times 4i} \\ \text{Residue of } f(z) \text{ at } z = 3i & \quad \frac{e^{-6}}{-5 \times 6i} \end{aligned}$$

RESIDUE THM:

$$\oint_C f(z) dz = 2\pi i \left\{ \frac{e^{-4}}{20i} - \frac{e^{-6}}{30i} \right\} = \frac{\pi}{5} \left(\frac{e^{-4}}{2} - \frac{e^{-6}}{3} \right)$$

$$\text{Now} \quad \int_C f(z) dz = \int_{H_R} f(z) dz + \int_{-R}^R \frac{e^{2ix}}{(x^2+4)(x^2+9)} dx$$

$$\lim_{R \rightarrow \infty} \int_{H_R} f(z) dz = 0 \quad \text{by Jordan's Lemma}$$

i) Only poles in upper $\frac{1}{2}$ -plane

ii) $n=2 > 0$

iii) $\frac{1}{(z^2+4)(z^2+9)} \rightarrow 0 \quad R \rightarrow \infty$

In the limit $R \rightarrow \infty$,

$$\frac{\pi}{5} \left(\frac{e^{-4}}{2} - \frac{e^{-6}}{3} \right) = \int_{-\infty}^{\infty} \frac{\cos 2x + i \sin 2x}{(x^2+4)(x^2+9)} dx$$

Since part of integral zero as it is odd.

$$\int_{-\infty}^{\infty} \frac{\cos 2x}{(x^2+4)(x^2+9)} dx = \frac{\pi}{5} \left(\frac{e^{-4}}{2} - \frac{e^{-6}}{3} \right)$$

Setter: J. D. Gibson


Setter's signature: J. D. Gibson

Checker: X. Wu

Checker's signature:

Xuesong

Please write on this side only, legibly and neatly, between the margins



$$\begin{aligned} H_R: & \theta: 0 \rightarrow \pi \\ C: & H_r: \theta: \pi \rightarrow 0 \\ x \text{ axis: } & (-R \rightarrow -r) \\ & (r \rightarrow R) \end{aligned}$$

$f(z) = \frac{e^{iz}}{z}$ has one simple pole at $z=0$.

The contour has been specifically chosen to exclude pole at $z=0$

Since C excludes the pole, $f(z)$ analytic everywhere in C .

$$\oint_C \frac{e^{iz}}{z} dz = 0 \quad \text{Cauchy's Thm.}$$

$$0 = \oint_C \frac{e^{iz}}{z} dz = \int_{H_R} \frac{e^{iz}}{z} dz + \int_{h_r} \frac{e^{iz}}{z} dz + \left(\int_{-R}^{-r} + \int_r^R \right) \frac{e^{ix}}{x} dx$$

$$H_r: \text{circle } z = r e^{i\theta} \quad \theta: \pi \rightarrow 0 \quad \int_{H_r} = \int_{\pi}^0 \frac{e^{i r e^{i\theta}}}{r e^{i\theta}} i r e^{i\theta} d\theta$$

$$\therefore \lim_{r \rightarrow 0} \int_{H_r} = -i \int_0^{\pi} d\theta = -i\pi$$

$H_R: \text{circle } z = R e^{i\theta} \quad \theta: 0 \rightarrow \pi$. Jordan's Lemma says that

$$\lim_{R \rightarrow \infty} \int_{H_R} \frac{e^{iz}}{z} dz = 0 \quad \text{provided } \begin{aligned} & \text{i) Only sing of } f(z) \text{ are poles} \\ & \text{ii) } m (=1) > 0 \quad \checkmark \\ & \text{iii) } \frac{1}{|z|} \rightarrow 0 \text{ as } R \rightarrow \infty \quad \checkmark. \end{aligned}$$

$$\therefore \lim_{\substack{r \rightarrow 0 \\ R \rightarrow \infty}} \left(\int_{-R}^{-r} + \int_r^R \right) \frac{e^{ix}}{x} dx = i\pi$$

$$\text{or} \quad \int_{-\infty}^{\infty} \frac{e^{ix}}{x} dx = i\pi$$

$$\int_{-\infty}^{\infty} \frac{\cos x}{x} dx = 0 \quad \text{as } \frac{\cos x}{x} \text{ odd.}$$

$$\therefore \int_{-\infty}^{\infty} \frac{\sin x}{x} dx = \pi$$

Setter : J.D. GIBSON

Setter's signature : J.D. Gibson

Checker : X.WU

Checker's signature :

Xuesong

Please write on this side only, legibly and neatly, between the margins

$$\begin{aligned}
 1) \quad \bar{\pi}(\omega) &= \int_{-\infty}^{\infty} e^{-i\omega t} \pi(t) dt \\
 &= \int_{-1/2}^{1/2} e^{-i\omega t} dt = \frac{1}{-i\omega} [e^{-i\omega/2} - e^{i\omega/2}] \\
 &= \frac{-2}{\omega} (e^{-i\omega/2} - e^{i\omega/2}) = \sin \omega/2 / \omega/2 = \text{sinc } \omega
 \end{aligned}$$

3

$$\begin{aligned}
 2) \quad \bar{\Lambda}(\omega) &= \int_{-\infty}^{\infty} e^{-i\omega t} \Lambda(t) dt = \int_{-1}^0 (1+t) e^{-i\omega t} dt + \int_0^1 (1-t) e^{-i\omega t} dt \\
 &= \left[\left(\frac{1}{\omega^2} + \frac{t}{\omega} \right) (1 - e^{-i\omega t}) + \frac{i}{\omega} e^{-i\omega t} \right]_{-1}^0 - \left[\frac{t}{\omega} - \frac{1}{\omega^2} (1 - e^{-i\omega t}) \right]_{0}^1 \\
 &= \frac{2}{\omega^2} - \frac{1}{\omega^2} (e^{i\omega} + e^{-i\omega}) = \frac{2}{\omega^2} (1 - \cos \omega) \\
 &= \frac{4}{\omega^2} \sin^2 \frac{1}{2} \omega = \text{sinc}^2 \omega
 \end{aligned}$$

6

[Comes from $\int t e^{-i\omega t} dt = \left[\frac{1}{\omega^2} + \frac{it}{\omega} \right] e^{-i\omega t}]$

$$3) \quad \int_{-\infty}^{\infty} \frac{e^{ipx}}{x} dx = \int_{-p}^p \frac{e^{iq}}{q} dq = \begin{cases} i\pi & p > 0 \\ -i\pi & p < 0 \end{cases}$$

$$\therefore \int_{-\infty}^{\infty} \frac{e^{-i\omega t} \sin t/2}{t/2} dt = 2 \int_{-\infty}^{\infty} \frac{e^{i\theta}}{\theta} e^{-2i\omega\theta} d\theta \quad t/2 = \theta$$

3

$$= \frac{1}{i} \int_{-\infty}^{\infty} \frac{e^{i\theta p_1}}{\theta} d\theta - \frac{1}{i} \int_{-\infty}^{\infty} \frac{e^{i\theta p_2}}{\theta} d\theta \quad \begin{matrix} p_1 = 1-2\omega \\ p_2 = -1-2\omega \end{matrix}$$

$$= \begin{cases} 0 & p_1, p_2 \text{ same sign; } \omega < -1/2, \omega > 1/2 \\ & (p_1, p_2 > 0) \quad (p_1 < 0, p_2 < 0) \\ \frac{2\pi i}{i} = 2\pi & p_1 > 0, p_2 < 0 \text{ (opposite signs)} \quad -1/2 < \omega < 1/2 \\ & \text{(The case } p_1 < 0, p_2 > 0 \text{ is not possible)} \end{cases}$$

3

$$= 2\pi \begin{cases} 1 & -1/2 < \omega < 1/2 \\ 0 & \text{otherwise} \end{cases} = 2\pi \pi(\omega)$$

Setter : J.D. GIBSON

Setter's signature : J.D. Gibson

Checker : X. WU

Checker's signature :

Xuesong

Please write on this side only, legibly and neatly, between the margins

Consider $\bar{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \bar{f}(\omega) e^{i\omega t} d\omega$$

$$\therefore f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} f(t') e^{-i\omega t'} dt' \right) e^{i\omega t} d\omega$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} e^{i\omega(t-t')} d\omega \right) f(t') dt'$$

Now the δ -function has the property:

$$f(t) = \int_{-\infty}^{\infty} \delta(t-t') f(t') dt'$$

$$\text{so } \textcircled{*} \quad 2\pi \delta(t-t') = \int_{-\infty}^{\infty} e^{i\omega(t-t')} d\omega$$

Signs irrelevant

Consider $e^{+it(\omega-\omega')}$
if $\bar{f}(\omega) \leftrightarrow f(t)$
reversed.

Bookwork

$$\therefore \int_{-\infty}^{\infty} f(t) g^*(t) dt = \int_{-\infty}^{\infty} dt \left(\int_{-\infty}^{\infty} \bar{f}(\omega) e^{i\omega t} d\omega \right) \left(\int_{-\infty}^{\infty} \bar{g}(\omega') e^{i\omega' t} d\omega' \right)^* \frac{1}{4\pi^2}$$

$$= \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} e^{i(\omega-\omega')t} dt \right) \bar{f}(\omega) \bar{g}^*(\omega') d\omega d\omega'$$

$$= \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} 2\pi \delta(\omega-\omega') \bar{f}(\omega) \bar{g}^*(\omega') d\omega d\omega' \quad \text{Using } \textcircled{*} \text{ above}$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} \bar{f}(\omega) \bar{g}^*(\omega) d\omega$$

If $f(t) = e^{-|t|} = \begin{cases} e^{-t} & t > 0 \\ e^t & t < 0 \end{cases} \Rightarrow \bar{f}(\omega) = \int_0^{\infty} e^{-t-i\omega t} dt + \int_{-\infty}^0 e^{t-i\omega t} dt$

$$\therefore \bar{f}(\omega) = \frac{1}{1+i\omega} + \frac{1}{1-i\omega} = \frac{2}{1+\omega^2}$$

Unseen

$$\bar{g}(\omega) = \frac{2\pi}{2} [\delta(\omega-\Omega) + \delta(\omega+\Omega)]$$

$$\therefore \int_{-\infty}^{\infty} e^{-|t|} \cos \Omega t dt = \frac{2\pi}{2\pi} \int_{-\infty}^{\infty} \frac{2}{1+\omega^2} [\delta(\omega-\Omega) + \delta(\omega+\Omega)] d\omega$$

$$= \frac{2}{1+\Omega^2} \quad \text{or directly from the } t\text{-integral.}$$

Setter : J.D. GIBSON

Setter's signature : J.D. Gibson

Checker :

X. WU

Checker's signature :

Xuesong

Please write on this side only, legibly and neatly, between the margins

$$\ddot{y} + 5\dot{y} + 6y = f(t)$$

$$\mathcal{L} \dot{y} = s \bar{y}(s) - y(0)$$

$$\mathcal{L} \ddot{y} = s^2 \bar{y}(s) - s y(0) - \dot{y}(0)$$

Take,

$$\therefore s^2 \bar{y}(s) - s y(0) - \dot{y}(0) + 5s \bar{y}(s) - 5y(0) + 6 \bar{y}(s) = \bar{f}(s)$$

$$(s^2 + 5s + 6) \bar{y}(s) = (s+5) \bar{y}(0) + \dot{y}(0) + \bar{f}(s)$$

$$= (s+3) \bar{y}(0) + \bar{f}(s) \quad \text{if } \dot{y}(0) = -2y(0)$$

$$\therefore \bar{y}(s) = \frac{\bar{f}(s)}{(s+3)(s+2)} + \frac{\bar{y}(0)}{s+2}$$

Now inverse transform: shift theorem $\mathcal{L}^{-1}\left(\frac{1}{s+2}\right) = e^{-2t}$

$$\frac{\bar{f}(s)}{(s+3)(s+2)} = \frac{\bar{f}(s)}{s+2} - \frac{\bar{f}(s)}{s+3}$$

$$\therefore y(t) = y(0) e^{-2t} + \mathcal{L}^{-1}\left(\frac{\bar{f}(s)}{s+2}\right) - \mathcal{L}^{-1}\left(\frac{\bar{f}(s)}{s+3}\right)$$

Now the convolution theorem says that

$$\mathcal{L}^{-1}(\bar{f}(s) \bar{g}(s)) = f(t) * g(t)$$

$$g_1(s) = \frac{1}{s+2}$$

$$g_2(s) = \frac{1}{s+3}$$

$$g_1(t) = e^{-2t} \quad g_2(t) = e^{-3t}$$

$$\therefore f(t) * g_1(t) = \int_0^t e^{-2(t-u)} f(u) du$$

$$f(t) * g_2(t) = \int_0^t e^{-3(t-u)} f(u) du$$

Altogether; $\bar{y}(s) = y(0) e^{-2t} + \int_0^t [e^{-2(t-u)} - e^{-3(t-u)}] f(u) du$

Setter : J.D. Gibson

Setter's signature : J.D. Gibson

Checker :

X.WU


Checker's signature :

Xuesong

Please write on this side only, legibly and neatly, between the margins

Consider a curve C with unit
tangent vector \hat{T} & normal \hat{n}

$$\hat{T} = \frac{dr}{ds} = \hat{i} \left(\frac{dx}{ds} \right) + \hat{j} \left(\frac{dy}{ds} \right)$$



but, because $\hat{n} \cdot \hat{T} = 0 \Rightarrow \hat{n} = \pm \left(\hat{i} \frac{dy}{ds} - \hat{j} \frac{dx}{ds} \right)$

Choose a vector \underline{u} such that $\underline{u} = (Q\hat{i} - P\hat{j})$

$$\therefore \underline{u} \cdot \hat{n} = \left(P \frac{dx}{ds} + Q \frac{dy}{ds} \right) \Rightarrow P dx + Q dy = \underline{u} \cdot \hat{n} ds$$

and $\text{div } \underline{u} = Q_x - P_y$. Hence, Green's Theorem gives

$$\oint_C \underline{u} \cdot \hat{n} ds = \iint_R \text{div } \underline{u} dx dy \quad \text{if } C \text{ is closed.}$$

$$\underline{u} = x^2 + y^2 \Rightarrow \text{div } \underline{u} = 2(x+y)$$

$$\therefore \iint_R \text{div } \underline{u} dx dy = 2 \iint_R (x+y) dx dy$$

$$= 2 \int_0^2 \left\{ \int_{\frac{1}{2}x^2}^{\sqrt{2}x} (x+y) dy \right\} dx$$


$$= 2 \int_0^2 \left(xy + \frac{1}{2}y^2 \right)_{\frac{1}{2}x^2}^{\sqrt{2}x} dx$$

$$= 2 \int_0^2 \left(\sqrt{2}x^{3/2} + x - \frac{1}{2}x^3 - \frac{1}{8}x^4 \right) dx$$

$$= 2 \left[\sqrt{2} \cdot \frac{2}{5} \cdot x^{5/2} + \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{40}x^5 \right]_0^2$$

$$= 2 \left[2^{4/5} + 2 - 2 - \frac{3^2}{40} \right]$$

$$= \frac{8}{5} [4 - 1] = \frac{24}{5}$$



Setter : B. D. GIBSON

Setter's signature : J. D. Gibson

Checker :

Checker's signature :


Please write on this side only, legibly and neatly, between the margins

(i) Transformation of boundaries:


$$\bullet \quad x+y=1 \Rightarrow v=1$$

$$\bullet \quad x=0 \quad (u=-y^2, v=y) \Rightarrow u=-v^2$$

$$\bullet \quad y=0 \quad (u=x^2, v=x) \Rightarrow u=v^2$$



\Rightarrow



$$\text{Jacobian: } \frac{\partial(u,v)}{\partial(x,y)} = \begin{vmatrix} 2x & -2y \\ 1 & 1 \end{vmatrix} = 2(x+y) = 2v, \quad \frac{\partial(x,y)}{\partial(u,v)} = \frac{1}{2v}$$

$$I = \iint_{R'} v^n f(u) \frac{1}{2v} du dv = \frac{1}{2} \int_0^1 v^{n-1} \left\{ \int_{-v^2}^{v^2} f(u) du \right\} dv$$

(ii) For $n=2$, $f(u) = e^u$

$$I = \frac{1}{2} \int_0^1 v \left\{ \int_{-v^2}^{v^2} e^u du \right\} dv = \frac{1}{2} \int_0^1 v (e^{v^2} - e^{-v^2}) dv$$

$$= \frac{1}{4} (e^{v^2} + e^{-v^2}) \Big|_0^1 = \frac{1}{4} (e + e^{-1} - 2)$$

(iii) For $n=0$, $f \equiv 1$

$$I = \frac{1}{2} \int_0^1 v^{-1} \left\{ \int_{-v^2}^{v^2} du \right\} dv = \int_0^1 v dv = \frac{1}{2}$$

$$\therefore I = \frac{1}{2}$$

$$\text{In this case } I = \iint_R dxdy = \underline{\text{Area of } R} = \frac{1}{2} \times 1 \times 1 = \frac{1}{2}$$

(15)

Setter : X.WU

Setter's signature :

Checker : R.L.JACOBS

Checker's signature :

Xuesong
R.L.JACOBS

Please write on this side only, legibly and neatly, between the margins

$$(i) \operatorname{div} \underline{F} = \underline{\nabla} \cdot \underline{F} = 2 \sin z + ze^y - x^2 \sin z = (2 - x^2) \sin z + ze^y$$

$$\operatorname{curl} \underline{F} = \underline{\nabla} \times \underline{F} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2x \sin z & ze^y & x^2 \cos z + ae^y \end{vmatrix}$$

$$= (a-1)e^y \underline{i} - (2x \cos z - 2x \cos z) \underline{j} + 0 \underline{k}$$

$$\operatorname{curl} \underline{F} = (a-1)e^y \underline{i}$$

$$(ii) a = 1. \quad \frac{\partial \phi}{\partial x} = 2x \sin z \Rightarrow \phi = x^2 \sin z + f(y, z)$$

$$\frac{\partial \phi}{\partial y} = ze^y \Rightarrow \frac{\partial f}{\partial y} = ze^y, \quad f = ze^y + g(z)$$

$$\therefore \phi = x^2 \sin z + ze^y + g(z)$$

$$\frac{\partial \phi}{\partial z} = x^2 \cos z + e^y \Rightarrow x^2 \cos z + e^y + g'(z) = x^2 \cos z + e^y$$

$$\text{i.e. } g'(z) = 0, \quad g = \text{const.}$$

$$\therefore \underline{\phi} = x^2 \sin z + ze^y + C$$

$$(iii) (\underline{F} \cdot \underline{\nabla}) \phi = 2x \sin z \frac{\partial \phi}{\partial x} + ze^y \frac{\partial \phi}{\partial y} + (x^2 \cos z + e^y) \frac{\partial \phi}{\partial z} \\ = 4x^2 \sin^2 z + z^2 e^{2y} + (x^2 \cos z + e^y)^2$$

$$\nabla^2 \phi = \frac{\partial^2}{\partial x^2} \phi + \frac{\partial^2}{\partial y^2} \phi + \frac{\partial^2}{\partial z^2} \phi = 2 \sin z + ze^y - x^2 \sin z$$

$$\therefore \nabla^2 \phi = (2 - x^2) \sin z + ze^y \quad (= \underline{\nabla} \cdot \underline{F})$$

Setter : X. WU

Setter's signature :

Checker : R. L. JACOBS

Checker's signature :

Xuesong
R. L. Jacobs

Please write on this side only, legibly and neatly, between the margins

(i) Consider two paths C_1 and C_2 with same endpoints A and B . Consider composite path

$C = C_1 \bar{C}_2$ taken counter clockwise



Then
$$\oint_C (F_1 dx + F_2 dy) = \iint_R \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) dx dy$$

In this case $F_1 = y \cos x$
 $F_2 = \sin x + 3y$

$\therefore \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} = 0$

$\therefore \oint_C = 0 \Rightarrow \int_{C_1} (F_1 dx + F_2 dy) = \int_{C_2} (F_1 dx + F_2 dy)$

Hence integral is independent of path.

Now $\frac{\partial q}{\partial x} = y \cos x$, $\frac{\partial q}{\partial y} = \sin x + 3y$
 (1) (2)

(1) $\Rightarrow q = y \sin x + h(y)$. Subst. into (2) to get
 $\sin x + \frac{dh}{dy} = \sin x + 3y \Rightarrow h = \frac{3}{2} y^2 + C$

$\therefore \phi(x, y) = y \sin x + \frac{3}{2} y^2 + C$

$\therefore \int_A^B = \phi \Big|_{(0,0)}^{(\pi/2, 1)} - \phi(0, 0) = 1 + \frac{3}{2} + C - C = 5/2$

Setter :

R-L. Jacob

Setter's signature :

R-L. Jacob


Checker :

J-D. Bishon

Checker's signature :

Please write on this side only, legibly and neatly, between the margins

$$\begin{aligned}
 (ii) \quad \oint_C (y^2 dx - x^2 dy) &= \iint_R (-2x - 2y) dx dy \\
 &= -2 \int_{y=0}^1 \left[\int_{x=0}^{1+2(1-y)^{1/2}} (x+y) dx \right] dy \\
 &= -2 \int_{y=0}^1 \left[2(1-y)^{1/2} + y 2(1-y)^{1/2} \right] dy
 \end{aligned}$$



$$\begin{aligned}
 &= -2 \left[2y - \frac{2y^3}{3} - \frac{2}{3}(1-y)^{3/2} \right]_0^1 \\
 &= -2 \left[2 - \frac{2}{3} - 0 - 0 + 0 + \frac{2}{3} \right] \\
 &= -4
 \end{aligned}$$

Setter : R-L. JACOBS

Setter's signature : R-L. Jacobs

Checker : J. D. Robinson

Checker's signature :

5

EXAMINATION QUESTION / SOLUTION

2003-2004

QUESTION

11

Please write on this side only, legibly and neatly, between the margins

SOLUTION

11

1.

$$\begin{aligned} P(\text{system failure}) &= P\{(c1 \text{ fails} \cup c2 \text{ fails}) \cap (c3 \text{ fails})\} \\ &= \{1 - P(c1 \text{ does not fail} \cap c2 \text{ does not fail})\} P(c3 \text{ fails}) \\ &= \{1 - (1 - p_1)(1 - p_2)\} p_3. \end{aligned}$$

5

$$\begin{aligned} E(\text{cost without } c3) &= k + \{1 - (1 - p_1)(1 - p_2)\}m \\ E(\text{cost with } c3) &= k + l + \{1 - (1 - p_1)(1 - p_2)\}p_3m \end{aligned}$$

5

$$\begin{aligned} \text{difference} &= \{1 - (1 - p_1)(1 - p_2)\}(1 - p_3)m - l \\ &> 0 \text{ if } l/m < \{1 - (1 - p_1)(1 - p_2)\}(1 - p_3) \end{aligned}$$

5

15

Setter: MT CROWDER

Setter's signature: MT Crowder

Checker: R COLEMAN

Checker's signature: R Coleman

EXAMINATION QUESTION / SOLUTION

2003-2004

QUESTION

12

Please write on this side only, legibly and neatly, between the margins

SOLUTION

12

2.

- (i) marginal X_1 : $p_1(1) = 0.3$, $p_1(2) = 0.1$, $p_1(3) = 0.6$
 marginal X_2 : $p_2(1) = 0.4$, $p_2(2) = 0.2$, $p_2(3) = 0.4$

4

- (ii) conditional $X_1 | X_2 = 3$:

$$p_{1|2}(1) = 0.13/0.4 = 0.325,$$

$$p_{1|2}(2) = 0.02/0.4 = 0.05,$$

$$p_{1|2}(3) = 0.25/0.4 = 0.625,$$

4

- (iii) $E(X_1) = \sum x_1 p_1(x_1) = 2.3$, $E(X_2) = 2.0$

$$\text{var}(X_1) = \sum x_1^2 p_1(x_1) - \mu_1^2 = 6.1 - 2.3^2 = 0.81, \text{var}(X_2) = 4.8 - 2.0^2 = 0.8$$

$$E(X_1 X_2) = \sum x_1 x_2 p(x_1, x_2) = 4.62$$

$$\text{cov}(X_1, X_2) = 4.62 - (2.3 \times 2.0) = 0.02$$

5

- (iv) Yes, because covariance not zero.

No, because covariance not zero, or e.g. $p(1, 2) \neq p_1(1)p_2(2)$

2

(15)

Setter: *MJ CROWDER*

Setter's signature: *MJ Crowder*

Checker: *A COLMAN*

Checker's signature: *A Colman*

