Comment on "Convergence Properties of the Likelihood of Computed Dynamic Models": A Corrected Proposition 2

Robert Kirkby*

October 2026

Draft

Abstract

Proposition 2 of Fernández-Villaverde et al. [2006] bounds the difference between the exact and the computed likelihood of a dynamic model by a term linear in the sample size. Ackerberg et al. [2009] showed by counterexample that the proposition is false, and argued that a corrected bound on likelihood levels would be uninformative about estimation; they did not locate the error in the proof or state a corrected result for the dynamic setting. We locate the error, a single step in which a difference of two T-fold products of densities is bounded as if the other factors were absent, and show that the proposition holds, with the same form of bound, for the logarithm of the likelihood: $|\log L_j(y^T) - \log L(y^T)| \le (T\chi/\xi + K/(1-\alpha))\,\delta$ under a pathwise version of the paper's contraction condition. The dependence on T is sharp. The average log likelihood is therefore within $O(\delta)$ of its exact value uniformly in the sample size, so the effect of a fixed approximation error on point estimates does not grow with T; relative to sampling uncertainty it matters once $\sqrt{T}\delta$ is not small. FRS's Bayesian corollaries hold with explicit, if conservative, constants.

Keywords: Approximation error, likelihood, computed dynamic models, Markov chains, coupling.

1 Introduction

Likelihood-based estimation of a dynamic model uses an approximated policy function, so the researcher evaluates the likelihood L_j of the computed model rather than the likelihood L of the exact one. Fernández-Villaverde et al. [2006] (henceforth, FRS) asked how the two are related, in particular how numerical error incurred in solving the dynamic model affects likelihood-based estimation of that model. Their Proposition 1 shows that $L_j \to L$ for a fixed sample as the computed transition and measurement equations converge to the exact

*Victoria University of Wellington. Every result in this comment has been verified in the Lean 4 proof assistant with the Mathlib library; Appendix C lists the correspondence, and the code is at `https://github.com/LeanEconomics/LeanEconomics`. The formal development and a first draft of this comment were produced with the assistance of Claude (Anthropic), an AI model.

ones. Their Proposition 2 quantifies the speed: if the computed equations are within δ of the exact ones in the sup norm, then $|L_j(y^T) - L(y^T)| < (TB\chi + L_F/(1-\alpha))\delta$, with constants independent of the sample size T.[1] FRS drew three conclusions from it, emphasised in their abstract and introduction: that "second order approximation errors in the policy function ... have first order effects on the likelihood function"; that the error "gets compounded with the size of the sample", so the accuracy of the solution must improve as T grows if the computed likelihood is to stay close to the exact one; and hence that linearisation, which fixes δ, "is due to fail as the sample size grows". They also used the proposition to motivate a likelihood-ratio diagnostic for solution accuracy.

Ackerberg et al. [2009] (henceforth, AGH), following an unpublished note by Geweke [2007], showed that Proposition 2 is false. In the static Gaussian location model $y_t = \gamma + v_t$, approximated by $y_t = \gamma + \delta + v_t$, they computed a sharp bound on $|L_j - L|$ of order $(\sigma\sqrt{2\pi})^{-(T-1)}$. This grows exponentially in T when $\sigma < 1/\sqrt{2\pi}$ and vanishes exponentially when $\sigma > 1/\sqrt{2\pi}$, so FRS's linear bound is too small in one case and too large in the other. Their contribution on Proposition 2 is this counterexample, together with the argument that bounds on likelihood *levels* are the wrong object for inference, since in their example the effect of δ on the maximum likelihood estimator depends on neither T nor σ. They then proved new results for a static i.i.d. model, under a direct assumption that the log densities and their first two derivatives are within Δ_j of the exact ones: the pseudo-maximum-likelihood estimator has probability limit within $O(\Delta_j)$ of the truth, and is asymptotically normal if $\sqrt{T}\Delta_j \to 0$. AGH did not identify which step of FRS's proof fails and did not state a corrected version of Proposition 2; they remarked that seeking a valid bound on likelihoods in more general models "might not be a fruitful endeavor". Their constructive results are for the static model, and they take Δ_j as given rather than deriving it from the policy-function error δ and the dynamics of the state.

This comment locates the step of FRS's proof that fails and proves a corrected Corrected Proposition 2′ that keeps their intent. The correction is to bound the logarithm of the likelihood rather than its level. That is the scale on which likelihoods are used in estimation, and it turns FRS's bound on the absolute error into a bound on the relative error $|L_j/L - 1|$. The bound keeps FRS's form: under a pathwise version of their contraction condition, the log-likelihood error is at most $(T\chi/\xi + K/(1-\alpha))\delta$, linear in T, and this rate cannot be improved. FRS's three conclusions then hold in modified form. First, an approximation error of order δ in the policy function moves the log likelihood by $O(T\delta)$, but the average log likelihood by only $O(\delta)$, uniformly in T; this is AGH's "second order errors have second order effects", now derived in FRS's dynamic setting rather than assumed. Second, the effect on point estimates does not compound with the sample size. Sampling uncertainty, however, shrinks like $1/\sqrt{T}$, so approximation error matters for inference, classical or Bayesian, once $\sqrt{T}\delta$ is not small. Third, linearisation, which fixes δ, does not fail as the sample grows in the sense FRS claimed, but its bias does grow relative to standard errors. Their likelihood-ratio diagnostic survives: the log likelihood ratio between two approximations differs by $O(T\delta)$,

[1] Here L_F is a Lipschitz constant of an integrated conditional likelihood (Section 2). FRS denote it L, the same symbol as the likelihood; we write L_F throughout to avoid the clash.

against sampling noise of order $\sqrt{T}$, so the diagnostic detects approximation error in roughly the regime where it matters.

Section 2 locates the error: step (14) of FRS's appendix bounds a difference of two T-fold products of densities by $TB\chi\delta$, but the telescoping identity places $T-1$ density factors beside each per-period error, so B is really the $(T-1)$-th power of a bound on the density. It also gives a counterexample that, unlike AGH's, satisfies FRS's compact-support assumption. Section 3 states and proves Corrected Proposition 2′. Section 4 draws the consequences for estimation and for Bayesian inference, where FRS's corollaries hold with explicit constants. Appendix A derives the assumptions of the corrected proposition from primitives in the case where the measurement shock alone is recovered from the data.

2 The setting and the error in the proof

FRS consider the state-space system

$$S_t = \varphi(S_{t-1}, W_t; \gamma), \qquad Y_t = g(S_t, V_t; \gamma),$$

with compact supports for the state and the i.i.d. shocks, and a unique invariant distribution μ^* of the state. The shock is partitioned as $W_t = (W_{1t}, W_{2t})$, where (W_{2t}, V_t) is recovered from the data by inverting the system (FRS's Assumption 3). Here T is the number of observations in the data $y^T = (y_1, \ldots, y_T)$, and the likelihoods below are the exact integrals implied by each model, not simulation estimates of them; the error from evaluating them by simulation, for example with a particle filter, is a separate matter that neither FRS nor this comment addresses. We fix γ and the data y^T and suppress both. Write $\omega = W_1^T$ for the unobserved shock path, with law Q, and

$$p_t(s, \omega) = p\big(y_t \mid W_1^t = \omega_{\le t},\, S_0 = s,\, y^{t-1}\big)$$

for the conditional density of the t-th observation, with $p_{j,t}$ (FRS's $p_j(y_t \mid W_1^t, S_0, y^{t-1})$) its counterpart under the computed equations (φ_j, g_j), whose invariant distribution is μ_j^*. The two likelihoods are[2]

$$L(y^T) = \int_S \int_\Omega \prod_{t=1}^{T} p_t(s,\omega)\, Q(\mathrm{d}\omega)\, \mu^*(\mathrm{d}s), \qquad L_j(y^T) = \int_S \int_\Omega \prod_{t=1}^{T} p_{j,t}(s,\omega)\, Q(\mathrm{d}\omega)\, \mu_j^*(\mathrm{d}s).$$

FRS's Assumptions 4 and 8 bound the densities below by $\xi > 0$; their Lemma 6 gives $|p_{j,t} - p_t| \le \chi\delta$ when $\|\varphi_j - \varphi\| \le \delta$ and $\|g_j - g\| \le \delta$; and their Condition 1 is the mean contraction $\int \|\varphi(s,w) - \varphi(s',w)\|\, Q(\mathrm{d}w) \le \alpha\|s - s'\|$ with $\alpha < 1$.

FRS's Proposition 2. *Under Assumptions 1–11 and Condition 1, if $\|\varphi_j - \varphi\| \le \delta$ and $\|g_j - g\| \le \delta$, there are positive constants B and L_F such that, for all T, $|L_j(y^T) - L(y^T)| <$*

[2] FRS's displays write the integral over W_1^T period by period, as a product of integrals. Since p_t depends on the whole history $\omega_{\le t}$, the likelihood is the integral of the product. Nothing below depends on the distinction.

$(TB\chi + L_F/(1-\alpha))\delta$.

The proof writes $F(s) = \int \prod_t p_t(s,\cdot)\,\mathrm{d}Q$ and $F_j(s) = \int \prod_t p_{j,t}(s,\cdot)\,\mathrm{d}Q$ and splits

$$|L_j - L| \le \Big|\int F\,\mathrm{d}\mu_j^* - \int F\,\mathrm{d}\mu^*\Big| + \int |F_j - F|\,\mathrm{d}\mu_j^*.$$

The first term is bounded in FRS's (12) by $L_F\delta/(1-\alpha)$, using a result of Santos and Peralta-Alva [2005] with L_F a Lipschitz constant of F. The second is bounded in their (14) by $TB\chi\delta$, from Lemma 6 and a bound B on the densities. Step (14) is where the proof fails. A difference of two products satisfies the telescoping identity

$$\prod_{t=1}^{T} p_{j,t} - \prod_{t=1}^{T} p_t = \sum_{t=1}^{T} \Big(\prod_{s<t} p_{j,s}\Big)\,(p_{j,t} - p_t)\,\Big(\prod_{s>t} p_s\Big), \tag{1}$$

so each of the T per-period errors is multiplied by $T-1$ densities. With M an upper bound on the densities, the bound that follows is $TM^{T-1}\chi\delta$, and M^{T-1} depends on T. No T-free constant can take its place:

Proposition 1. *Let $M > 1$ and $\varepsilon > 0$. There is no constant B such that $|(M+\varepsilon)^T - M^T| \le TB\varepsilon$ for every $T \in \mathbb{N}$.*

Proof. By Bernoulli's inequality, $(M+\varepsilon)^{n+1} - M^{n+1} = M^{n+1}[(1+\varepsilon/M)^{n+1} - 1] \ge (n+1)\varepsilon M^n$. The bound would therefore give $M^n \le B$ for all n, which is false since $M > 1$. □

Two further steps let constants depend on T without comment. In (12), the Lipschitz constant of F is "different for each t, but since t is finite, we can set a global L" (our L_F), although the proposition is asserted for all T. And Lemma 6 asserts χ for all t "since the model is stationary", although the recovered (v_t, s_t, w_{2t}) depend on a t-fold composition of the transition, whose sensitivity to δ can accumulate with t. The corrected proposition below makes both requirements explicit, and Appendix A verifies them from primitives in one case.

AGH's counterexample uses a Gaussian measurement shock, which does not have the compact support FRS assume. The failure does not depend on this.

Example 2 (A counterexample with compact support)**.** Let f be a smooth density on $[0,1]$ equal to a constant $c_0 > 0$ on $[\frac{1}{4}, \frac{3}{4}]$. Take the static model (S_t and W_t trivial) $y_t = g(v_t) = \gamma v_t$ with $v_t \sim f$ and $0 < \gamma < c_0$, approximated by $g_j(v) = (1+\delta)\gamma v$, so that $\|g_j - g\| \le \gamma\delta \le \delta$. FRS's assumptions hold. For data in $[\frac{1}{4}\gamma(1+\delta), \frac{3}{4}\gamma]$ the exact and computed densities are $M = c_0/\gamma > 1$ and $M/(1+\delta)$, so

$$L(y^T) - L_j(y^T) = M^T\big(1 - (1+\delta)^{-T}\big),$$

which for fixed δ grows exponentially in T and eventually exceeds any bound linear in T. With $\gamma > c_0$ instead, $M < 1$ and the difference vanishes exponentially, as in AGH. In both cases $\log L(y^T) - \log L_j(y^T) = T\log(1+\delta)$ exactly.

The last line of the example is the point of the next section: the level of the likelihood moves like M^T, but the log-likelihood gap is linear in T, as FRS's argument suggested.

3 The corrected proposition

The accumulation of error that FRS describe is multiplicative. It holds for the log likelihood, equivalently for the likelihood ratio L_j/L, and not for the level. We state the result under three assumptions; Appendix A derives all three from primitives.

Assumption A1 (Per-period error, uniformly in t)**.** There are $\xi > 0$ and $\varepsilon \geq 0$ such that $p_t(s,\omega) \geq \xi$, $p_{j,t}(s,\omega) \geq \xi$ and $|p_{j,t}(s,\omega) - p_t(s,\omega)| \leq \varepsilon$ for all t, s and ω.

In FRS's notation $\varepsilon = \chi\delta$; the lower bounds are their Assumptions 4 and 8.

Assumption A2 (Lipschitz dependence on the initial state)**.** There are $\ell_1, \ldots, \ell_T \geq 0$ such that $\log p_t(s',\omega) - \log p_t(s,\omega) \leq \ell_t\, d(s,s')$ for all t, s, s' and ω. Write $K_T = \sum_{t=1}^{T} \ell_t$.

Only the exact densities need this, not the computed ones.

Assumption A3 (Coupled initial distributions)**.** There is a probability space (X, P) and measurable maps $a, b : X \to S$ with laws μ^* and μ_j^* such that $d(a(x), b(x)) \leq \rho$ for P-almost every x.

Assumption A3 replaces FRS's use of Condition 1. It holds with $\rho = \delta/(1-\alpha)$ if Condition 1 is strengthened from contraction in mean to contraction for every shock, $d(\varphi(s,w), \varphi(s',w)) \leq \alpha\, d(s,s')$ for all w, and the computed chain's invariant distribution is unique (Theorem 9 in the appendix).

Corrected Proposition 2′. *Under Assumptions A1–A3,*

$$e^{-D}\, L(y^T) \;\leq\; L_j(y^T) \;\leq\; e^{D}\, L(y^T), \qquad D = \frac{T\varepsilon}{\xi} + K_T\, \rho,$$

equivalently $|\log L_j(y^T) - \log L(y^T)| \leq D$.

The proof uses two lemmas.

Lemma 3. *If $b \geq \xi > 0$ and $|a - b| \leq \varepsilon$, then $a \leq e^{\varepsilon/\xi}\, b$. Hence, under Assumption A1, for all s and ω,*

$$\prod_{t=1}^{T} p_{j,t}(s,\omega) \leq e^{T\varepsilon/\xi} \prod_{t=1}^{T} p_t(s,\omega) \qquad \text{and} \qquad \prod_{t=1}^{T} p_t(s,\omega) \leq e^{T\varepsilon/\xi} \prod_{t=1}^{T} p_{j,t}(s,\omega).$$

Proof. $a \leq b + \varepsilon \leq b + (\varepsilon/\xi)\, b = (1 + \varepsilon/\xi)\, b \leq e^{\varepsilon/\xi}\, b$, using $b/\xi \geq 1$ and $1 + x \leq e^x$. The product bounds follow by multiplying T such inequalities between nonnegative numbers; the second uses $p_{j,t} \geq \xi$. □

Lemma 4. *Under Assumption A2, $\prod_{t=1}^{T} p_t(s',\omega) \leq e^{K_T d(s,s')} \prod_{t=1}^{T} p_t(s,\omega)$ for all s, s' and ω.*

Proof. Multiply the inequalities $p_t(s',\omega) \leq e^{\ell_t d(s,s')}\, p_t(s,\omega)$ over t. □

Proof of Corrected Proposition 2′. Integrate out the shock path:

$$F(s) = \int \prod_t p_t(s,\omega)\, Q(\mathrm{d}\omega), \qquad F_j(s) = \int \prod_t p_{j,t}(s,\omega)\, Q(\mathrm{d}\omega).$$

By Assumption A3 and a change of variables, $L = \int_X F(a(x))\, P(\mathrm{d}x)$ and $L_j = \int_X F_j(b(x))\, P(\mathrm{d}x)$.

Upper bound. For all s, s' and ω, Lemma 3 and then Lemma 4 give

$$\prod_t p_{j,t}(s',\omega) \le e^{T\varepsilon/\xi} \prod_t p_t(s',\omega) \le e^{T\varepsilon/\xi + K_T d(s,s')} \prod_t p_t(s,\omega).$$

Integrating over ω, $F_j(s') \le e^{T\varepsilon/\xi + K_T d(s,s')} F(s)$. Put $s = a(x)$ and $s' = b(x)$. Since $d(a(x), b(x)) \le \rho$ almost surely and $F \ge 0$, $F_j(b(x)) \le e^D F(a(x))$ for P-almost every x, and integrating against P gives $L_j \le e^D L$.

Lower bound. Take the two moves in the other order: first move the initial state within the exact model, then change model. For all s, s' and ω,

$$\prod_t p_t(s,\omega) \le e^{K_T d(s',s)} \prod_t p_t(s',\omega) \le e^{K_T d(s',s) + T\varepsilon/\xi} \prod_t p_{j,t}(s',\omega),$$

by Lemma 4 and the second part of Lemma 3. Integrating over ω and then, with $s = a(x)$ and $s' = b(x)$, over P gives $L \le e^D L_j$. This ordering is why Assumption A2 is needed only for the exact model.

Finally, L and L_j are at least $\xi^T > 0$, since the densities are at least ξ and Q and P are probability measures, so taking logarithms gives the second form. □

Remark 5 (Sharpness). The dependence of D on T cannot be improved. With every exact density equal to M and every computed density equal to $M+\varepsilon$, $\log L_j - \log L = T \log(1+\varepsilon/M)$, which is linear in T; Example 2 is an instance within FRS's framework. In AGH's Gaussian model restricted to a compact support, data with every $y_t = \gamma + c$ give $\log L_j - \log L = T(c\delta - \delta^2/2)/\sigma^2$, again linear in T.

When the per-period Lipschitz constants decay geometrically, as they do when the exact chain forgets its initial state at rate α, the constant K_T is bounded uniformly in T and the bound takes exactly FRS's form, with the level replaced by the log.

Corollary 6. *If $\ell_t \le c\,\alpha^t$ with $0 \le \alpha < 1$ and $\rho = \delta/(1-\alpha)$, then for every T,*

$$|\log L_j(y^T) - \log L(y^T)| \le \frac{T\varepsilon}{\xi} + \frac{c\,\delta}{(1-\alpha)^2}, \qquad \frac{1}{T}|\log L_j(y^T) - \log L(y^T)| \le \frac{\varepsilon}{\xi} + \frac{c\,\delta}{T(1-\alpha)^2}.$$

Proof. $K_T \le c \sum_{t\ge 0} \alpha^t = c/(1-\alpha)$; substitute into Corrected Proposition 2′. □

In the case where the measurement shock alone is recovered from the data ($\dim V_t = \dim Y_t$), Theorem 11 in the appendix derives Assumptions A1–A3 and the geometric decay from primitives. If the computed transition is within δ of a pathwise α-contraction, the

per-state densities lie in $[\xi, M]$, differ by at most η and are c-Lipschitz in the state, and each chain has a unique invariant distribution, then

$$|\log L_j(y^T) - \log L(y^T)| \le \frac{T}{\xi}\Big(\eta + \frac{c\,\delta}{1-\alpha}\Big) + \frac{c\,\alpha}{\xi}\,\frac{\delta}{(1-\alpha)^2}.$$

When W_{2t} is also recovered from the data, Assumptions A1 and A2 amount to a stability condition on the inverted filter. FRS's Condition 1 does not imply it, and we know of no primitive condition in their setting that does. The closest results, on forgetting of the initial condition in hidden Markov models [Bickel et al., 1998, Douc et al., 2004] and on uniform-in-time approximation of nonlinear filters [Le Gland and Oudjane, 2004], assume a transition density bounded above and below, which a state that moves by a deterministic map of the shocks does not have.

4 Implications

The size of the error. FRS's statement that the error is of order $T\delta$ is correct for the relative error: $|L_j/L - 1| \le e^D - 1$, which is approximately D when D is small. It is false for the absolute error, which inherits the factor M^T carried by the level of the likelihood itself.

Estimation. By Corollary 6, the average log likelihood $\frac{1}{T}\log L_j$ is within $\varepsilon/\xi + O(\delta/T)$ of $\frac{1}{T}\log L$, uniformly in T. This is the dynamic analogue of the level part of AGH's approximation error Δ_j, now derived from the policy-function error and the dynamics of the state rather than assumed. It agrees with AGH's conclusion that approximation error of order δ moves the estimator's probability limit by $O(\delta)$, and not by more as T grows. A bound on the level of the objective alone delivers only $O(\sqrt{\delta})$ for the maximiser under quadratic identification; AGH's $O(\Delta_j)$ rate also uses closeness of the first two derivatives, which our result does not address.

Bayesian inference. FRS's Corollaries 1–4 assert that likelihood ratios, marginal likelihoods, posteriors and posterior expectations of the computed model converge to their exact counterparts. Corrected Proposition 2′ makes these quantitative. Suppose its conclusion holds at every parameter value with a common D, and let $\pi(\cdot \mid y^T) = L/Z$ and $\pi_j(\cdot \mid y^T) = L_j/Z_j$ be the posterior densities with respect to a prior π_0, where $Z = \int L\,\mathrm{d}\pi_0$ and $Z_j = \int L_j\,\mathrm{d}\pi_0$.

Proposition 7. *If $L, L_j > 0$ are π_0-integrable and $Z > 0$, then*

(i) $e^{-D}Z \le Z_j \le e^{D}Z$;

(ii) $L_j(\gamma')/L_j(\gamma) \le e^{2D}\,L(\gamma')/L(\gamma)$ *for all γ, γ', so every Metropolis–Hastings acceptance ratio is off by at most a factor e^{2D};*

(iii) $e^{-2D}\pi(\gamma \mid y^T) \le \pi_j(\gamma \mid y^T) \le e^{2D}\pi(\gamma \mid y^T)$;

(iv) the total variation distance between the posteriors is at most $(e^{2D} - 1)/2$;

(v) for measurable h with $|h| \le B$, $|\mathbb{E}_j[h \mid y^T] - \mathbb{E}[h \mid y^T]| \le B(e^{2D} - 1)$.

The proof is in Appendix B. Part (v) replaces FRS's Corollary 4, which asserts convergence without a rate, with an explicit bound. The closest result we know is the approximation theorem of Dashti and Stuart [2017, Theorem 4.9 in the arXiv version], building on Cotter et al. [2010]: if the negative log likelihoods satisfy $|\Phi - \Phi_N| \le M_2(\|u\|)\psi(N)$, the Hellinger distance between the posteriors is at most $C\psi(N)$ for N large. Corrected Proposition 2′ supplies that hypothesis with $\Phi = -\log L$ and $\psi = \delta$. Proposition 7 is the version with explicit constants, and it holds for every δ.

What remains of FRS's warnings. The claim that linearisation "is due to fail as the sample size grows" because the exact and computed likelihoods diverge does not survive as stated: both levels go to zero or infinity like M^T anyway, and the log-likelihood gap grows no faster than the log likelihood itself. For classical estimation based on the average log likelihood, a fixed δ produces a bias in point estimates that does not grow with T, and the same is true of posterior means in AGH's example. What grows is the bias relative to sampling uncertainty, which shrinks like $1/\sqrt{T}$: AGH's Theorem 4 requires $\sqrt{T}\Delta_j \to 0$ for the approximation to leave the asymptotic distribution unaffected, and in regular models the Bernstein–von Mises theorem gives the same condition for posteriors. The bounds of Proposition 7, which degrade like e^{2D} with $D \propto T\delta$, are conservative by comparison: in AGH's Gaussian example the total variation distance between the exact and computed posteriors is of order $\sqrt{T}\delta/\sigma$, not $T\delta$. Likelihood ratios themselves do move by a factor of order $e^{T\delta}$; in Example 2 the ratio between the exact and the computed likelihood is exactly $(1+\delta)^T$. In corrected form, then, FRS's warning stands with $\sqrt{T}$ in place of T: approximation error matters for inference, classical or Bayesian, once $\sqrt{T}\delta$ is not small, and for comparisons of likelihood levels once $T\delta$ is not small.

A The assumptions from primitives

This appendix derives Assumptions A1–A3 in the case $\dim V_t = \dim Y_t$, so that $W_1 = W$ and the measurement shock alone is recovered from the data. The state path, defined by s_0 given and $s_t = \varphi(s_{t-1}, \omega_t)$, is then a function $s_t(s_0, \omega)$ of the initial state and the shock path only, and $p_t(s_0, \omega) = f_t(s_t(s_0, \omega))$ for a function f_t of the current state: the density of V at $g^{-1}(y_t; s)$ times the Jacobian. Similarly $p_{j,t} = f_{j,t}(s_{j,t})$, with $s_{j,t}$ the computed path. We assume throughout that the state space S is a nonempty compact metric space, that φ and φ_j are jointly measurable and continuous in the state, that φ is a pathwise α-contraction, $d(\varphi(s,w), \varphi(s',w)) \le \alpha\, d(s,s')$ for all w with $0 \le \alpha < 1$, and that $d(\varphi_j(s,w), \varphi(s,w)) \le \delta$ with $\delta \ge 0$.

A.1 The coupling

Lemma 8 (Krylov–Bogolyubov for a random map)**.** *Let X be a nonempty compact metric space, ν a probability measure on a measurable space W, and $F : X \times W \to X$ jointly*

measurable with $F(\cdot, w)$ continuous for each w. Then there is a probability measure μ on X with $(\mu \otimes \nu) \circ F^{-1} = \mu$.

Proof. Write $T\mu = (\mu \otimes \nu) \circ F^{-1}$ and $(Mh)(x) = \int h(F(x, w))\, \nu(\mathrm{d}w)$ for bounded continuous h. By change of variables and Fubini's theorem, $\int h \,\mathrm{d}(T\mu) = \int Mh \,\mathrm{d}\mu$. By dominated convergence Mh is continuous, so T is weakly continuous. Start from a Dirac measure μ_0 and form the averages $\bar{\mu}_n = \frac{1}{n+1} \sum_{k=0}^{n} T^k \mu_0$. The difference $\int Mh \,\mathrm{d}\bar{\mu}_n - \int h \,\mathrm{d}\bar{\mu}_n$ telescopes to $\frac{1}{n+1}(\int h \,\mathrm{d}T^{n+1}\mu_0 - \int h \,\mathrm{d}\mu_0)$, which is at most $2\|h\|/(n+1)$ in absolute value. By Prokhorov's theorem the probability measures on X form a compact space, so $(\bar{\mu}_n)$ has a cluster point μ, and by continuity $\int Mh \,\mathrm{d}\mu = \int h \,\mathrm{d}\mu$ for every bounded continuous h. Hence $\int h \,\mathrm{d}(T\mu) = \int h \,\mathrm{d}\mu$ for all such h, and $T\mu = \mu$. □

Theorem 9 (The coupling)**.** *If μ^* is the unique invariant distribution of φ and μ_j^* the unique invariant distribution of φ_j, then Assumption A3 holds with $\rho = \delta/(1-\alpha)$.*

Proof. Let $B_\rho = \{(s, s') \in S \times S : d(s', s) \le \rho\}$, which is closed, hence compact, and contains the diagonal. Define $G((s, s'), w) = (\varphi(s, w), \varphi_j(s', w))$, both chains driven by the same shock. G maps B_ρ into itself, since

$$d(\varphi_j(s', w), \varphi(s, w)) \le d(\varphi_j(s', w), \varphi(s', w)) + d(\varphi(s', w), \varphi(s, w)) \le \delta + \alpha\rho = \rho.$$

It is jointly measurable and continuous in the pair, so Lemma 8 gives an invariant probability measure P on B_ρ. Because the first coordinate of G is φ applied to the first coordinate, taking the first marginal commutes with one step of the chain: $(P_1 \otimes \nu) \circ \varphi^{-1} = ((P \otimes \nu) \circ G^{-1})_1 = P_1$. So P_1 is invariant for φ and equals μ^* by uniqueness; likewise $P_2 = \mu_j^*$. Take $X = B_\rho$ and a, b the coordinate projections. □

Without uniqueness of the computed chain's invariant distribution, the conclusion holds for some invariant distribution of the computed chain, namely the one the coupling selects.

A.2 The densities

Lemma 10. *For all s_0, s_0', ω and t: (a) $d(s_{j,t}(s_0, \omega), s_t(s_0, \omega)) \le \delta/(1-\alpha)$; (b) $d(s_t(s_0, \omega), s_t(s_0', \omega)) \le \alpha^t d(s_0, s_0')$.*

Proof. (a) With $d_t = d(s_{j,t}, s_t)$, the triangle inequality as in the proof of Theorem 9 gives $d_{t+1} \le \alpha d_t + \delta$, so $d_t \le \alpha^t d_0 + \delta/(1-\alpha)$ by induction, and $d_0 = 0$. (b) Induction on t using the contraction. □

Theorem 11 (Corrected Proposition 2′ from primitives)**.** *Suppose each chain has a unique invariant distribution, the measurable functions $f_t, f_{j,t}$ take values in $[\xi, M]$ with $\xi > 0$, $|f_{j,t} - f_t| \le \eta$, and each f_t is c-Lipschitz. Then for every T and every law Q of the shock path,*

$$|\log L_j(y^T) - \log L(y^T)| \le \frac{T}{\xi}\Big(\eta + \frac{c\,\delta}{1-\alpha}\Big) + \frac{c\,\alpha}{\xi}\,\frac{\delta}{(1-\alpha)^2}.$$

Proof. Assumption A1. With $x = s_t(s_0, \omega)$ and $y = s_{j,t}(s_0, \omega)$, $|f_{j,t}(y) - f_t(x)| \le |f_{j,t}(y) - f_t(y)| + |f_t(y) - f_t(x)| \le \eta + c\, d(y, x)$, and $d(y, x) \le \delta/(1 - \alpha)$ by Lemma 10(a). So $\varepsilon = \eta + c\delta/(1 - \alpha)$, uniformly in t.

Assumption A2. With $x = s_t(s_0, \omega)$ and $x' = s_t(s_0', \omega)$, $|f_t(x') - f_t(x)| \le c\, \alpha^t d(s_0, s_0')$ by Lemma 10(b). Since $f_t(x) \ge \xi$, the first part of Lemma 3 gives $f_t(x') \le e^{(c/\xi)\alpha^t d(s_0, s_0')} f_t(x)$. This is Assumption A2 with $\ell_t = (c/\xi)\alpha^t$, so $K_T \le (c/\xi) \sum_{t \ge 1} \alpha^t = c\alpha/(\xi(1 - \alpha))$.

Assumption A3 is Theorem 9, with $\rho = \delta/(1 - \alpha)$. Corrected Proposition 2′ now gives $D = T\varepsilon/\xi + K_T \rho$, which is at most the stated bound.

Measurability. The state path is jointly measurable in (s_0, ω), by induction on t, so $\prod_t f_t(s_t)$ is jointly measurable and, by Fubini's theorem, its integral over ω is measurable in s_0. The bounds $0 \le \prod_t f_t \le M^T$ give integrability against every probability measure. □

What this leaves informal is the reduction of FRS's measurement equation to the per-state densities f_t, and the bound $\eta = O(\delta)$ on their error at a fixed state, which is FRS's Lemma 6 argument without the time dimension.

B Proof of Proposition 7

(i) Integrate $L_j \le e^D L$ and $L \le e^D L_j$ against π_0. (ii) $L_j(\gamma') \le e^D L(\gamma')$ and $1/L_j(\gamma) \le e^D / L(\gamma)$. (iii) Divide $L_j(\gamma) \le e^D L(\gamma)$ by $Z_j \ge e^{-D} Z$, and symmetrically. (iv) Write $a = \pi_j(\gamma \mid y^T)$ and $b = \pi(\gamma \mid y^T)$. By (iii), $a \le e^{2D} b$ and $b \le e^{2D} a$, so $a - b \le (e^{2D} - 1)b$ and $b - a \le (1 - e^{-2D})b \le (e^{2D} - 1)b$. Hence $|a - b| \le (e^{2D} - 1)b$, and integrating against π_0, using $\int b \, \mathrm{d}\pi_0 = 1$, gives $\int |\pi_j - \pi| \, \mathrm{d}\pi_0 \le e^{2D} - 1$, which is twice the total variation distance. (v) $|\int h\,(a - b) \, \mathrm{d}\pi_0| \le B \int |a - b| \, \mathrm{d}\pi_0 \le B(e^{2D} - 1)$. □

C Formal verification

Every result above has been verified in Lean 4 with Mathlib, in the repository cited on the title page. The development builds without unproved steps (`sorry`), and every result depends only on Lean's three standard axioms (`propext`, `Classical.choice`, `Quot.sound`). The formal likelihood is $\int_X \int_\Omega \prod_t p_t(a(x), \omega) \, \mathrm{d}Q \, \mathrm{d}P$ for an arbitrary finite set of periods; Theorem 11 converts it to the form against μ^* by change of variables. Paths are relative to `LeanEconomics/`.

Result	Lean name
Econometrics/ApproxLikelihood	
Proposition 1	`not_exists_linear_bound_prod_sub`
Log ratio linear in T (Example 2, Remark)	`log_prod_div_prod_const`
Lemma 3	`le_exp_div_mul`, `prod_le_exp_card_mul_prod`
Lemma 4	`prod_le_exp_sum_mul_prod`
Corrected Proposition 2′, the two bounds	`lik_le_exp_mul_lik`, `..._rev`
Corrected Proposition 2′, log form	`abs_log_likelihood_sub_le`
Corollary 6	`abs_log_likelihood_sub_le_of_geometric`
Coupled chain: band, path bound	`dist_step_le_of_le`, `dist_iterate_le`
Distribution/RandomMap	
Lemma 8	`RandomMap.exists_isInvariant`
Econometrics/ApproxLikelihoodModel	
Theorem 9	`exists_coupling`
Lemma 10	`dist_statePath_le`, `dist_statePath_exact_le`
Theorem 11, Assumption A1	`abs_densPath_sub_le`
Theorem 11, Assumption A2	`densPath_le_exp_mul`
Theorem 11	`abs_log_likelihood_sub_le_of_primitives`
Econometrics/ApproxPosterior	
Proposition 7 (i)–(iii)	`evidence_le_exp_mul`, `postDens_le_exp_mul`
Proposition 7 (iv)	`integral_abs_postDens_sub_le`
Proposition 7 (v)	`abs_integral_postDens_sub_le`

Example 2 is verified only through its algebraic core (`log_prod_div_prod_const`, and Proposition 1); the construction of the density f is not formalised.

References

Daniel Ackerberg, John Geweke, and Jinyong Hahn. Comments on "Convergence properties of the likelihood of computed dynamic models". *Econometrica*, 77(6):2009–2017, 2009. doi: 10.3982/ECTA7669.

Peter J. Bickel, Ya'acov Ritov, and Tobias Rydén. Asymptotic normality of the maximum-likelihood estimator for general hidden Markov models. *Annals of Statistics*, 26(4):1614–1635, 1998.

Simon L. Cotter, Masoumeh Dashti, and Andrew M. Stuart. Approximation of Bayesian inverse problems for PDEs. *SIAM Journal on Numerical Analysis*, 48(1):322–345, 2010. doi: 10.1137/090770734.

Masoumeh Dashti and Andrew M. Stuart. The Bayesian approach to inverse problems. In Roger Ghanem, David Higdon, and Houman Owhadi, editors, *Handbook of Uncertainty Quantification*. Springer, 2017. arXiv:1302.6989.

Randal Douc, Éric Moulines, and Tobias Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime. *Annals of Statistics*, 32 (5):2254–2304, 2004.

Jesús Fernández-Villaverde, Juan F. Rubio-Ramírez, and Manuel S. Santos. Convergence properties of the likelihood of computed dynamic models. *Econometrica*, 74(1):93–119, 2006.

John Geweke. Convergence properties of the likelihood of computed dynamic models: Comment. Unpublished manuscript, University of Iowa, 2007.

François Le Gland and Nadia Oudjane. Stability and uniform approximation of nonlinear filters using the Hilbert metric and application to particle filters. *Annals of Applied Probability*, 14(1):144–187, 2004.

Manuel S. Santos and Adrian Peralta-Alva. Accuracy of simulations for stochastic dynamic models. *Econometrica*, 73(6):1939–1976, 2005.