

Search for Heavy Neutral Leptons with IceCube DeepCore

Dissertation

zur Erlangung des akademischen Grades
doctor rerum naturalium
(Dr. rer. nat.)

im Fach: Physik
Spezialisierung: Experimentalphysik

eingereicht an der
Mathematisch-Naturwissenschaftlichen Fakultät
der Humboldt-Universität zu Berlin

von

Leander Fischer M. Sc.
geboren am 24. Oktober 1992
in Heidelberg

Präsidentin der Humboldt-Universität zu Berlin
Prof. Dr. Julia von Blumenthal

Dekanin der Mathematisch-Naturwissenschaftlichen Fakultät
Prof. Dr. Caren Tischendorf

Copyright Notice

This book is released into the public domain using the CC-BY-4.0 code.

To view a copy of the CC-BY-4.0 code, visit:

<https://creativecommons.org/licenses/by/4.0/>

Colophon

This document was typeset with the help of KOMA-Script and L^AT_EX using the open-source kaobook template class.

The source code of this thesis is available at:

https://github.com/LeanderFischer/phd_thesis

Zusammenfassung

Zusammenfassung ...

Abstract

Abstract ...

Todo list

Re-write/re-formulate this section (copied from HNL technote).	1
add varied total cross-section for a few background HNL events	2
Add comparisons of SM cross sections between NuXSplMkr and genie	3
add information about the matter profile used	9
add information about the oscillation probability calculation and the software used for it	9
get correct final level rates from my pipeline(s)	9
add rate and Poisson error for HNL samples	9
maybe just pick one mixing?	9
add 3D expectation and/or S/\sqrt{B} plots	10
Do I want/need to include the description of the KDE muon estimation?	10
Add table with all systematic uncertainties used in this analysis (in the analysis chapter).	10
add final level effects of varying the axial mass parameters (or example of one)	10
add final level effects of varying the DIS parameter (or example of one)	10
Do I want additional plots for this (fit diff, LLH distr, minim. stats, param. fits)?	11
Add bin-wise TS distribution? Add 3D TS maps?	11

Contents

Abstract	iii
Contents	vii
1 Heavy Neutral Lepton Signal Simulation	1
1.1 Model Independent Simulation	1
1.1.1 Generator Functions	1
1.1.2 Simplistic Sets	1
1.1.3 Realistic Set	1
1.2 Model Specific Simulation	1
1.2.1 Custom LeptonInjector	1
1.2.2 MadGraph5 3-Body Decays	4
1.2.3 Sampling Distributions	5
1.2.4 Weighting Scheme	5
1.3 Model Dependent Simulation Distributions	6
1.3.1 Generation Level	6
1.3.2 Final Level	6
2 Detecting Low Energetic Double Cascades	7
2.1 Reconstruction	7
2.1.1 Table-Based Minimum Likelihood Algorithms	7
2.1.2 Double Cascade Hypothesis	7
2.1.3 Modification to Low Energy Events	7
2.2 Performance	7
2.2.1 Energy/Decay Length Resolution	7
2.2.2 Double Cascade Classification	7
3 Search for an Excess of Heavy Neutral Lepton Events	9
3.1 Final Level Sample	9
3.1.1 Expected Rates/Events	9
3.1.2 Analysis Binning	10
3.2 Statistical Analysis	10
3.2.1 Test Statistic	10
3.2.2 Systematic Uncertainties	10
3.3 Analysis Checks	11
3.3.1 Minimization Robustness	11
3.3.2 Ensemble Tests	11
3.4 Results	12
3.4.1 Best Fit Parameters	12
3.4.2 Upper Limits	12
3.4.3 Post-Fit Data/MC Agreement	12
3.4.4 Likelihood Coverage	12
APPENDIX	13
A Analysis Checks	15
A.0.1 Minimization Robustness	15
A.0.2 Ensemble Tests	15
Bibliography	17

List of Figures

1.1	Custom HNL total cross sections for the four target masses compared to the total ($\nu_\tau/\bar{\nu}_\tau$ neutral current) cross section used for SM neutrino simulation production with GENIE.	2
1.2	Branching ratios of the HNL within the mass range considered, calculated based on the results from [3]. Given the existing constraints on $ U_{e4} ^2$ and $ U_{\mu4} ^2$, we consider that the corresponding decay modes are negligible.	3
3.1	Asimov inject/recover test (0.6 GeV)	12
3.2	Pseudo-data trials TS distribution (0.6 GeV)	12
A.1	Asimov inject/recover test (0.3 GeV, 1.0 GeV)	15
A.2	Pseudo-data trials TS distribution (0.3 GeV, 1.0 GeV)	15

List of Tables

1.1	xx	3
1.2	Sampling distributions of HNL simulation generation.	5
3.1	Final level event/rate expectation	9
3.2	Analysis binning	10
3.3	Staged minimization settings	11

Heavy Neutral Lepton Signal Simulation

1

1.1 Model Independent Simulation

1.1.1 Generator Functions

1.1.2 Simplistic Sets

1.1.3 Realistic Set

1.2 Model Specific Simulation

1.2.1 Custom LeptonInjector

Signal events are simulated using a **custom LeptonInjector (LI) tool** [1], modified from its standard version to include the HNL particle and the description of the HNL decays needed to produce the double cascade signature (currently only ν_τ related). In its SM work mode, LI injects a lepton and a cascade (under the general name *Hadrons*) at the interaction vertex of the neutrino. Both objects have the same (x,y,z,t) coordinates. In the modified version, the lepton at the interaction vertex is replaced by the HNL. After a chosen distance the HNL is forced to decay. The decay is sampled from the kinematically accessible decay modes shown in Figure 1.2.

A big addition to the standard LI is that the decay products of the HNL are added to the list of particles in the I3MCTree with a displaced position and delayed time from the interaction vertex. These daughter particles form a second cascade, not in the form of a *Hadrons* object, but as the explicit particles forming the shower. The kinematics of the two-body decays are computed analytically, while the three-body decays are dealt with using MadGraph5. To do so, we randomly pick an event from a list that we generated for each three-body decay mode. Independent of the number of particles in the final state of the HNL decay, the kinematics are calculated/simulated at rest and then boosted along the HNL momentum. The decay mode is randomly chosen based on the mass dependent branching ratios shown in Figure 1.2.


Each file is produced by running the **generation level processing script** using the filename as random seed and the above settings for the sampling distributions. The main part is calling the *MultiLeptonInjector* module in *volume mode* adding two generators (for ν_τ and $\bar{\nu}_\tau$) with 50% of the events. The generators are provided with the custom double-differential/total cross section splines described in Section 1.2.1 and the parameters defining the sampling distributions. For each frame *OneWeight* and a reference weight are also calculated and stored using the **weighting functions** and a baseline atmospheric ν_τ flux + oscillation spline. The weight will later be calculated inside of the analysis framework

1.1	Model Independent Simulation	1
1.2	Model Specific Simulation	1
1.3	Model Dependent Simulation Distributions	6

Re-write/re-formulate this section (copied from HNL technote).

[1]: Abbasi et al. (2021), “LeptonInjector and LeptonWeighter: A neutrino event generator and weighter for neutrino observatories”

Figure 1.1: Custom HNL total cross sections for the four target masses compared to the total ($\nu_\tau/\bar{\nu}_\tau$ neutral current) cross section used for SM neutrino simulation production with GENIE.



PISA, based on the input OneWeight. In addition to the i3 file itself, a LeptonInjector configuration file is written which stores the needed information to produce event weights using LeptonWeighter. Optionally the script can also produce an hdf5 file with the same name in the same location. This will store a fixed set of keys, extracted from the i3 file.

We are using *volume mode*, for the injection of the primary particle on a cylindrical volume. The main generation/sampling happens in `VolumeLeptonInjector::DAQ` inside

`LeptonInjector.cxx`. After writing the config (s) frame (currently not kept), the energy is sampled from a power law distribution, then the cosine(zenith) and azimuth angles are sampled from uniform distributions. The (x,y) position is sampled uniform in r, ϕ (for position on disk) and the z position is sampled from a uniform distribution. After the primary properties have been sampled the *EventProperties* is created and handed over to the `FillTree` functions which is where the custom HNL simulation happens:

add varied total cross-section for a few background HNL events

Cross Sections

The cross sections are calculated using a **modified version** of Carlos Argüelles' **NuXSplMkr**, which is a tool to calculate neutrino cross sections from parton distribution functions (PDFs) and then produce splines that can be read and used with IceCube software. The main modification to calculate the cross sections for the ν_τ neutral current interaction into the new heavy mass state is the addition of a kinematic condition to ensure that there is sufficient energy to produce the heavy mass state. It is the same condition that needs to be fulfilled for the charged current case, where the outgoing lepton mass is non-zero. Following [2] (equation 7), the condition

$$(1 + x\delta_N)h^2 - (x + \delta_4)h + x\delta_4 \leq 0, \quad (1.1)$$

is implemented for the neutral current case. Here $\delta_4 = \frac{m_4^2}{s-M^2}$, $\delta_N = \frac{M^2}{s-M^2}$, and $h \stackrel{\text{def}}{=} xy + \delta_4$, with x, y being the Bjorken variables, m_4 and M the mass of the heavy state and the target nucleon, respectively, and s the center of mass energy squared. Since the (SM) neutrino background simulation used for this analysis was created using GENIE (version 2.12.8), interfaced through the IceCube software package *genie-icetray*, with the **GRV98LO** PDFs, those were added as *GRV98lo_patched* to the

[2]: Levy (2009), "Cross-section and polarization of neutrino-produced tau's made simple"




Figure 1.2: Branching ratios of the HNL within the mass range considered, calculated based on the results from [3]. Given the existing constraints on $|U_{e4}|^2$ and $|U_{\mu4}|^2$, we consider that the corresponding decay modes are negligible.

Channel	Opens [MeV]	Max BR [%]
$\nu_4 \rightarrow \nu_\tau \nu_\alpha \bar{\nu}_\alpha$	0	100.0
$\nu_4 \rightarrow \nu_\tau e^+ e^-$	1	?
$\nu_4 \rightarrow \nu_\tau \pi^0$	135	?
$\nu_4 \rightarrow \nu_\tau \mu^+ \mu^-$	211	?
$\nu_4 \rightarrow \nu_\tau \eta$	548	?
$\nu_4 \rightarrow \nu_\tau \rho^0$	770	?
$\nu_4 \rightarrow \nu_\tau \omega$	783	?
$\nu_4 \rightarrow \nu_\tau \eta'$	958	?

Table 1.1: xx

cross section spline maker, to ensure the best possible agreement. Double-differential ($dsdx dy$) and total (σ) cross sections were produced for the four target HNL masses and then splined. The produced cross section splines are stored in the resources of the custom **LeptonInjector** module. Figure 1.1 shows the total cross sections that were produced compared to the cross section used for the production of the SM $\nu_\tau/\bar{\nu}_\tau$ neutral current background simulation .

Add comparisons of SM cross sections between NuXSS-plMkr and genie

Decay Channels

The accessible decay channels are dependent on the mass of the HNL and the allowed mixing. For this analysis, where only $|U_{\tau4}|^2 \neq 0$, the considered decay channels are listed in Table 1.1 and the corresponding branching ratios are shown in Figure 1.2. The individual branching ratio for a specific mass is calculated as $BR_i(m_4) = \Gamma_i(m_4)/\Gamma_{\text{total}}(m_4)$, where $\Gamma_{\text{total}}(m_4) = \sum \Gamma_i(m_4)$. The formulas to calculate the decay width show up in multiple references, but we chose to match them to [3], which also discusses the discrepancies in previous literature.

[3]: Coloma et al. (2021), “GeV-scale neutrinos: interactions with mesons and DUNE sensitivity”

2-Body Decay Widths The decay to a neutral pseudoscalar mesons is

$$\Gamma_{\nu_4 \rightarrow \nu_\tau P} = |U_{\tau4}|^2 \frac{G_F^2 m_4^3}{32\pi} f_P^2 (1 - x_p^2)^2, \quad (1.2)$$

with $x_P = m_P/m_4$ and

$$f_{\pi^0} = 0.130 \text{ GeV}, \quad f_\eta = 0.0816 \text{ GeV}, \quad C_2 = f_{\eta'} = -0.0946 \text{ GeV}, \quad (1.3)$$

while the decay to a neutral vector meson is given by

$$\Gamma_{\nu_4 \rightarrow \nu_\tau V} = |U_{\tau 4}|^2 \frac{G_F^2 m_4^3}{32\pi} \left(\frac{f_V}{m_V} \right)^2 g_V^2 (1 + 2x_V^2)(1 - x_V^2)^2, \quad (1.4)$$

with $x_V = m_V/m_4$,

$$f_{\rho^0} = 0.171 \text{ GeV}^2, \quad f_\omega = 0.155 \text{ GeV}^2, \quad (1.5)$$

and

$$g_{\rho^0} = 1 - 2 \sin^2 \theta_w, \quad g_\omega = \frac{-2 \sin^2 \theta_w}{3}, \quad \sin^2 \theta_w = 0.2229 \quad (1.6)$$

[4].

[4]: Tiesinga et al. (2021), "CODATA recommended values of the fundamental physical constants: 2018"

3-Body Decay Widths The (invisible) decay to three neutrinos is

$$\Gamma_{\nu_4 \rightarrow \nu_\tau \nu_\alpha \bar{\nu}_\alpha} = |U_{\tau 4}|^2 \frac{G_F^2 m_4^5}{192\pi^3}, \quad (1.7)$$

while the decay to two charged leptons (using $x_\alpha = (m_\alpha/m_4)^2$) of the same flavor reads

$$\Gamma_{\nu_4 \rightarrow \nu_\tau l_\alpha^+ l_\alpha^-} = |U_{\tau 4}|^2 \frac{G_F^2 m_4^5}{192\pi^3} [C_1 f_1(x_\alpha) + C_2 f_2(x_\alpha)], \quad (1.8)$$

with the constants defined as

$$C_1 = \frac{1}{4}(1 - 4s_w^2 + 8s_w^4), \quad C_2 = \frac{1}{2}(-s_w^2 + 2s_w^4), \quad (1.9)$$

the functions as

$$f_1(x_\alpha) = (1 - 14x_\alpha - 2x_\alpha^2 - 12x_\alpha^3)\sqrt{1 - 4x_\alpha} + 12x_\alpha^2(x_\alpha^2 - 1)L(x_\alpha), \quad (1.10)$$

$$f_2(x_\alpha) = 4[x_\alpha(2 + 10x_\alpha - 12x_\alpha^2)\sqrt{1 - 4x_\alpha} + 6x_\alpha^2(1 - 2x_\alpha + 2x_\alpha^2)L(x_\alpha)], \quad (1.11)$$

and

$$L(x) = \ln \left(\frac{1 - 3x_\alpha - (1 - x_\alpha)\sqrt{1 - 4x_\alpha}}{x_\alpha(1 + \sqrt{1 - 4x_\alpha})} \right). \quad (1.12)$$

1.2.2 MadGraph5 3-Body Decays

The code to produce the 3-body decay kinematics is [MadGraph4 v3.4.0](#) based on the decay diagrams calculated with [FeynRules 2.0](#) using the Lagrangians derived in [3]. The Universal FeynRules Output (UFO) from `effective_HeavyN_Majorana_v103` were used for our calculation. For each meass and corresponding decay channel, we produce 1e06 decay kinematic variations (rest frame) and store those in a text file.

[3]: Coloma et al. (2021), "GeV-scale neutrinos: interactions with mesons and DUNE sensitivity"

variable	distribution	range
energy	E^{-2}	$[2, 10^4]$ GeV
zenith	uniform (in $\cos(\theta)$)	$[180^\circ, 80^\circ]$
azimuth	uniform	$[0^\circ, 360^\circ]$
vertex (x, y)	uniform	$r = 600$ m
vertex z	uniform	$[-600, 0]$ m
m_{HNL}	fixed	$[0.3, 0.6, 1.0]$ GeV
L_{decay}	L^{-1}	$[0.0004, 1000.0]$ m / $[1.0, 1000.0]$ m

Table 1.2: Sampling distributions of HNL simulation generation.

1.2.3 Sampling Distributions

This is the description of the signal simulation generator used to (re-)start simulation production in December 2023. The underlying sampling distributions are listed in Table 1.2. Judging from how the generation/processing efficiency was for the 190607 set, we target $1e04$ files per set with $5e05$ events per file at generation, resulting in a maximum of $5e09$ events per set at generation level. Note here that the actual number of events per set at generation might be a little lower since some events won't be allowed if they don't have enough energy to produce the HNL.

1.2.4 Weighting Scheme

The weighting for the HNL signal simulation happens in a **custom stage of PISA**. The only input is the stored OneWeight and the variable physics parameter $|U_{\tau 4}|^2$, which is the mixing strength of the new heavy mass state and the tau sector. The custom re-weighting is needed to go from the used sampling PDF ($1/L$ with fixed range in lab frame decay length) to the target PDF (exponential defined by proper lifetime of the HNL). For each event the re-weighting factor is calculated using the gamma factor

$$\gamma = \frac{\sqrt{E_{\text{kin}}^2 + m_{\text{HNL}}^2}}{m_{\text{HNL}}}, \quad (1.13)$$

with the HNL mass m_{HNL} and it's kinetic energy E_{kin} . The speed of the HNL is calculated as

$$v = c \cdot \sqrt{1 - \frac{1}{\gamma^2}}, \quad (1.14)$$

where c is the speed of light. With these the lab frame decay length range can be converted into the rest frame lifetime range for each event

$$\tau_{\text{min/max}} = \frac{s_{\text{min/max}}}{v \cdot \gamma}. \quad (1.15)$$

The proper lifetime of each HNL event can be calculated using the total decay width Γ_{total} shown in Figure ?? and the chosen mixing strength $|U_{\tau 4}|^2$ as

$$\tau_{\text{proper}} = \frac{\hbar}{\Gamma_{\text{total}}(m_{\text{HNL}}) \cdot |U_{\tau 4}|^2}, \quad (1.16)$$

where \hbar is the reduced Planck constant. Since the decay length/lifetime of the events is sampled from an inverse distribution instead of an exponential as it would be expected from a particle decay we have to re-weight accordingly to achieve the correct decay length/lifetime

distribution. This is done by using the wanted exponential distribution

$$\text{PDF}_{\text{exp}} = \frac{1}{\tau_{\text{proper}}} \cdot e^{\frac{-\tau}{\tau_{\text{proper}}}}, \quad (1.17)$$

and the inverse distribution that was sampled from

$$\text{PDF}_{\text{inv}} = \frac{1}{\tau \cdot (\ln(\tau_{\text{max}}) - \ln(\tau_{\text{min}}))}. \quad (1.18)$$

The lifetime re-weighting factor is calculated as

$$w_{\text{lifetime}} = \frac{\text{PDF}_{\text{exp}}}{\text{PDF}_{\text{inv}}} = \frac{\Gamma_{\text{total}}(m_{\text{HNL}}) \cdot |U_{\tau 4}|^2}{\hbar} \cdot \tau \cdot (\ln(\tau_{\text{max}}) - \ln(\tau_{\text{min}})) \cdot e^{\frac{-\tau}{\tau_{\text{proper}}}}. \quad (1.19)$$

Adding another factor of $|U_{\tau 4}|^2$ to account for the mixing at the interaction vertex the total re-weighting factor becomes

$$w_{\text{total}} = |U_{\tau 4}|^2 \cdot w_{\text{lifetime}}, \quad (1.20)$$

which can be applied on top of flux and oscillation weight to get the final HNL weight for a given mixing (and mass).

1.3 Model Dependent Simulation Distributions

1.3.1 Generation Level

1.3.2 Final Level

Detecting Low Energetic Double Cascades

2

2.1 Reconstruction

2.1 Reconstruction 7

2.2 Performance 7

2.1.1 Table-Based Minimum Likelihood Algorithms

2.1.2 Double Cascade Hypothesis

2.1.3 Modification to Low Energy Events

2.2 Performance

2.2.1 Energy/Decay Length Resolution

2.2.2 Double Cascade Classification

Search for an Excess of Heavy Neutral Lepton Events

3

The measurement performed in this thesis is the search for an excess of HNL events in the 10 years of IceCube DeepCore data. In principle the two physics parameters to be probed are the mass of the HNL, m_4 , and the mixing between the fourth heavy mass state and the SM τ sector, $|U_{\tau 4}|^2$. Since the mass itself influences the production and decay kinematics of the event and the accessible decay modes, individual mass sets were produced as described in Section 1.2. The mass slightly influences the energy distribution, while the mixing both changes the overall scale of the HNL events and the shape in energy and PID. IceCube DeepCore is suited to measure the excess which appears around and below 20 GeV, due to its production from the atmospheric tau neutrinos, although a reduced lower energy threshold could improve the analysis. The measurement will be performed for the three mass sets individually, while the mixing is the parameter that can be varied continuously and will be measured in the fit.

3.1	Final Level Sample	9
3.2	Statistical Analysis	10
3.3	Analysis Checks	11
3.4	Results	12

3.1 Final Level Sample

The final level sample of this analysis always consists of the neutrino and muon MC introduced in Section ?? and one of the three HNL samples explained in Section 1.2. All of those simulation sets and the 10 years of IceCube DeepCore data are processed through the full processing and event selection chain described in Section ?? leading to the final level sample. Since applying the last cuts from Section ?? leaves an insignificant amount of pure noise events in the sample, the noise simulation is not included in the analysis and won't be listed here.

add information about the matter profile used

add information about the oscillation probability calculation and the software used for it

3.1.1 Expected Rates/Events

The rates and the expected events in 10 years are shown in Table 3.1. For the HNL the expectation depends on the mass and the mixing. Shown here are two example mixings for all the three masses. A mixing of 0.0 would result in a rate of 0.0 and therefore no HNL events.

get correct final level rates from my pipeline(s)

add rate and Poisson error for HNL samples

maybe just pick one mixing?

Type	Rate [mHz]	Events (in 10 years)	
ν_μ^{CC}	0.3522	103063 ± 113	
ν_e^{CC}	0.1411	41299 ± 69	
ν_τ^{CC}	0.0348	10187 ± 22	
ν_{NC}	0.0667	968 ± 57	
μ	0.0033	19522 ± 47	
HNL		$ U_{\tau 4} ^2 = 10^{-3}$	$ U_{\tau 4} ^2 = 10^{-1}$
$m_4=0.3 \text{ GeV}$	x.xxx	2.5	1342.5
$m_4=0.6 \text{ GeV}$	x.xxx	9.0	1207.0
$m_4=1.0 \text{ GeV}$	x.xxx	9.6	966.5

Table 3.1: Final level rates and event expectation of the SM background particle types and the HNL signal for all three masses and two example mixing values.

Table 3.2: Three dimensional binning used in the analysis. All variables are from the FLERCNN reconstruction explained in Section ??.

Variable	N_{bins}	Edges	Step
P_ν	3	[0.00, 0.25, 0.55, 1.00]	linear
E	12	[5.00, 100.00]	logarithmic
$\cos(\theta)$	8	[-1.00, 0.04]	linear

3.1.2 Analysis Binning

[5]: Yu et al. (2023), “Recent neutrino oscillation result with the IceCube experiment”

The identical binning to the analysis performed in [5] is used. It was chosen such that the track-like bin has the largest ν_μ -CC fraction. Extend the binning towards lower energies or increasing the number of bins did not improve the HNL sensitivities significantly. It also has to be considered that sufficient data events need to end up in the individual bins to result in a good fit, which was already investigated in the previous analysis. To mitigate the low data statistics, a few bins were not taken into account in the analysis. There are three bins in PID (cascade-like, mixed and track-like), 12 bins in reconstructed energy, and 8 bins in cosine of the reconstructed zenith angle as specified in Table 3.2. Originally, there were two more bins in $\cos(\theta)$, which were removed to reduce muons coming from the horizon and some low energy bins in the cascade-like bin are removed due to the low event expectation.

add 3D expectation and/or S/\sqrt{B} plots

3.2 Statistical Analysis

3.2.1 Test Statistic

The measurements are performed by comparing the weighted MC to the data. Through variation of the nuisance and physics parameters that govern the weights, the best matching set of parameters can be found. The comparison is done using a modified χ^2 defined as

$$\chi_{\text{mod}}^2 = \sum_{i \in \text{bins}} \frac{(N_i^\nu + N_i^\mu + N_i^{\text{HNL}} - N_i^{\text{obs}})^2}{N_i^\nu + N_i^\mu + N_i^{\text{HNL}} + (\sigma_i^\nu)^2 + (\sigma_i^\mu)^2 + (\sigma_i^{\text{HNL}})^2} + \sum_{j \in \text{syst}} \frac{(s_j - \hat{s}_j)^2}{\sigma_{s_j}^2}, \quad (3.1)$$

as the test statistic (TS), where N_i^ν , N_i^μ , and N_i^{HNL} are the expected number of events in bin i from neutrinos, atmospheric muons, and HNL, while N_i^{obs} is the observed number of events in bin i . The expected number of events from each particle type is calculated by summing the weights of all events in the bin $N_i^{\text{type}} = \sum_i^{\text{type}} \omega_i$, with the statistical uncertainty being $(\sigma_i^{\text{type}})^2 = \sum_i^{\text{type}} \omega_i^2$. The expected Poisson error is calculated using the combined expectation of neutrinos, atmospheric muons, and HNL events. The additional term in Equation 3.1 is included to apply a penalty term for prior knowledge of the systematic uncertainties of the parameters where they are known. s_j are the systematic parameters that are varied in the fit, while \hat{s}_j are their nominal values and σ_{s_j} are the known uncertainties.

Do I want/need to include the description of the KDE muon estimation?

Add table with all systematic uncertainties used in this analysis (in the analysis chapter).

add final level effects of varying the axial mass parameters (or example of one)

add final level effects of varying the DIS parameter (or example of one)

3.2.2 Systematic Uncertainties

3.3 Analysis Checks

Fitting to data will be performed in a *blind* manner, where the analyzer does not immediately see the fitted physics and nuisance parameter values, but first checks that a set of pre-defined *goodness of fit* (GOF) criteria are fulfilled. If those criteria are met to satisfaction the fit results are unblinded and the full result can be revealed. Before these blind fits to data are run, the robustness of the analysis method is tested using pseudo-data that is generated using the MC sets.

3.3.1 Minimization Robustness

To find the set of parameters that describes the data best, a staged minimization routine is used. In the first stage, a fit with coarse minimizer settings is performed to find a rough estimate of the *best fit point* (BFP). In the second stage, the fit is performed again in both octants¹ of θ_{23} , starting from the BFP of the coarse fit. For each individual fit the MIGRAD routine of MINUIT [6] is used to minimize the χ^2 TS defined in Equation 3.1. Iminuit is a fast, python compatible minimizer based on the MINUIT2 C++ library [7]. The individual minimizer settings are shown in Table 3.3.

To test the minimization routine and to make sure it consistently recovers any injected physics parameters, pseudo-data sets are produced from the MC by choosing the nominal nuisance parameters and specific physics parameters, without adding any statistical or systematic fluctuations to it. These so-called *Asimov*² data sets are then fit back with the full analysis chain. This type of test is called *Asimov inject/recover test*. A set of mixing values between 10^{-3} and 10^0 is injected and fit back. Even though this range is well within the excluded regions by other experiments, discussed in Section ??, this covers the current sensitive region of the analysis in IceCube DeepCore. Without fluctuations the fit is expected to always recover the injected parameters (both physics and nuisance parameters). The fitted mixing values from the Asimov inject/recover tests are compared to the true injected values in Figure 3.1 for the 0.6 GeV set. As expected, the fit is always able to recover the injected physics parameter and the nuisance parameters. Additional plots for the other mass sets can be found in Section A.0.1.

3.3.2 Ensemble Tests

To estimate the goodness of fit, pseudo-data is generated from the MC by injecting the BFP parameters as true parameters and then fluctuating the expected bin counts using Poisson fluctuation. The resulting pseudo-data sets are then fit back with the analysis chain. By comparing the distribution of TS values from this *ensemble* of pseudo-data trials to the TS of the fit to real data, a p-value can be calculated. The p-value is the probability of finding a TS value at least as large as the one from the data fit. Figure 3.2 shows the TS distribution from the ensemble tests for the 0.6 GeV mass set and the observed TS value from the fit, resulting in a p-value of 1.23 %³. Plots for the additional two mass sets are shown in Section A.0.2.

1: There is a degeneracy between the lower octant ($\theta_{23} < 45^\circ$) and the upper octant ($\theta_{23} > 45^\circ$), which can lead to TS minima (local and global) at two positions that are mirrored around 45° in θ_{23} .

[6]: Dembinski et al. (2022), *scikit-hep/iminuit: v2.17.0*

[7]: James et al. (1975), “Minuit: A System for Function Minimization and Analysis of the Parameter Errors and Correlations”

Fit	Err.	Prec.	Tol.
Coarse	1e-1	1e-8	1e-1
Fine	1e-5	1e-14	1e-5

Table 3.3: Migrad settings for the two stages in the minimization routine. *Err.* are the step size for the numerical gradient estimation, *Prec.* is the precision with which the LLH is calculated, and *Tol.* is the tolerance for the minimization.

2: A pseudo-data set without statistical fluctuations is called Asimov data set.

Do I want additional plots for this (fit diff, LLH distr, minim. stats, param. fits)?

Add bin-wise TS distribution? Add 3D TS maps?

3: The p-values for the 0.3 GeV and 1.0 GeV are 1.23 % and 1.23 %, respectively




Figure 3.1: Asimov inject/recover test for the 0.6 GeV mass set. Mixing values between 10^{-3} and 10^0 are injected and fit back with the full analysis chain. The injected parameter is always recovered within the statistical uncertainty.




Figure 3.2: Observed fit TS and TS distribution from pseudo-data trials for the 0.6 GeV mass set.

3.4 Results

3.4.1 Best Fit Parameters

3.4.2 Upper Limits

3.4.3 Post-Fit Data/MC Agreement

3.4.4 Likelihood Coverage

APPENDIX

Analysis Checks

A.0.1 Minimization Robustness

Figure A.1 shows additional Asimov inject/recover tests for the 0.3 GeV and 1.0 GeV mass sets. The tests were described in Section 3.3.1.




Figure A.1: Asimov inject/recover test for the 0.3 GeV (left) and 1.0 GeV (right) mass sets. Mixing values between 10^{-3} and 10^0 are injected and fit back with the full analysis chain. The injected parameter is always recovered within the statistical uncertainty.

A.0.2 Ensemble Tests

Figure A.2 shows additional TS distributions from pseudo-data trials and the observed TS from the fit to the data for the ensemble for the 0.3 GeV and 1.0 GeV mass sets. The tests were described in Section 3.3.2.




Figure A.2: Observed fit TS and TS distribution from pseudo-data trials for the 0.3 GeV (left) and 1.0 GeV (right) mass set.

Bibliography

Here are the references in citation order.

- [1] R. Abbasi et al. “LeptonInjector and LeptonWeighter: A neutrino event generator and weighter for neutrino observatories”. In: *Comput. Phys. Commun.* 266 (2021), p. 108018. doi: [10.1016/j.cpc.2021.108018](https://doi.org/10.1016/j.cpc.2021.108018) (cited on page 1).
- [2] J.-M. Levy. “Cross-section and polarization of neutrino-produced tau’s made simple”. In: *J. Phys. G* 36 (2009), p. 055002. doi: [10.1088/0954-3899/36/5/055002](https://doi.org/10.1088/0954-3899/36/5/055002) (cited on page 2).
- [3] P. Coloma et al. “GeV-scale neutrinos: interactions with mesons and DUNE sensitivity”. In: *Eur. Phys. J. C* 81.1 (2021), p. 78. doi: [10.1140/epjc/s10052-021-08861-y](https://doi.org/10.1140/epjc/s10052-021-08861-y) (cited on pages 3, 4).
- [4] E. Tiesinga et al. “CODATA recommended values of the fundamental physical constants: 2018”. In: *Rev. Mod. Phys.* 93 (2 July 2021), p. 025010. doi: [10.1103/RevModPhys.93.025010](https://doi.org/10.1103/RevModPhys.93.025010) (cited on page 4).
- [5] S. Yu and J. Micallef. “Recent neutrino oscillation result with the IceCube experiment”. In: *38th International Cosmic Ray Conference*. July 2023 (cited on page 10).
- [6] H. Dembinski et al. *scikit-hep/iminuit: v2.17.0*. Version v2.17.0. Sept. 2022. doi: [10.5281/zenodo.7115916](https://doi.org/10.5281/zenodo.7115916) (cited on page 11).
- [7] F. James and M. Roos. “Minuit: A System for Function Minimization and Analysis of the Parameter Errors and Correlations”. In: *Comput. Phys. Commun.* 10 (1975), pp. 343–367. doi: [10.1016/0010-4655\(75\)90039-9](https://doi.org/10.1016/0010-4655(75)90039-9) (cited on page 11).