

First Search for Heavy Neutral Leptons with IceCube DeepCore

Dissertation

zur Erlangung des akademischen Grades
doctor rerum naturalium
(Dr. rer. nat.)

im Fach: Physik
Spezialisierung: Experimentalphysik

eingereicht an der
Mathematisch-Naturwissenschaftlichen Fakultät
der Humboldt-Universität zu Berlin

von

Leander Fischer M. Sc.
geboren am 24. Oktober 1992
in Heidelberg

Präsidentin der Humboldt-Universität zu Berlin
Prof. Dr. Julia von Blumenthal

Dekanin der Mathematisch-Naturwissenschaftlichen Fakultät
Prof. Dr. Caren Tischendorf

Copyright Notice

This book is released into the public domain using the CC-BY-4.0 code.

To view a copy of the CC-BY-4.0 code, visit:

<https://creativecommons.org/licenses/by/4.0/>

Colophon

This document was typeset with the help of KOMA-Script and L^AT_EX using the open-source kaobook template class.

The source code of this thesis is available at:

https://github.com/LeanderFischer/phd_thesis

Abstract

first search for heavy neutral leptons in the GeV mass range with IceCube DeepCore extending the three flavor neutrino model by adding a fourth heavy mass state, considering three mass values m_4 of 0.3 GeV, 0.6 GeV, and 1.0 GeV and allowing only mixing with the tau neutrino through the mixing parameter $|U_{\tau 4}|^2$ the strength of the mixing is tested using atmospheric neutrinos as a production source of HNLs, using ten years of data taken between 2011 and 2021 to constrain the mixing parameter to $|U_{\tau 4}|^2 < 0.09 (m_4 = 0.3 \text{ GeV})$, $|U_{\tau 4}|^2 < 0.21 (m_4 = 0.6 \text{ GeV})$, and $|U_{\tau 4}|^2 < 0.24 (m_4 = 1.0 \text{ GeV})$ at 68% confidence level no significant signal of HNLs is observed for any of the tested masses, and the best fit mixing values obtained are consistent with the null hypothesis of no mixing thorough investigation of unique low energy double cascade signature of HNLs in IceCube benchmark reconstruction performance, with a well established reconstruction tool, after optimizing it for low energy double cascades identify the limitations to detect low energy double cascades and their origins lays fundamental groundwork for future searches for HNLs in IceCube

Zusammenfassung

Zusammenfassung ...

Todo list

highlight a few more neutrino related open questions, to circle back to related to the HNL searches maybe? (YELLOW)	4
add Majorana condition and mention what this means for interactions (LNV of 2) (ORANGE)	5
Discuss lepton number conservation (pure dirac) and lepton number violation (dirac+majorana) (ORANGE)	5
elaborate on Leptogenesis in ν MSM and sterile neutrino DM, or link some papers? (ORANGE)	6
I think here I'd want the extended leptonic EW lagrangien, so I can explain the mass mixing and the interactions it opens up (RED)	6
fix caption and check I mentioned all experiments (RED)	7
fix caption and check I mentioned all experiments (RED)	7
fix caption and check I mentioned all experiments (RED)	7
Say something about atmospheric neutrino flux uncertainties, based on recent JP/Anatoli papers. (YELLOW)	12
say something about matter effect? (ORANGE)	14
say something about mass ordering? (ORANGE)	14
fix up-scattering feynman diagram and put into the margin (RED)	15
fix the decay feynman diagrams and put into the margin (RED)	15

Contents

Abstract	iii
Contents	vii
1 Standard Model Neutrinos and Beyond	1
1.1 The Standard Model	1
1.1.1 Fundamental Fields	1
1.1.2 Electroweak Symmetry Breaking	2
1.1.3 Fermion Masses	3
1.1.4 Leptonic Weak Interactions after Symmetry Breaking	3
1.2 Beyond the Standard Model	3
1.2.1 Mass Mechanisms	4
1.2.2 Minimal Extensions and the ν MSM	5
1.2.3 Observational Avenues for Right-Handed Neutrinos	6
1.3 Atmospheric Neutrinos as Source of Heavy Neutral Leptons	11
1.3.1 Production of Neutrinos in the Atmosphere	11
1.3.2 Neutrino Oscillations	12
1.3.3 Neutrino Interactions with Nuclei	14
1.3.4 Heavy Neutral Lepton Production and Decay	15
Bibliography	17

List of Figures

1.1	Feynman diagrams of neutrino weak interactions	3
1.2	Current $ U_{e4}^2 - m_4$ limits	8
1.3	Current $ U_{\mu 4}^2 - m_4$ limits	9
1.4	Current $ U_{\tau 4}^2 - m_4$ limits	10
1.5	Atmospheric neutrino fluxes	11
1.6	Neutrino-nucleon deep inelastic scattering	14
1.7	Total inclusive neutrino-nucleon cross-sections	15
1.8	HNL decay widths	16
1.9	Theoretical mean HNL decay length (0.6 GeV mass)	16

List of Tables

1.1	Standard model fermions	2
1.2	Global fit neutrino mixing parameter results	13

Standard Model Neutrinos and Beyond

1

1.1 The Standard Model

The *Standard Model (SM)* of particle physics is a Yang-Mills theory [1] providing very accurate predictions of weak, strong, and *electromagnetic (EM)* interactions. It is a relativistic quantum field theory that relies on gauge invariance, where all matter is made up of fermions, which are divided into quarks and leptons, and bosons describe the interactions between the fermions that have to fulfil the overall symmetry of the theory. Leptons are excitations of Dirac-type fermion fields.

The initial idea of the theory is associated with the works of Weinberg [2], Glashow [3], and Salam [4], that proposed a unified description of EM and weak interactions as a theory of a spontaneously broken $SU(2) \times U(1)$ symmetry for leptons, predicting a neutral massive vector boson Z^0 , a massive charged vector boson W^\pm , and a massless photon γ as the gauge bosons. The Higgs mechanism [5], describing the breaking of the symmetry, predicts the existence of an additional scalar particle, the Higgs boson, giving the W^\pm and Z^0 bosons their mass. The Higgs boson was discovered in 2012 at the LHC [6, 7].

Gell-Mann and Zweig proposed the quark model in 1964 [8, 9], which was completed by the discovery of non-abelian gauge theories [10] to form the $SU(3)$ symmetry of the strong interaction called *quantum chromodynamics (QCD)*. QCD describes the interaction between quarks and gluons which completed the full picture of the SM in the mid-1970s. Together with the electroweak theory, the SM is a $SU(3)_C \times SU(2)_L \times U(1)_Y$ local gauge symmetry, with the conserved quantities C , *color*, L , *left-handed chirality*, and Y , *weak hypercharge*.

In the following, the basic properties of the SM are described, following the derivations of [11, 12].

1.1.1 Fundamental Fields

Fermions in the SM are Weyl fields with either *left-handed (LH)* or *right-handed (RH)* chirality, meaning they are eigenvectors of the chirality operator γ_5 with $\gamma_5 \psi_{R/L} = \pm \psi_{R/L}$. Only LH particles transform under $SU(2)_L$. The Higgs field is a complex scalar field, a doublet of $SU(2)_L$, which is responsible for the spontaneous symmetry breaking of $SU(2)_L \times U(1)_Y$ to $U(1)_{EM}$. Local gauge transformations of the fields are given by

$$\psi \rightarrow e^{ig\theta^a(x)T^a} \psi, \quad (1.1)$$

where g is the coupling constant, $\theta^a(x)$ are the parameters of the transformation, and T^a are the generators of the group, with a counting them. The number of bosons is dependent on the generators of the symmetry groups, while the strength is defined by the coupling constants. There are eight massless gluons corresponding to the generators of the $SU(3)_C$ group. These

1.1	The Standard Model	1
1.2	Beyond the Standard Model	3
1.3	Atmospheric Neutrinos as Source of Heavy Neutral Leptons . . .	11

[1]: Yang et al. (1954), "Conservation of Isotopic Spin and Isotopic Gauge Invariance"

[2]: Weinberg (1967), "A Model of Leptons"

[3]: Glashow (1961), "Partial-symmetries of weak interactions"

[5]: Higgs (1964), "Broken symmetries, massless particles and gauge fields"

[6]: Chatrchyan et al. (2012), "Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC"

[7]: Aad et al. (2012), "Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC"

[8]: Gell-Mann (1964), "A Schematic Model of Baryons and Mesons"

[9]: Zweig (1964), "An $SU(3)$ model for strong interaction symmetry and its breaking. Version 2"

[11]: Giunti et al. (2007), *Fundamentals of Neutrino Physics and Astrophysics*

[12]: Schwartz (2013), *Quantum Field Theory and the Standard Model*

mediate the strong force which conserves color charge. The W_1, W_2, W_3 , and B boson fields of the $SU(2)_L \times U(1)_Y$ group are mixed into the massive bosons through spontaneous symmetry breaking as

$$W^\pm = \frac{1}{\sqrt{2}}(W_1 \mp iW_2) \quad (1.2)$$

and

$$Z^0 = \cos \theta_W W_3 - \sin \theta_W B, \quad (1.3)$$

with θ_W being the *Weinberg angle*. The massless photon field is given by

$$A = \sin \theta_W W_3 + \cos \theta_W B \quad (1.4)$$

and its conserved quantity is the EM charge Q , which depends on the weak hypercharge, Y , and the third component of the weak isospin, T_3 , as $Q = T_3 + Y/2$.

	Type			Q
quarks	u	c	t	+2/3
	d	s	b	-1/3
leptons	ν_e	ν_μ	ν_τ	0
	e	μ	τ	-1

Table 1.1: Fermions in the Standard Model. Shown are all three generations of quarks and leptons with their electric charge Q .

Fermions are divided into six quarks and six leptons. Weak, strong, and EM force act on the quarks, and they are always found in bound form as baryons or mesons. Leptons do not participate in the strong interaction and only the electrically charged leptons are massive and are effected by the EM force, while neutrinos are massless and only interact via the weak force. Each charged lepton has an associated neutrino, which it interacts with in *charged-current* (CC) weak interactions, that will be explained in more detail in Section 1.1.4. The fermions are listed in Table 1.1.

1.1.2 Electroweak Symmetry Breaking

To elaborate the process of spontaneous symmetry breaking through which the gauge bosons of the weak interaction acquire their masses, the Lagrangian of the Higgs field is considered as

$$\mathcal{L}_{\text{Higgs}} = (D_\mu \Phi^\dagger)(D^\mu \Phi) - \lambda \left(\Phi^\dagger \Phi - \frac{v^2}{2} \right)^2, \quad (1.5)$$

with parameters λ and v , where λ is assumed to be positive. Φ is the Higgs doublet, which is defined as

$$\Phi = \begin{pmatrix} \Phi^+ \\ \Phi^0 \end{pmatrix}, \quad (1.6)$$

with the charged component Φ^+ and the neutral component Φ^0 . The covariant derivative is given by

$$D_\mu = \partial_\mu - ig_2 \frac{\sigma^i}{2} W_\mu^i - \frac{1}{2} ig_1 B_\mu, \quad (1.7)$$

with the Pauli matrices σ^i and the gauge boson fields W_μ^i and B_μ of the $SU(2)_L$ and $U(1)_Y$ groups, respectively. The coupling constants g_2 and g_1 are the respective coupling constants which are related to the Weinberg angle as $\tan \theta_W = \frac{g_1}{g_2}$. The Higgs potential has a non-zero *vacuum expectation value* (v) at the minimum of the potential at $\Phi^\dagger \Phi = \frac{v^2}{2}$. Since the vacuum is electrically neutral, it can only come from a neutral component of the Higgs

doublet as

$$\Phi_{\text{vev}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}. \quad (1.8)$$

1.1.3 Fermion Masses

The mass term for charged fermions with spin-1/2 is given by

$$\mathcal{L}_{\text{Dirac}} = m(\bar{\Psi}_R \Psi_L - \bar{\Psi}_L \Psi_R), \quad (1.9)$$

composed of the product of LH and RH Weyl spinors $\Psi_{L/R}$. This term is not invariant under $SU(2)_L \times U(1)_Y$ gauge transformations, but adding a Yukawa term

$$\mathcal{L}_{\text{Yukawa}} = -Y^e \bar{L}_L \Phi e_R + h.c., \quad (1.10)$$

coupling the fermion fields e_R to the Higgs field Φ , recovers the invariance and gives the fermions their masses. Here, Y^e is the Yukawa coupling constant and \bar{L}_L is the $SU(2)_L$ doublet. With the vev, this results in the mass term for the charged leptons and down-type quarks of $-m_e(\bar{e}_L e_R + \bar{e}_R e_L)$ with $m_e = \frac{Y^e v}{\sqrt{2}}$. With $\tilde{\Phi} = i\sigma_2 \Phi^*$, a similar Yukawa term can be written as $-Y^u \bar{L}_L \tilde{\Phi} u_R + h.c.$, which leads to the masses of the up-type quarks.

1.1.4 Leptonic Weak Interactions after Symmetry Breaking

After the spontaneous symmetry breaking, the leptonic part of the electroweak Lagrangian can be written as

$$\begin{aligned} \mathcal{L}_{\text{EW}}^\ell = & \frac{g}{\sqrt{2}} W^+ \sum_{\alpha=e,\mu,\tau} \bar{\nu}_\alpha \gamma^\mu P_L \ell_\alpha + \frac{g}{4c_w} Z \\ & \times \left\{ \sum_{\alpha=e,\mu,\tau} \bar{\nu}_\alpha \gamma^\mu P_L \nu_\alpha + \sum_\alpha \bar{\ell}_\alpha \gamma^\mu [2s_w^2 P_R - (1 - 2s_w^2) P_L] \ell_\alpha \right\} + h.c., \end{aligned} \quad (1.11)$$

where $c_w \equiv \cos \theta_w$, $s_w \equiv \sin \theta_w$, P_L and P_R are the left and right projectors, respectively, while ν_α and ℓ_α are the neutrino and charged lepton weak eigenstates. The W^\pm and Z bosons are the massive gauge bosons of the weak interaction. The large boson masses $m_W \sim 80 \text{ GeV}$ and $m_Z \sim 90 \text{ GeV}$ result in a short range of the force of about $1 \times 10^{-18} \text{ m}$. Interactions carried out by the W^\pm bosons are called *charged current (CC)* interactions, as they propagate a charge, therefore changing the interacting lepton to its charged/neutral counterpart. *Neutral current (NC)* interactions are those mediated by the Z^0 boson, where no charge is transferred. NC interactions couple neutrinos to neutrinos and charged leptons to charged leptons, but not to each other. The Feynman diagrams for CC and NC interactions are shown in Figure 1.1.




Figure 1.1: Feynman diagrams of charged-current (top) and neutral-current (bottom) neutrino weak interactions, modified from [13].

1.2 Beyond the Standard Model

The fundamentals of the SM described above **are not** enough to explain all observed phenomena. Gravity cannot be explained by the SM, as it is incompatible with general relativity, neither can some of the cosmological observations like *dark matter (DM)*, and the matter-antimatter asymmetry be explained. But most importantly, the SM does not predict neutrinos to

[14]: Deruelle et al. (2018), *Relativity in Modern Physics*

[15]: Workman et al. (2022), "Review of Particle Physics"

[16]: Fukugita et al. (1986), "Baryogenesis without grand unification"

[17]: Davis et al. (1968), "Search for Neutrinos from the Sun"

[18]: Fukuda et al. (1998), "Evidence for Oscillation of Atmospheric Neutrinos"

[19]: Ahmad et al. (2002), "Direct Evidence for Neutrino Flavor Transformation from Neutral-Current Interactions in the Sudbury Neutrino Observatory"

[20]: Alam et al. (2021), "Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory"

[21]: Aghanim et al. (2020), "Planck2018 results: VI. Cosmological parameters"

[22]: Aker et al. (2022), "Direct neutrino-mass measurement with sub-electronvolt sensitivity"

highlight a few more neutrino related open questions, to circle back to related to the HNL searches maybe? (YEL-Low)

have mass, which is experimentally proven by neutrino oscillations, so some extension to the SM is needed in order to explain the observed phenomena.

The standard cosmological model Λ CDM [14] assumes that equal amounts of matter and anti-matter were produced in the early universe. However, the universe today is dominantly made up of matter. This so-called *baryon asymmetry of the universe (BAU)* can be measured by the difference between the number densities of baryons and anti-baryons normalized to the number density of photons as

$$\eta_B = \frac{n_B - n_{\bar{B}}}{n_\gamma}, \quad (1.12)$$

where n_B , $n_{\bar{B}}$, and n_γ are the number densities of baryons, anti-baryons, and photons, respectively. Baryons are the dominant component with η_B being observed to be at the order of 10^{-9} [15]. Leptogenesis and EW baryogenesis are scenarios that could explain this phenomenon, where the former could be realized by the existence of heavy RH neutrinos [16].

The observation of neutrino flavor conversions and neutrino oscillations in a multitude of experiments [17–19] is the strongest evidence for physics *beyond the standard model (BSM)* measured in laboratories to date. The observation that neutrinos change their flavor while they propagate through space can only be explained, if at least two neutrinos have a non-zero mass. From those measurements we know the mass differences are very small as compared to the lepton masses, but neither their existence, nor their smallness is predicted by the SM. There are upper limits on the sum of all neutrino masses from cosmological observations at 1.2 eV [20, 21] and at 0.8 eV from the KATRIN experiment [22]. Adding RH neutrino states to the theory could explain the origin of the observed non-zero neutrino masses and could be tested for by searching for corresponding signatures in experiments.

1.2.1 Mass Mechanisms

Since there are no RH neutrinos in the SM, the mass mechanism described in Section 1.1.3, which couples the Higgs field to LH and RH Weyl fields, predicts the LH neutrinos to be massless. From experimental observations it is known that at least two of the three neutrino generations need to have a non-zero mass. Assuming the existence of RH neutrinos fields ν_R , one way of producing the neutrino masses is by adding a Yukawa coupling term similar to the one for up-type quarks mentioned in Section 1.1.3, to write the full Yukawa Lagrangian as

$$\mathcal{L}_{\text{Yukawa}} = -Y_{ij}^e \bar{L}_L^i \Phi e_R^j - Y_{ij}^\nu \bar{L}_L^i \tilde{\Phi} \nu_R^j + h.c., \quad (1.13)$$

with i, j running over the three generations of leptons e, μ , and τ , and Y^e and Y^ν being the Yukawa coupling matrices. Diagonalizing the Yukawa coupling matrices through unitary transformations U^e and U^ν leads to the **Dirac mass term** in the mass basis as

$$\mathcal{L}_{\text{Dirac}}^{\text{mass}} = \frac{v}{\sqrt{2}} (\bar{e}_L M_e e_R - \bar{\nu}_L M_\nu \nu_R), \quad (1.14)$$

where M_e and M_ν are the diagonal mass matrices of leptons and neutrinos, respectively. A purely Dirac mass term would not explain the smallness of the neutrino masses in a straightforward way. Only fine-tuning the Yukawa coupling constants to small values would lead to small neutrino masses.

An additional way of generating neutrino masses is by adding a Majorana mass term of the form

$$\mathcal{L}_{\text{Majorana}} = -\frac{1}{2} M_{ij} (\nu_R^i)^c \nu_R^j + h.c. , \quad (1.15)$$

with M_{ij} being the Majorana mass matrix and the indices i, j running over all n_R RH neutrino generations. The superscript c denotes the charge conjugate field. Combining the charge conjugated RH neutrino fields with the LH neutrino fields as

$$\mathbf{N} = \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix} , \quad (1.16)$$

with ν_R containing the n_R RH fields. The full neutrino mass Lagrangian is then given by the combined **Dirac and Majorana mass term** as

$$\mathcal{L}_{\text{Dirac+Majorana}}^{\text{mass},\nu} = \frac{1}{2} \mathbf{N}^T \hat{C} M^{\text{D+M}} \mathbf{N} + h.c. , \quad (1.17)$$

and the mass matrix is given by

$$M^{\text{D+M}} = \begin{pmatrix} 0 & (M^{\text{D}})^T \\ M^{\text{D}} & M^{\text{R}} \end{pmatrix} . \quad (1.18)$$

On top of explaining the origin of neutrino masses itself, a combined Dirac and Majorana mass term could also solve the question of their smallness. If the mass of the RH neutrinos is very large, the masses of the active neutrino flavors is suppressed, which is known as *see-saw mechanism*.

add Majorana condition and mention what this means for interactions (LNV of 2) (ORANGE)

1.2.2 Minimal Extensions and the ν MSM

So far we have described neutrinos in their flavor eigenstates, which are relevant for weak interactions, where the three weak flavor states ν_e, ν_μ , and ν_τ are related to the charged leptons they interact with in CC interactions. In order to *just* explain the three oscillating flavor eigenstates, three mass states are needed, which are related to the flavor eigenstates by the unitary, 3×3 *Pontecorvo-Maki-Nakagawa-Sakata* (PMNS) mixing matrix U , where the flavor states are a superposition of the mass states as

$$|\nu_\alpha\rangle = \sum_k U_{\alpha k}^* |\nu_k\rangle , \quad (1.19)$$

with the weak flavor states $|\nu_\alpha\rangle$, $\alpha = e, \mu, \tau$, and the mass states $|\nu_k\rangle$ with $k = 1, 2, 3$. In its generic form the PMNS matrix is given by

$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} \end{pmatrix} , \quad (1.20)$$

which will be the basis for the discussion of neutrino oscillations in Section 1.3.2.

This however is not enough to explain the neutrino masses observed in oscillation experiments. The most minimal model required to give rise to two non-zero active neutrino masses, is an additional two RH neutrinos, assuming the mass of the lightest SM neutrino is zero. If the additional neutrino states have masses $\gg \text{eV}$ they are referred to as *heavy neutral leptons*

Discuss lepton number conservation (pure dirac) and lepton number violation (dirac+majorana) (ORANGE)

[23]: Asaka et al. (2005), “The nuMSM, dark matter and neutrino masses”

[24]: Asaka et al. (2005), “The ν MSM, dark matter and baryon asymmetry of the universe”

elaborate on Leptogenesis in ν MSM and sterile neutrino DM, or link some papers? (ORANGE)

[25]: Minkowski (1977), “ $\mu \rightarrow e \gamma$ at a rate of one out of 10^9 muon decays?”

[26]: Yanagida (1980), “Horizontal Symmetry and Masses of Neutrinos”

[27]: Glashow (1980), “The Future of Elementary Particle Physics”

[28]: Gell-Mann et al. (1979), “Complex Spinors and Unified Theories”

[29]: Mohapatra et al. (1980), “Neutrino Mass and Spontaneous Parity Nonconservation”

I think here I'd want the extended leptonic EW lagrangien, so I can explain the mass mixing and the interactions it opens up (RED)

[30]: Aartsen et al. (2020), “eV-Scale Sterile Neutrino Search Using Eight Years of Atmospheric Muon Neutrino Data from the IceCube Neutrino Observatory”

[13]: Trettin (2023), “Search for eV-scale sterile neutrinos with IceCube DeepCore”

[31]: Tastet et al. (2021), “Reinterpreting the ATLAS bounds on heavy neutral leptons in a realistic neutrino oscillation model”

(HNL), which are almost sterile, with a small mass mixing with the active neutrinos.

But the SM also fails to explain additional observations of physics beyond the standard model (BAU, DM), which could be solved by the *neutrino minimal standard model* (ν MSM) [23, 24]. In the ν MSM, three RH neutrinos are added, where two of them are heavy, to explain the observed neutrino masses and oscillations, and a third one is light and serves as a DM candidate. The mixing between mass and flavor eigenstates is then described by an extended 6x6 mixing matrix as

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ N_1 \\ N_2 \\ N_3 \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} & U_{e5} & U_{e6} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} & U_{\mu4} & U_{\mu5} & U_{\mu6} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} & U_{\tau4} & U_{\tau5} & U_{\tau6} \\ U_{N_11} & U_{N_12} & U_{N_13} & U_{N_14} & U_{N_15} & U_{N_16} \\ U_{N_21} & U_{N_22} & U_{N_23} & U_{N_24} & U_{N_25} & U_{N_26} \\ U_{N_31} & U_{N_32} & U_{N_33} & U_{N_34} & U_{N_35} & U_{N_36} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \\ \nu_4 \\ \nu_5 \\ \nu_6 \end{pmatrix}, \quad (1.21)$$

where N_i and ν_{i+3} ($i \in [1, 2, 3]$) are the sterile flavor states and the additional RH mass states, respectively. In the ν MSM, the two heavy RH neutrinos generate the active neutrino masses through the type I seesaw mechanism [25–29], where they are assumed to be SM scalars and couple to the Higgs field as such.

1.2.3 Observational Avenues for Right-Handed Neutrinos

If the RH neutrinos have masses at the eV scale, they can be observed through distortion effects in measurements of neutrino oscillation experiments. Several analyses looking for these so-called light sterile neutrinos exist in IceCube, where [30] is using atmospheric neutrinos in the higher energy range of 500 GeV to 10 000 GeV and [13] is using the lower energy region of 6 GeV to 156 GeV. The latter work includes a detailed description of the expected oscillation effects and the various anomalies observed in oscillation experiments that could be explained by the existence of a light sterile neutrino, which is not covered in this work.

Here, the focus will be on heavy RH neutrinos, interchangeably also called heavy sterile neutrinos, or HNLs. A defining property is that they are too massive to be produce in oscillations and to be observed as distortions thereof. Several ways to observe HNLs are possible through direct production and decay experiments, which will be discussed in the following. Most of the existing searches assume the minimal model, where only one coupling between the new mass states and the SM neutrinos is non-zero and the coupling is just through mass mixing in a type I seesaw scenario, but more complex scenarios are of course also possible and might produce various additional signatures, or stronger signals.

In general, the constraints discussed in the following are based on models, where only the coupling between the HNL and one SM flavor is non-zero. While this is the straight forward approach to test the mixing parameters individually, this might make the constraints stronger than they would be in a more complex scenario, where the HNLs couple to more than one SM flavor as was show in [31] for collider bounds.

Extracted Beamline Searches

Protons interacting with a target or a beam dump can produce pions, kaons, and heavy-quark hadrons, whose subsequent decays would also produce HNLs. The energies of the HNLs produced in those interactions are between 1 MeV and 4 GeV and could decay at several distances, depending on their lifetime, which is model dependent. Experiments along the extracted beamline, which are using a spectrometer with particle identification, can search for unique decay signatures at displaced vertices. Example signatures are $\nu_4 \rightarrow l_\alpha \pi$, $\nu_4 \rightarrow l_\alpha^+ l_\alpha^-$, or $\nu_4 \rightarrow \nu \pi^0$ (or other neutral mesons), which cannot be explained by SM neutrinos. Depending on the decay channel, a specific mixing can be probed. The other way of searching for HNLs with these interactions is to look for peaks in the missing mass spectrum, measured around the production vertex at the target, which usually is not possible for beam dumps, as the beam dump region is not calorimetrically instrumented.

HNL search pioneer work was done by experiments at extracted beam lines, with first results from *PS191* [32] and *CHARM* [33], reporting bounds from direct production and decay of HNLs for $|U_{e4}|^2$, $|U_{\mu 4}|^2$, and combinations of them, at masses from 10 MeV to 500 MeV at orders of 10^{-3} to 10^{-6} . Since then, there has been and still is a large activity of searches for HNLs at extracted beamlines and the strongest bounds on $|U_{e4}|^2$ and $|U_{\mu 4}|^2$ are currently set by *PIENU* [34–36], *TRIUMF* [37], *NA62* [38], *T2K* [39], *MicroBooNE* [40], and *NuTeV* [41] in the mass range from 1 MeV to 4 GeV, reaching from 10^{-4} at the lower mass to 10^{-9} at 4 at the highest masses. The current strongest bounds are shown in Figure 1.4 and Figure ??, which also shows bounds from other experiments, which will be discussed in the following.

Especially noteworthy are the results of analyses probing the mixing with the third lepton generation, $|U_{\tau 4}|^2$, from *NOMAD* [42] and reinterpretations of the *CHARM* results and the *BEBC* results in the context of the mixing $|U_{\tau 4}|^2$, where the latter is place the most stringent limits from 10^{-3} to 10^{-6} in the 0.1 GeV to 2 GeV range [43–45]. In Figure ?? the current strongest bounds on $|U_{\tau 4}|^2$ are shown.

Collider Searches

So far, collider searches have been conducted at the *large electron positron collider (LEP)* and at the *large hadron collider (LHC)* in proton-proton mode. Strongest results are from the *ATLAS* and *CMS* experiments, which are nearly hermetic, general purpose detectors around the interaction point, and from the *DELPHI* and the *LHCb* experiments, which are forward detectors that can be used to search for new particles in decays of heavy particles produced. In the minimal model, HNLs in the GeV mass range can be produced through mass mixing in decays of heavy mesons, tau leptons, Z/W bosons, H bosons, or top quarks originating from the collisions. Depending on the dirac or majorana nature of the HNL, they can decay to lepton number conserving or lepton number violating channels.

Analyzing Z boson decays, DELPHI has set constraints on

Using prompt and displaced decays of the HNL, ATLAS has set constraints on $|U_{\mu 4}|^2$ at the level of 10^{-5} to 10^{-6} in the mass range of 4 GeV to 10 GeV [47]. Similarly, CMS constrained both $|U_{e4}|^2$ and $|U_{\mu 4}|^2$ at the level of 10^{-5}

[32]: Bernardi et al. (1986), “Search for Neutrino Decay”

[33]: Bergsma et al. (1983), “A Search for Decays of Heavy Neutrinos”

[34]: Ito et al. (2021), “Search for heavy neutrinos in $\pi^+ \rightarrow \mu^+ \nu$ decay and status of lepton universality test in the PIENU experiment”

[35]: Aguilar-Arevalo et al. (2018), “Improved search for heavy neutrinos in the decay $\pi \rightarrow e \nu$ ”

[36]: Bryman et al. (2019), “Constraints on Sterile Neutrinos in the MeV to GeV Mass Range”

[37]: Britton et al. (1992), “Improved search for massive neutrinos in $\pi + \rightarrow e + \nu$ decay”

[38]: Parkinson et al. (2022), “Search for heavy neutral lepton production at the NA62 experiment”

[39]: Abe et al. (2019), “Search for heavy neutrinos with the T2K near detector ND280”

[40]: Abratenko et al. (2024), “Search for Heavy Neutral Leptons in Electron-Positron and Neutral-Pion Final States with the MicroBooNE Detector”

[41]: Vaitaitis et al. (1999), “Search for neutral heavy leptons in a high-energy neutrino beam”

[42]: Astier et al. (2001), “Search for heavy neutrinos mixing with tau neutrinos”

[43]: Orloff et al. (2002), “Limits on the mixing of tau neutrino to heavy neutrinos”

[44]: Boiarska et al. (2021), “Blast from the past: constraints from the CHARM experiment on Heavy Neutral Leptons with tau mixing”

[45]: Barouki et al. (2022), “Blast from the past II: Constraints on heavy neutral leptons from the BEBC WA66 beam dump experiment”

fix caption and check I mentioned all experiments (RED)

fix caption and check I mentioned all experiments (RED)

fix caption and check I mentioned all experiments (RED)

[47]: Aad et al. (2019), “Search for heavy neutral leptons in decays of W bosons produced in 13 TeV pp collisions using prompt and displaced signatures with the ATLAS detector”




Figure 1.2: Current $|U_{e4}|^2 - m_4$ limits from xx. Modified from [46].

[48]: Tumasyan et al. (2022), “Search for long-lived heavy neutral leptons with displaced vertices in proton-proton collisions at $\sqrt{s} = 13$ TeV”

[49]: Sirunyan et al. (2018), “Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV”

[50]: Shuve et al. (2016), “Revision of the LHCb Limit on Majorana Neutrinos”

[51]: Aaij et al. (2021), “Search for heavy neutral leptons in $W^+ \rightarrow \mu^+ \mu^\pm \text{jet}$ decays”

in the mass range of 10 GeV to 60 GeV [48]. Another CMS search, looking for final states with three leptons, has set constraints on $|U_{\mu 4}|^2$ at the level of 10^{-7} in the mass range of 8 GeV to 14 GeV [49].

The LHCb experiment has HNL search results at HNL masses below and above the W boson mass, where the low mass searches are using the decay channel $B^- \rightarrow \pi^+ \mu^- \mu^-$, setting limits at the 10^{-3} level for $|U_{\mu 4}|^2$ in the mass range of 0.5 GeV to 3.5 GeV [50]. At high masses, the $W^+ \rightarrow \mu^- \mu^\pm \text{jet}$ channel is used to set limits at the order of 10^{-3} to 10^{-2} for $|U_{\mu 4}|^2$ in the mass range of 5 GeV to 50 GeV in the LNC channel and at the order of 10^{-4} to 10^{-3} in the LNV channel [51].

Nuclear Decays Measurements

A novel approach of searching for irregularities in energy-momentum conservation measurements in nuclear reactions might be a viable way of searching for HNLs, as they could be interpreted as constraints on $|U_{e4}|^2$ and m_4 .

[52]: Osipowicz et al. (2001), “KATRIN: A Next generation tritium beta decay experiment with sub-eV sensitivity for the electron neutrino mass. Letter of intent”

[53]: Mertens et al. (2019), “A novel detector system for KATRIN to search for keV-scale sterile neutrinos”

[54]: Aker et al. (2023), “Search for keV-scale sterile neutrinos with the first KATRIN data”

[55]: Abi et al. (2020), “Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume II: DUNE Physics”

Kinks in **beta decay** spectra would show up at $Q - m_4 c^2$, where the HNL mass, m_4 , can be measured between the lower energy detection threshold and the energy released in the decay, which is called Q value. Analyses using the tritium decay, with $Q = 18.6$ keV, are planned in *KATRIN* [52] and *TRISTAN* [53] in the 1 keV to 18 keV range. Their projected statistical limits are around 10^{-7} for $|U_{e4}|^2$, but will require further detector upgrades [53]. A first result from KATRIN measurements during commissioning sets limits at the order of 10^{-2} to 10^{-3} in the mass range of 0.1 keV to 1.6 keV [54]. *DUNE* is planning to measure the ionization charge of atmospheric argon decays, with $Q = 565$ keV, to probe $|U_{e4}|^2$ at in the 20 keV to 450 keV mass range. The projected sensitivity is at the 10^{-5} level, and might improve to 10^{-7} with additional detector improvements [55].




Figure 1.3: Current $|U_{\mu 4}^2| - m_4$ limits from xx. Modified from [46].

To test for the existence of HNLs using **electron capture** measurements, total energy-momentum reconstruction of all non-neutrino final states is needed. Electron capture is a pure two body decay process, where the recoiling atom and the electron neutrino are the only final state particles, but additional energy is carried away by the de-excitation x-ray or auger electron. The energy-momentum conservation can be probed by measuring the atom and the associated de-excitation products. The mixing $|U_{e4}|^2$ can be probed by looking for a separated non-zero missing mass peak. The *BeEST* experiment has set limits at the 10^{-4} level in the 100 keV to 850 keV mass range, using berillium-7, which has a Q value of 862 keV. After planned upgrades to the experiment, the sensitivity is expected to improve to the 10^{-7} level [56].

Reactor searches up to 12 MeV in mass are possible at short baseline experiments using commercial or research reactors, which are a strong source of electron antineutrinos and could therefore also produce HNLs if $|U_{e4}|^2$ is non-zero. Visible decay channels at these energies are $\nu_4 \rightarrow \nu_e e^+ e^-$, $\nu_4 \rightarrow \nu \gamma$, and $\nu_4 \rightarrow \nu \gamma \gamma$, where the first dominates. The first analysis in this field, reports limits at the 10^{-4} level in the 2 MeV to 7 MeV mass range [57].

Atmospheric and Solar

Natural sources of neutrinos are provided up to 20 MeV by the sun and up to 100s of GeV by neutrino production in the atmosphere. Both fluxes contain all flavors of neutrinos, due to mixing and oscillations, and can therefore be used to directly probe the mixings with ν_e , ν_μ , and ν_τ . Depending on the HNL mass and the strength of the mixing, which both govern the decay length, different signatures can be used to experimentally access large regions of the HNL parameter space. The strength of the mixing defines the total rate of HNL events, which is additionally affected by whether solely the minimal mass mixing is assumed, or also more complicated mixing scenarios, like the dipole portal, are considered.

[56]: Friedrich et al. (2021), “Limits on the Existence of sub-MeV Sterile Neutrinos from the Decay of ^7Be in Superconducting Quantum Sensors”

[57]: Hagner et al. (1995), “Experimental search for the neutrino decay $\nu_3 + \nu_j + e^+ + e^-$ and limits on neutrino mixing”




Figure 1.4: Current $|U_{\tau 4}|^2 - m_4$ limits from xx. Modified from [46].

So far, only very few analyses exist, which are performed by the experimental collaborations themselves. Several external theoretical groups have predicted the expected sensitivities to HNLs, produced from solar or atmospheric neutrinos, based on various coupling scenarios and decay lengths. A selection of the potential analyses will be discussed in the following.

For very long-lived particles, **production inside the sun** can be used as a source to search for HNLs in detectors on earth. This will only allow production through non-zero $|U_{e4}|^2$, because the initial solar neutrino flux is only ν_e . By searching for HNL decays to a SM neutrino and an electron positron pair $\nu_4 \rightarrow \nu_e e^+ e^-$ and comparing to the expected inter planetary positron flux, *Borexino* has placed the strongest limits on the mixing $|U_{e4}|^2$ at the order of 10^{-5} in the few MeV mass range [58].

[58]: Bellini et al. (2013), “New limits on heavy sterile neutrino mixing in B8 decay obtained with the Borexino detector”

For HNL decay length scales of the order of the Earth’s diameter, HNL **up-scattering outside the detector** is possible, where a neutrino from the solar or the atmospheric neutrino flux scatters in the Earth and transfers some kinetic energy into the mass of the HNL, which can then later decay inside the detector. For HNL masses below 18 MeV produced from solar neutrinos, (external) limits were derived using the *Borexino* data for purely tau coupling through mass mixing [59] and for all flavor coupling through the dipole portal [60]. At similar decay length scales, the HNL could also be produced directly in the atmosphere, but neither this channel, nor the production anywhere in the Earth from atmospheric neutrinos has been investigated yet.

[59]: Plestid (2021), “Luminous solar neutrinos I: Dipole portals”

[60]: Plestid (2021), “Luminous solar neutrinos II: Mass-mixing portals”

[61]: Coloma et al. (2017), “Double-Cascade Events from New Physics in Icecube”

[62]: Coloma (2019), “Icecube/Deep-Core tests for novel explanations of the MiniBooNE anomaly”

[63]: Atkinson et al. (2022), “Heavy Neutrino Searches through Double-Bang Events at Super-Kamiokande, DUNE, and Hyper-Kamiokande”

[64]: Coloma et al. (2021), “GeV-scale neutrinos: interactions with mesons and DUNE sensitivity”

If the HNL decay lengths are sufficiently short, **production and decay in the detector** can happen and the observation of two vertices could be used to constrain the mixing parameters. In principle, this could be possible with any neutrino flavor produced in the sun or the atmosphere, but so far only theoretical studies have been performed for mass-mixing and dipole-portal couplings for the atmospheric neutrino detectors *IceCube* [61, 62] and *Super-K*, *Hyper-K*, and *Dune* [63, 64]. Due to the high complexity of these experiments, several simplified assumptions were made in the studies, which might not hold in reality, and the results should be taken with caution. For reliable

sensitivity estimates and limits the collaborations should perform their own analyses.

1.3 Atmospheric Neutrinos as Source of Heavy Neutral Leptons

This work focuses on the search for HNLs using atmospheric neutrinos as source for the production and decay inside the IceCube detector. The following sections will give a brief overview of the production of neutrinos in the atmosphere and the oscillations they undergo, before discussing the expected signatures of HNLs in the detector, where they are produced from the incoming neutrinos and subsequently decay.

1.3.1 Production of Neutrinos in the Atmosphere

The analysis performed in this work is based on the sample of neutrinos observed in IceCube DeepCore at energies below 100 GeV. At these energies, the flux exclusively originates in the Earth's atmosphere. Highly relativistic cosmic rays (protons and heavier nuclei [65]) interact in the upper atmosphere, producing showers of secondary particles. Neutrinos are produced in decays of charged pions and kaons (π and K mesons) present in those showers, where the dominant contribution comes from the decay chain

$$\begin{aligned}\pi^\pm &\rightarrow \mu^\pm + \nu_\mu(\bar{\nu}_\mu), \\ \mu^\pm &\rightarrow e^\pm + \bar{\nu}_\mu(\nu_\mu) + \nu_e(\bar{\nu}_e),\end{aligned}\quad (1.22)$$

where muon neutrinos ν_μ and muons μ^\pm are produced in the first decay and both electron and muon neutrinos $\nu_{e/\mu}$ are produced in the second decay. Atmospheric muons, which are also produced in these decays, are the main background component for IceCube DeepCore analyses.

The different atmospheric flux components are shown in Figure 1.5 (left), for a much broader energy range than relevant for this work. Both neutrinos and antineutrino fluxes are shown for electron and muon neutrinos and all fluxes are the directionally averaged expectation calculated at the South Pole. Muon neutrinos are dominating the flux and from Equation 1.22 the naive assumption would be that the ratio between muon and electron neutrinos is

[65]: Tanabashi et al. (2018), "Review of Particle Physics"




Figure 1.5: The atmospheric fluxes of different neutrino flavors as a function of energy (left) and the ratios between muon neutrinos and electron neutrinos as well as the ratios between neutrinos and antineutrinos for both those flavors (right). Results from the calculations performed for the geographic South Pole, taken from [66].

$(\nu_\mu + \bar{\nu}_\mu)/(\nu_e + \bar{\nu}_e) = 2$. This is roughly true at energies below 1 GeV, where all muons decay in flight, but at larger energies muons can reach the detector before decaying, which increases the ratio to approximately 10:1 at around 100 GeV. Additionally, kaon decays start to contribute which also increases the number of muons and muon neutrinos. The increasing ratio can be seen in Figure 1.5 (right), which also shows the ration between neutrinos and antineutrinos for both flavors.

[67]: Fedynitch et al. (2015), "Calculation of conventional and prompt lepton fluxes at very high energy"

Say something about atmospheric neutrino flux uncertainties, based on recent JP/Anatoli papers. (YELLOW)

Charged mesons or tau particles can also be produced in cosmic ray interactions. Their decays lead to the production of tau neutrinos. At the energies relevant for this work however, the resulting tau neutrino flux is negligible as compared to the muon neutrino flux [67] and is not considered in the analysis. This is because both charged mesons and tau particles are much heavier than pions and kaons and therefore their production is suppressed at high energies.

1.3.2 Neutrino Oscillations

Describing neutrinos in their mass state as introduced in Section ?? is crucial to understand their propagation through space and time and to explain neutrino oscillations. Oscillations mean that a neutrino changes from its initial flavor, that it was produced with, to another flavor and back after traveling a certain distance.

The neutrino propagation in vacuum can be expressed by applying a plane wave approach, where the mass eigenstates evolve as

$$|v_k(t)\rangle = e^{-iE_k t/\hbar} |v_k\rangle . \quad (1.23)$$

The energy of the mass eigenstate $|v_k\rangle$ is $E_k = \sqrt{\vec{p}^2 c^2 + m_k^2 c^4}$, with momentum \vec{p} and mass m_k , \hbar is the reduced Planck constant, and c is the speed of light in vacuum. A neutrino is produced as a flavor eigenstate $|v_\alpha\rangle$ in a CC weak interaction, but its propagation happens as the individual mass states it is composed of. The probability of finding the neutrino with initial flavor $|v_\alpha\rangle$ in the flavor state $|v_\beta\rangle$ after the time t is calculated as

$$P_{v_\alpha \rightarrow v_\beta}(t) = |\langle v_\beta | v_\alpha(t) \rangle|^2 , \quad (1.24)$$

[68]: Dirac (1927), "The Quantum Theory of the Emission and Absorption of Radiation"

by applying Fermi's Golden Rule [68], which defines the transition rate from one eigenstate to another by the strength of the coupling between them. This coupling strength is the square of the matrix element and using the fact that the mixing matrix is unitary ($U^{-1} = U^\dagger$) to describe the mass eigenstates as flavor eigenstates, we find the time evolution of the flavor state $|v_\alpha(t)\rangle$, which can be inserted into Equation 1.24 to find the probability as

$$P_{v_\alpha \rightarrow v_\beta}(t) = \sum_{j,k} U_{\beta j}^* U_{\alpha j} U_{\beta k} U_{\alpha k}^* e^{-i(E_k - E_j)t/\hbar} . \quad (1.25)$$

The indices j and k run over the mass eigenstates.

We can approximate the energy as

$$E_k \approx E + \frac{c^4 m_k^2}{2E} \longrightarrow E_k - E_j \approx \frac{c^4 \Delta m_{kj}^2}{2E} , \quad (1.26)$$

for small neutrino masses compared to their kinetic energy. Here, $\Delta m_{kj}^2 = m_k^2 - m_j^2$ is the mass-squared splitting between states k and j . Replacing the time in Equation 1.25 by the distance traveled by relativistic neutrinos $t \approx L/c$ we get

$$P_{\nu_\alpha \rightarrow \nu_\beta}(t) = \delta_{\alpha\beta} - 4 \sum_{j>k} \text{Re}(U_{\beta j}^* U_{\alpha j} U_{\beta k} U_{\alpha k}^*) \sin^2\left(\frac{c^3 \Delta m_{kj}^2}{4E\hbar} L\right) + 2 \sum_{j>k} \text{Im}(U_{\beta j}^* U_{\alpha j} U_{\beta k} U_{\alpha k}^*) \sin^2\left(\frac{c^3 \Delta m_{kj}^2}{4E\hbar} L\right), \quad (1.27)$$

which is called the survival probability if $\alpha = \beta$, and the transition probability if $\alpha \neq \beta$. Once again, this probability is only non-zero if there are neutrino mass eigenstates with masses greater than zero. Additionally, there must be a mass-squared difference Δm^2 and non-zero mixing between the states. Since we assumed propagation in vacuum in Equation 1.23, the transition and survival probabilities correspond to vacuum mixing.

The mixing matrix can be parameterized as [65]

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta_{CP}} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta_{CP}} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (1.28)$$

where $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$ are cosine and sine of the mixing angle θ_{ij} , that defines the strength of the mixing between the mass eigenstates i and j , and δ_{CP} is the neutrino CP-violating phase. Experiments are sensitive to different mixing parameters, depending on the observed energy range, neutrino flavor, and the distance between the source and the detector L , commonly referred to as *baseline*. To be able to resolve oscillations the argument

$$\frac{\Delta m^2 L}{4E} \quad (1.29)$$

should be at the order of 1. This divides experiments into ones that are sensitive to very slow oscillations from $\Delta m_{21}^2 \approx \mathcal{O}(10^{-5} \text{eV}^2)$ and ones that are sensitive to faster oscillations from $\Delta m_{31}^2 \approx \mathcal{O}(10^{-3} \text{eV}^2)$. Relevant for this work are the parameters that can be measured at the earth's surface using atmospheric neutrinos, which are Δm_{31}^2 , θ_{23} , and θ_{13} , because the flux is primarily composed of muon neutrinos and antineutrinos. Applying the parameterization from Equation 1.28 to Equation 1.27 and using the fact that θ_{13} is small and θ_{12} is close to $\pi/4$, the survival probability of muon neutrinos can be approximated as

$$P_{\nu_\mu \rightarrow \nu_\mu} \simeq 1 - 4|U_{\mu 3}|^2(1 - |U_{\mu 3}|^2) \sin^2\left(\frac{\Delta m_{31}^2 L}{4E}\right) \simeq 1 - \sin^2(2\theta_{23}) \sin^2\left(\frac{\Delta m_{31}^2 L}{4E}\right), \quad (1.30)$$

while the tau neutrino appearance probability is

$$P_{\nu_\mu \rightarrow \nu_\tau} \simeq 4|U_{\mu 3}|^2|U_{\tau 3}|^2 \sin^2\left(\frac{\Delta m_{31}^2 L}{4E}\right) \simeq \sin^2(2\theta_{23}) \sin^2\left(\frac{\Delta m_{31}^2 L}{4E}\right). \quad (1.31)$$

[65]: Tanabashi et al. (2018), "Review of Particle Physics"

Parameter	Global Fit
θ_{12} [°]	$33.41^{+0.75}_{-0.72}$
θ_{13} [°]	$8.54^{+0.11}_{-0.12}$
θ_{23} [°]	$49.1^{+1.0}_{-1.3}$
Δm_{21}^2 [10^{-5}eV^2]	$7.41^{+0.21}_{-0.20}$
Δm_{31}^2 [10^{-3}eV^2]	$2.511^{+0.028}_{-0.027}$
δ_{CP} [°]	197^{+42}_{-25}

Table 1.2: Results from the latest global fit of neutrino mixing parameters from [69].

[69]: Esteban et al. (2020), “The fate of hints: updated global analysis of three-flavor neutrino oscillations”

say something about matter effect? (ORANGE)

say something about mass ordering? (ORANGE)

The latest global fit [69] of all the parameters is shown in Table 1.2.

1.3.3 Neutrino Interactions with Nuclei

The neutrino detection principle of IceCube DeepCore is explained in Chapter ?? and relies on the weak interaction processes between neutrinos and the nuclei of the Antarctic glacial ice. At neutrino energies above 5 GeV, the cross-sections are dominated by *deep inelastic scattering (DIS)*, where the neutrino is energetic enough to resolve the underlying structure of the nucleons and interact with one of the composing quarks individually. As a result the nucleon breaks and a shower of hadronic secondary particles is produced. Depending on the type of interaction, the neutrino either remains in the final state for NC interactions or is converted into its charged lepton counterpart for CC interactions. The CC DIS interactions have the form


$$\begin{aligned} \nu_l + N &\rightarrow l^- + X, \\ \bar{\nu}_l + N &\rightarrow l^+ + X, \end{aligned} \quad (1.32)$$

where $\nu_l/\bar{\nu}_l$ and l^-/l^+ are the neutrino/antineutrino and its corresponding lepton/antilepton, and l can be either an electron, muon, or tau. N is the nucleon and X stands for any set of final state hadrons. The NC DIS interactions are

$$\begin{aligned} \nu_l + N &\rightarrow \nu_l + X \text{ and} \\ \bar{\nu}_l + N &\rightarrow \bar{\nu}_l + X. \end{aligned} \quad (1.33)$$

Figure 1.6 shows the Feynman diagrams for both processes DIS interactions

Figure 1.6: Feynman diagrams for deep inelastic scattering of a neutrino with a nucleon via charged-current (left) and neutral current (right) interactions. p_{ν_i} , p_N and p_{ν_f} , p_l , p_N are the input and output four-momenta, while q is the momentum transfer. Taken from [70].



have a roughly linear energy dependent cross-section above ~ 20 GeV and are well measured and easy to theoretically calculate. They are the primary interaction channel for neutrinos detected with IceCube.

At energies below 5 GeV, *quasi-elastic scattering (QE)* and *resonant scattering (RES)* become important. At these energies the neutrinos interact with the approximately point-like nucleons, without breaking them up in the process. RES describes the process of a neutrino scattering off a nucleon producing an excited state of the nucleon in addition to a charged lepton. It is the dominant process at 1.5 GeV to 5 GeV for neutrinos and 1.5 GeV to 8 GeV for antineutrinos. Below 1.5 GeV QE is the main process, where protons are converted to neutrons in antineutrino interactions and vice-versa for neutrino interactions. Additionally, a charged lepton corresponding to the neutrino/antineutrino flavor is produced. The cross-sections of QE and RES scattering processes are not linear in energy and the transition region

from QE/RES to DIS is poorly understood. The total cross-sections and their composition is shown in Figure 1.7. It can be seen that the interaction cross-sections are very small at the order of 10^{-38} cm^2 . This is the reason why very large volume detectors are required to measure atmospheric neutrinos with sufficient statistics to perform precision measurements of their properties. The interaction length of a neutrino with $E_\nu = 10 \text{ GeV}$ is of $\mathcal{O}(10 \times 10^{10} \text{ km})$, for example.




Figure 1.7: Total neutrino (left) and antineutrino (right) per nucleon cross-section divided by neutrino energy plotted against energy. The three main scattering processes quasi-elastic scattering (QE), resonant scattering (RES), and deep-inelastic scattering (DIS) are shown. Taken from [71].

1.3.4 Heavy Neutral Lepton Production and Decay

For the search conducted in this work, both production and decay are assumed to happen inside the detector, therefore probing decay lengths ranges at the scale of the detector size, which is below 1000 m. Since the mixing with the first two generations of leptons is already strongly constrained as was discussed in Section 1.3, only the mixing with the tau neutrino will be considered in the following. Due to the effect of oscillations, described in Section 1.3.2, the initial atmospheric muon neutrino flux provides a sizable tau neutrino flux at the detector.

For a non-zero $|U_{\tau 4}|^2$, the HNL can be produced through **up-scattering in the ice**. An incoming tau neutrinos scatters on an ice nucleus and transfers some of its kinetic energy to the heavy neutrino. The Feynman diagram of this process is shown in Figure ?? . The custom NC cross-sections calculated for this purpose are explained in more detail in Section ?? , but are similar to the SM tau neutrino NC cross-sections, with a reduction scaling with the mixing $|U_{\tau 4}|^2$ and energy dependent reductions, due to kinematic constraints because of the heavy neutrino mass. The scattering process produces a hadronic cascade, which will produce light in the detector.

After a certain distance, the HNL will **decay in the ice**, where the possible decay channels considered in this work are shown in Figure 1.8 and the underlying, explicit calculations are discussed in Section ?? . The decay can be a CC or NC and both purely leptonic and leptonic+mesonic modes are possible. The Feynman diagrams of the decays can be seen in Section ?? . Only the mass range relevant for this work is presented and mixing with $\nu_{e/\mu}$ is assumed to be negligible. Depending on the decay channel, an electromagnetic or a hadronic cascade is produced, while some energy

fix up-scattering feynman diagram and put into the margin (RED)

fix the decay feynman diagrams and put into the margin (RED)

Figure 1.8: Decay widths of the HNL within the mass range considered, calculated based on the results from [64]. Given the existing constraints on $|U_{e4}|^2$ and $|U_{\mu 4}|^2$, we consider that the corresponding decay modes are negligible.





Figure 1.9: Theoretical mean decay length of the HNL for a mass of 0.6 GeV and different mixing values.



1: A particle decay time follows an exponential distribution, with mean lifetime given by the proper lifetime. The proper lifetime is the lifetime in the rest frame of the particle.

is carried away by the invisible neutrino. The decay length of the HNL is defined by its proper lifetime¹, which is given by

$$\tau_{\text{proper}} = \frac{\hbar}{\Gamma_{\text{total}}(m_4) \cdot |U_{\tau 4}|^2} , \quad (1.34)$$

where \hbar is the reduced Planck constant, $\Gamma_{\text{total}}(m_4)$ is the total decay width of the HNL for the given mass, and $|U_{\tau 4}|^2$ is the mixing with the tau neutrino. The total decay width is the sum of the partial decay widths for all possible decay channels. The mean lab frame decay length is then given by

$$L_{\text{decay}} = \gamma v \tau_{\text{proper}} , \quad (1.35)$$

where γ is the Lorentz factor of the HNL, defined by the kinetic energy. This will be further discussed on Section ?? . Figure 1.9 shows the mean decay lengths for an example mass of $m_4 = 0.6$ GeV and several mixing values.

Bibliography

Here are the references in citation order.

- [1] C. N. Yang and R. L. Mills. “Conservation of Isotopic Spin and Isotopic Gauge Invariance”. In: *Physical Review* 96.1 (Oct. 1954), pp. 191–195. doi: [10.1103/PhysRev.96.191](https://doi.org/10.1103/PhysRev.96.191) (cited on page 1).
- [2] S. Weinberg. “A Model of Leptons”. In: *Phys. Rev. Lett.* 19 (21 Nov. 1967), pp. 1264–1266. doi: [10.1103/PhysRevLett.19.1264](https://doi.org/10.1103/PhysRevLett.19.1264) (cited on page 1).
- [3] S. L. Glashow. “Partial-symmetries of weak interactions”. In: *Nuclear Physics* 22.4 (Feb. 1961), pp. 579–588. doi: [10.1016/0029-5582\(61\)90469-2](https://doi.org/10.1016/0029-5582(61)90469-2) (cited on page 1).
- [4] R. Jackiw. “Physical Formulations: Elementary Particle Theory. Relativistic Groups and Analyticity. Proceedings of the eighth Nobel Symposium, Aspenäsgråden, Lerum, Sweden, May 1968. Nils Svartholm, Ed. Interscience (Wiley), New York, and Almqvist and Wiksell, Stockholm, 1969. 400 pp., illus. \$31.75.” In: *Science* 168.3936 (1970), pp. 1196–1197. doi: [10.1126/science.168.3936.1196.b](https://doi.org/10.1126/science.168.3936.1196.b) (cited on page 1).
- [5] P. Higgs. “Broken symmetries, massless particles and gauge fields”. In: *Physics Letters* 12.2 (1964), pp. 132–133. doi: [https://doi.org/10.1016/0031-9163\(64\)91136-9](https://doi.org/10.1016/0031-9163(64)91136-9) (cited on page 1).
- [6] S. Chatrchyan et al. “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC”. In: *Phys. Lett. B* 716 (2012), pp. 30–61. doi: [10.1016/j.physletb.2012.08.021](https://doi.org/10.1016/j.physletb.2012.08.021) (cited on page 1).
- [7] G. Aad et al. “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”. In: *Phys. Lett. B* 716 (2012), pp. 1–29. doi: [10.1016/j.physletb.2012.08.020](https://doi.org/10.1016/j.physletb.2012.08.020) (cited on page 1).
- [8] M. Gell-Mann. “A Schematic Model of Baryons and Mesons”. In: *Resonance* 24 (1964), pp. 923–925 (cited on page 1).
- [9] G. Zweig. “An SU(3) model for strong interaction symmetry and its breaking. Version 2”. In: *DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. VOL. 1. 1964 - 1978*. Ed. by D. B. Lichtenberg and S. P. Rosen. Feb. 1964, pp. 22–101 (cited on page 1).
- [10] D. J. Gross and F. Wilczek. “Ultraviolet Behavior of Non-Abelian Gauge Theories”. In: *PRL* 30.26 (June 1973), pp. 1343–1346. doi: [10.1103/PhysRevLett.30.1343](https://doi.org/10.1103/PhysRevLett.30.1343) (cited on page 1).
- [11] C. Giunti and C. W. Kim. *Fundamentals of Neutrino Physics and Astrophysics*. Oxford University Press, Mar. 2007 (cited on page 1).
- [12] M. D. Schwartz. *Quantum Field Theory and the Standard Model*. Cambridge University Press, 2013 (cited on page 1).
- [13] A. Trettin. “Search for eV-scale sterile neutrinos with IceCube DeepCore”. PhD thesis. Berlin, Germany: Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät, 2023. doi: <https://github.com/atrettin/PhD-Thesis> (cited on pages 3, 6).
- [14] N. Deruelle, J.-P. Uzan, and P. de Forcrand-Millard. *Relativity in Modern Physics*. Oxford University Press, Aug. 2018 (cited on page 4).
- [15] R. L. Workman et al. “Review of Particle Physics”. In: *Progress of Theoretical and Experimental Physics* 2022.8 (Aug. 2022), p. 083C01. doi: [10.1093/ptep/ptac097](https://doi.org/10.1093/ptep/ptac097) (cited on page 4).
- [16] M. Fukugita and T. Yanagida. “Baryogenesis without grand unification”. In: *Physics Letters B* 174.1 (1986), pp. 45–47. doi: [https://doi.org/10.1016/0370-2693\(86\)91126-3](https://doi.org/10.1016/0370-2693(86)91126-3) (cited on page 4).
- [17] R. Davis, D. S. Harmer, and K. C. Hoffman. “Search for Neutrinos from the Sun”. In: *Phys. Rev. Lett.* 20 (21 May 1968), pp. 1205–1209. doi: [10.1103/PhysRevLett.20.1205](https://doi.org/10.1103/PhysRevLett.20.1205) (cited on page 4).

- [18] Y. Fukuda et al. “Evidence for Oscillation of Atmospheric Neutrinos”. In: *Phys. Rev. Lett.* 81 (8 Aug. 1998), pp. 1562–1567. doi: [10.1103/PhysRevLett.81.1562](https://doi.org/10.1103/PhysRevLett.81.1562) (cited on page 4).
- [19] Q. R. Ahmad and other. “Direct Evidence for Neutrino Flavor Transformation from Neutral-Current Interactions in the Sudbury Neutrino Observatory”. In: *Phys. Rev. Lett.* 89 (1 June 2002), p. 011301. doi: [10.1103/PhysRevLett.89.011301](https://doi.org/10.1103/PhysRevLett.89.011301) (cited on page 4).
- [20] S. Alam et al. “Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory”. In: *Phys. Rev. D* 103 (8 Apr. 2021), p. 083533. doi: [10.1103/PhysRevD.103.083533](https://doi.org/10.1103/PhysRevD.103.083533) (cited on page 4).
- [21] N. Aghanim et al. “Planck2018 results: VI. Cosmological parameters”. In: *Astronomy & Astrophysics* 641 (Sept. 2020), A6. doi: [10.1051/0004-6361/201833910](https://doi.org/10.1051/0004-6361/201833910) (cited on page 4).
- [22] M. Aker et al. “Direct neutrino-mass measurement with sub-electronvolt sensitivity”. In: *Nature Phys.* 18.2 (2022), pp. 160–166. doi: [10.1038/s41567-021-01463-1](https://doi.org/10.1038/s41567-021-01463-1) (cited on page 4).
- [23] T. Asaka, S. Blanchet, and M. Shaposhnikov. “The nuMSM, dark matter and neutrino masses”. In: *Phys. Lett. B* 631 (2005), pp. 151–156. doi: [10.1016/j.physletb.2005.09.070](https://doi.org/10.1016/j.physletb.2005.09.070) (cited on page 6).
- [24] T. Asaka and M. Shaposhnikov. “The ν MSM, dark matter and baryon asymmetry of the universe”. In: *Phys. Lett. B* 620 (2005), pp. 17–26. doi: [10.1016/j.physletb.2005.06.020](https://doi.org/10.1016/j.physletb.2005.06.020) (cited on page 6).
- [25] P. Minkowski. “ $\mu \rightarrow e \gamma$ at a rate of one out of 10^9 muon decays?” In: *Physics Letters B* 67.4 (Apr. 1977), pp. 421–428. doi: [10.1016/0370-2693\(77\)90435-X](https://doi.org/10.1016/0370-2693(77)90435-X) (cited on page 6).
- [26] T. Yanagida. “Horizontal Symmetry and Masses of Neutrinos”. In: *Progress of Theoretical Physics* 64.3 (Sept. 1980), pp. 1103–1105. doi: [10.1143/PTP.64.1103](https://doi.org/10.1143/PTP.64.1103) (cited on page 6).
- [27] S. L. Glashow. “The Future of Elementary Particle Physics”. In: *NATO Sci. Ser. B* 61 (1980), p. 687. doi: [10.1007/978-1-4684-7197-7_15](https://doi.org/10.1007/978-1-4684-7197-7_15) (cited on page 6).
- [28] M. Gell-Mann, P. Ramond, and R. Slansky. “Complex Spinors and Unified Theories”. In: *Conf. Proc. C* 790927 (1979), pp. 315–321 (cited on page 6).
- [29] R. N. Mohapatra and G. Senjanovi ć. “Neutrino Mass and Spontaneous Parity Nonconservation”. In: *Phys. Rev. Lett.* 44 (14 Apr. 1980), pp. 912–915. doi: [10.1103/PhysRevLett.44.912](https://doi.org/10.1103/PhysRevLett.44.912) (cited on page 6).
- [30] M. G. Aartsen et al. “eV-Scale Sterile Neutrino Search Using Eight Years of Atmospheric Muon Neutrino Data from the IceCube Neutrino Observatory”. In: *Phys. Rev. Lett.* 125.14 (2020), p. 141801. doi: [10.1103/PhysRevLett.125.141801](https://doi.org/10.1103/PhysRevLett.125.141801) (cited on page 6).
- [31] J.-L. Tastet, O. Ruchayskiy, and I. Timiryasov. “Reinterpreting the ATLAS bounds on heavy neutral leptons in a realistic neutrino oscillation model”. In: *JHEP* 12 (2021), p. 182. doi: [10.1007/JHEP12\(2021\)182](https://doi.org/10.1007/JHEP12(2021)182) (cited on page 6).
- [32] G. Bernardi et al. “Search for Neutrino Decay”. In: *Phys. Lett. B* 166 (1986), pp. 479–483. doi: [10.1016/0370-2693\(86\)91602-3](https://doi.org/10.1016/0370-2693(86)91602-3) (cited on page 7).
- [33] F. Bergsma et al. “A Search for Decays of Heavy Neutrinos”. In: *Phys. Lett. B* 128 (1983). Ed. by J. Tran Thanh Van, p. 361. doi: [10.1016/0370-2693\(83\)90275-7](https://doi.org/10.1016/0370-2693(83)90275-7) (cited on page 7).
- [34] S. Ito et al. “Search for heavy neutrinos in $\pi^+ \rightarrow \mu^+ \nu$ decay and status of lepton universality test in the PIENU experiment”. In: *JPS Conf. Proc.* 33 (2021). Ed. by N. Saito, pp. 011131-1–011131-6. doi: [10.7566/JPSCP.33.011131](https://doi.org/10.7566/JPSCP.33.011131) (cited on page 7).
- [35] A. Aguilar-Arevalo et al. “Improved search for heavy neutrinos in the decay $\pi \rightarrow e \nu$ ”. In: *Phys. Rev. D* 97.7 (2018), p. 072012. doi: [10.1103/PhysRevD.97.072012](https://doi.org/10.1103/PhysRevD.97.072012) (cited on page 7).
- [36] D. A. Bryman and R. Shrock. “Constraints on Sterile Neutrinos in the MeV to GeV Mass Range”. In: *Phys. Rev. D* 100 (2019), p. 073011. doi: [10.1103/PhysRevD.100.073011](https://doi.org/10.1103/PhysRevD.100.073011) (cited on page 7).
- [37] D. Britton et al. “Improved search for massive neutrinos in $\pi \rightarrow e + \nu$ decay”. In: *Physical Review D* 46 (1992) (cited on page 7).
- [38] C. J. Parkinson et al. “Search for heavy neutral lepton production at the NA62 experiment”. In: *PoS EPS-HEP2021* (2022), p. 686. doi: [10.22323/1.398.0686](https://doi.org/10.22323/1.398.0686) (cited on page 7).

- [39] K. Abe et al. “Search for heavy neutrinos with the T2K near detector ND280”. In: *Phys. Rev. D* 100.5 (2019), p. 052006. doi: [10.1103/PhysRevD.100.052006](https://doi.org/10.1103/PhysRevD.100.052006) (cited on page 7).
- [40] P. Abratenko et al. “Search for Heavy Neutral Leptons in Electron-Positron and Neutral-Pion Final States with the MicroBooNE Detector”. In: *Phys. Rev. Lett.* 132.4 (2024), p. 041801. doi: [10.1103/PhysRevLett.132.041801](https://doi.org/10.1103/PhysRevLett.132.041801) (cited on page 7).
- [41] A. Vaitaitis et al. “Search for neutral heavy leptons in a high-energy neutrino beam”. In: *Phys. Rev. Lett.* 83 (1999), pp. 4943–4946. doi: [10.1103/PhysRevLett.83.4943](https://doi.org/10.1103/PhysRevLett.83.4943) (cited on page 7).
- [42] P. Astier et al. “Search for heavy neutrinos mixing with tau neutrinos”. In: *Phys. Lett. B* 506 (2001), pp. 27–38. doi: [10.1016/S0370-2693\(01\)00362-8](https://doi.org/10.1016/S0370-2693(01)00362-8) (cited on page 7).
- [43] J. Orloff, A. N. Rozanov, and C. Santoni. “Limits on the mixing of tau neutrino to heavy neutrinos”. In: *Phys. Lett. B* 550 (2002), pp. 8–15. doi: [10.1016/S0370-2693\(02\)02769-7](https://doi.org/10.1016/S0370-2693(02)02769-7) (cited on page 7).
- [44] I. Boiarska et al. “Blast from the past: constraints from the CHARM experiment on Heavy Neutral Leptons with tau mixing”. In: (July 2021) (cited on page 7).
- [45] R. Barouki, G. Marocco, and S. Sarkar. “Blast from the past II: Constraints on heavy neutral leptons from the BEBC WA66 beam dump experiment”. In: *SciPost Phys.* 13 (2022), p. 118. doi: [10.21468/SciPostPhys.13.5.118](https://doi.org/10.21468/SciPostPhys.13.5.118) (cited on page 7).
- [46] E. Fernandez-Martinez et al. “Effective portals to heavy neutral leptons”. In: *JHEP* 09 (2023), p. 001. doi: [10.1007/JHEP09\(2023\)001](https://doi.org/10.1007/JHEP09(2023)001) (cited on pages 8–10).
- [47] G. Aad et al. “Search for heavy neutral leptons in decays of W bosons produced in 13 TeV pp collisions using prompt and displaced signatures with the ATLAS detector”. In: *JHEP* 10 (2019), p. 265. doi: [10.1007/JHEP10\(2019\)265](https://doi.org/10.1007/JHEP10(2019)265) (cited on page 7).
- [48] A. Tumasyan et al. “Search for long-lived heavy neutral leptons with displaced vertices in proton-proton collisions at $\sqrt{s}=13$ TeV”. In: *JHEP* 07 (2022), p. 081. doi: [10.1007/JHEP07\(2022\)081](https://doi.org/10.1007/JHEP07(2022)081) (cited on page 8).
- [49] A. M. Sirunyan et al. “Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s}=13$ TeV”. In: *Phys. Rev. Lett.* 120.22 (2018), p. 221801. doi: [10.1103/PhysRevLett.120.221801](https://doi.org/10.1103/PhysRevLett.120.221801) (cited on page 8).
- [50] B. Shuve and M. E. Peskin. “Revision of the LHCb Limit on Majorana Neutrinos”. In: *Phys. Rev. D* 94.11 (2016), p. 113007. doi: [10.1103/PhysRevD.94.113007](https://doi.org/10.1103/PhysRevD.94.113007) (cited on page 8).
- [51] R. Aaij et al. “Search for heavy neutral leptons in $W^+ \rightarrow \mu^+ \mu^\pm \text{jet}$ decays”. In: *Eur. Phys. J. C* 81.3 (2021), p. 248. doi: [10.1140/epjc/s10052-021-08973-5](https://doi.org/10.1140/epjc/s10052-021-08973-5) (cited on page 8).
- [52] A. Osipowicz et al. “KATRIN: A Next generation tritium beta decay experiment with sub-eV sensitivity for the electron neutrino mass. Letter of intent”. In: (Sept. 2001) (cited on page 8).
- [53] S. Mertens et al. “A novel detector system for KATRIN to search for keV-scale sterile neutrinos”. In: *Journal of Physics G: Nuclear and Particle Physics* 46.6 (2019), p. 065203. doi: [10.1088/1361-6471/ab12fe](https://doi.org/10.1088/1361-6471/ab12fe) (cited on page 8).
- [54] M. Aker et al. “Search for keV-scale sterile neutrinos with the first KATRIN data”. In: *Eur. Phys. J. C* 83.8 (2023), p. 763. doi: [10.1140/epjc/s10052-023-11818-y](https://doi.org/10.1140/epjc/s10052-023-11818-y) (cited on page 8).
- [55] B. Abi et al. “Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume II: DUNE Physics”. In: (Feb. 2020) (cited on page 8).
- [56] S. Friedrich et al. “Limits on the Existence of sub-MeV Sterile Neutrinos from the Decay of ^7Be in Superconducting Quantum Sensors”. In: *Phys. Rev. Lett.* 126.2 (2021), p. 021803. doi: [10.1103/PhysRevLett.126.021803](https://doi.org/10.1103/PhysRevLett.126.021803) (cited on page 9).
- [57] C. Hagner et al. “Experimental search for the neutrino decay $\nu_3 + \nu_j + e^+ + e^-$ and limits on neutrino mixing”. English. In: 52.3 (1995), pp. 1343–1352. doi: [10.1103/PhysRevD.52.1343](https://doi.org/10.1103/PhysRevD.52.1343) (cited on page 9).
- [58] G. Bellini et al. “New limits on heavy sterile neutrino mixing in B_8 decay obtained with the Borexino detector”. In: *Phys. Rev. D* 88.7 (2013), p. 072010. doi: [10.1103/PhysRevD.88.072010](https://doi.org/10.1103/PhysRevD.88.072010) (cited on page 10).
- [59] R. Plestid. “Luminous solar neutrinos I: Dipole portals”. In: *Phys. Rev. D* 104 (2021), p. 075027. doi: [10.1103/PhysRevD.104.075027](https://doi.org/10.1103/PhysRevD.104.075027) (cited on page 10).

- [60] R. Plestid. “Luminous solar neutrinos II: Mass-mixing portals”. In: *Phys. Rev. D* 104 (2021). [Erratum: *Phys.Rev.D* 105, 099901 (2022)], p. 075028. doi: [10.1103/PhysRevD.104.075028](https://doi.org/10.1103/PhysRevD.104.075028) (cited on page 10).
- [61] P. Coloma et al. “Double-Cascade Events from New Physics in Icecube”. In: *Phys. Rev. Lett.* 119.20 (2017), p. 201804. doi: [10.1103/PhysRevLett.119.201804](https://doi.org/10.1103/PhysRevLett.119.201804) (cited on page 10).
- [62] P. Coloma. “Icecube/DeepCore tests for novel explanations of the MiniBooNE anomaly”. In: *Eur. Phys. J. C* 79.9 (2019), p. 748. doi: [10.1140/epjc/s10052-019-7256-8](https://doi.org/10.1140/epjc/s10052-019-7256-8) (cited on page 10).
- [63] M. Atkinson et al. “Heavy Neutrino Searches through Double-Bang Events at Super-Kamiokande, DUNE, and Hyper-Kamiokande”. In: *JHEP* 04 (2022), p. 174. doi: [10.1007/JHEP04\(2022\)174](https://doi.org/10.1007/JHEP04(2022)174) (cited on page 10).
- [64] P. Coloma et al. “GeV-scale neutrinos: interactions with mesons and DUNE sensitivity”. In: *Eur. Phys. J. C* 81.1 (2021), p. 78. doi: [10.1140/epjc/s10052-021-08861-y](https://doi.org/10.1140/epjc/s10052-021-08861-y) (cited on pages 10, 16).
- [65] M. Tanabashi et al. “Review of Particle Physics”. In: *Phys. Rev. D* 98 (3 Aug. 2018), p. 030001. doi: [10.1103/PhysRevD.98.030001](https://doi.org/10.1103/PhysRevD.98.030001) (cited on pages 11, 13).
- [66] M. Honda et al. “Atmospheric neutrino flux calculation using the NRLMSISE-00 atmospheric model”. In: *Phys. Rev. D* 92 (2 July 2015), p. 023004. doi: [10.1103/PhysRevD.92.023004](https://doi.org/10.1103/PhysRevD.92.023004) (cited on page 11).
- [67] A. Fedynitch et al. “Calculation of conventional and prompt lepton fluxes at very high energy”. In: *European Physical Journal Web of Conferences*. Vol. 99. European Physical Journal Web of Conferences. Aug. 2015, p. 08001. doi: [10.1051/epjconf/20159908001](https://doi.org/10.1051/epjconf/20159908001) (cited on page 12).
- [68] P. A. M. Dirac. “The Quantum Theory of the Emission and Absorption of Radiation”. In: *Proceedings of the Royal Society of London Series A* 114.767 (Mar. 1927), pp. 243–265. doi: [10.1098/rspa.1927.0039](https://doi.org/10.1098/rspa.1927.0039) (cited on page 12).
- [69] I. Esteban et al. “The fate of hints: updated global analysis of three-flavor neutrino oscillations”. In: *JHEP* 09 (2020), p. 178. doi: [10.1007/JHEP09\(2020\)178](https://doi.org/10.1007/JHEP09(2020)178) (cited on pages 13, 14).
- [70] A. Terliuk. “Measurement of atmospheric neutrino oscillations and search for sterile neutrino mixing with IceCube DeepCore”. PhD thesis. Berlin, Germany: Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät, 2018. doi: [10.18452/19304](https://doi.org/10.18452/19304) (cited on page 14).
- [71] J. A. Formaggio and G. P. Zeller. “From eV to EeV: Neutrino cross sections across energy scales”. In: *Rev. Mod. Phys.* 84 (3 Sept. 2012), pp. 1307–1341. doi: [10.1103/RevModPhys.84.1307](https://doi.org/10.1103/RevModPhys.84.1307) (cited on page 15).