

Commutative Algebra

Harry Armitage

October 27, 2020

Contents

0	Introduction	2
1	Noetherian Rings: Definitions and Examples	3
1.1	Ideal Structure	7
1.2	Minimal and Associated Primes	10
1.3	Annihilators and Associated Primes	11
2	Localisation	13
2.1	Modules	15
3	Tensor Products	17

0 Introduction

Commutative Algebra is the study of commutative rings and the spaces on which those rings act, namely modules. It was developed from two key sources: algebraic geometry, and algebraic number theory.

In algebraic geometry we are focused on polynomial rings over a field k , whilst in number theory we are focused on \mathbb{Z} , the ring of rational integers. Much of this work was done by Grothendieck, but the subject goes back much further, at least to Hilbert who wrote a series of papers on polynomial invariant theory in the late nineteenth century.

As an example, take Σ_n , the symmetric group on the set $\{1, 2, \dots, n\}$. Σ_n acts on $k[x_1, \dots, x_n]$ by permuting the variables, so that $(\sigma f)(x_1, \dots, x_n) = f(x_{\sigma^{-1}(1)}, \dots, x_{\sigma^{-1}(n)})$. σ_n acts here via ring automorphisms, and it is then natural to consider the *ring of invariants*, given by $\{f \in k[x] : \sigma f = f \ \forall \sigma \in \Sigma_n\} := S$. S is a ring, *the ring of symmetric polynomials*. We can consider the elementary symmetric functions, which are:

$$\begin{aligned} e_1(x_1, \dots, x_n) &= x_1 + \dots + x_n \\ e_2(x_1, \dots, x_n) &= \sum_{i < j} x_i x_j \\ &\vdots \\ e_n(x_1, \dots, x_n) &= x_1 \dots x_n \end{aligned}$$

In fact, S is generated as a ring by these e_i , and there are canonical maps $k[y_1, \dots, y_n] \rightarrow S$ such that $Y_i \mapsto e_i$, which is a ring isomorphism.

Hilbert showed that S is finitely generated, and moreover for many other groups, not just symmetric groups.

Along the way, he proved four very deep theorems:

- Basis theorem
- Nullstellensatz
- The polynomial nature of the Hilbert function (leading to the beginnings of dimension theory)
- The syzygy theorem (leading to the beginnings of homological theory of polynomial rings)

In 1921 Emmy Noether extracted the key property that made the basis theorem, namely that a commutative ring is *noetherian* if every ideal is finitely generated (there are several equivalent definitions).

Theorem 0.1 (Hilbert's Basis Theorem). *If R is a commutative noetherian ring, then $R[x]$ is also noetherian.*

Corollary 0.2. *If k is a field, then $k[x_1, \dots, x_n]$ is noetherian.*

Noether developed a theory of ideals for noetherian rings, for example the existence of primary decomposition, which generalises factorisation into primes in noetherian rings.

Link between Commutative Algebra and Algebraic Geometry

The starting point for this link is the *fundamental theorem of algebra*, which says that $f \in \mathbb{C}[x]$ is determined up to scalar multiples by its zeros up to multiplicity. Given $f \in \mathbb{C}[x_1, \dots, x_n]$, there is a polynomial function $\mathbb{C}^n \rightarrow \mathbb{C}$ given by $(a_1, \dots, a_n) \mapsto f(a_1, \dots, a_n)$.

Different polynomials will yield different functions, and so $\mathbb{C}[x_1, \dots, x_n]$ can be viewed as a ring of polynomial functions on complex affine n -space.

More specifically, given $I \subseteq \mathbb{C}[x_1, \dots, x_n]$, we can define the *set of common zeros*, $Z(I) = \{(a_1, \dots, a_n) \in \mathbb{C}^n : f(a_1, \dots, a_n) = 0 \ \forall f \in I\}$, called an *(affine) algebraic set*.

Remarks:

- One can replace I by the ideal generated by I , and you get the same algebraic set. Similarly, replacing an ideal by a generating set of the ideal leaves the algebraic set. The basis theorem asserts that any algebraic set is the set of common zeros of some *finite* set of polynomials.
- $\bigcap_j Z(I_j) = Z(\bigcup_j I_j)$, $\bigcup_{j=1}^n Z(I_j) = Z(\prod_{j=1}^n I_j)$, for ideal I_j . If we define a topology on \mathbb{C}^n by calling these algebraic sets the closed sets, we get the *Zariski topology*, which is a rather coarser topology on \mathbb{C}^n than the usual topology.
- For $S \subseteq \mathbb{C}^n$, we can define $I(S) = \{f \in \mathbb{C}[x_1, \dots, x_n] : f(a_1, \dots, a_n) = 0 \ \forall (a_1, \dots, a_n) \in S\}$. This is an *ideal* of $\mathbb{C}[x_1, \dots, x_n]$, and it is *radical*, i.e. $f^r \in I(S) \implies f \in I(S)$. The Nullstellensatz is a family of results asserting that the correspondence

$$\begin{aligned} I &\mapsto Z(I) \\ I(S) &\leftrightarrow S \end{aligned}$$

gives a bijection between the radical ideals in $\mathbb{C}[x_1, \dots, x_n]$ and the algebraic subsets of \mathbb{C}^n . In particular, the maximal ideals of $\mathbb{C}[x_1, \dots, x_n]$ correspond to points in \mathbb{C}^n .

Dimension

A large portion of the course deals with the dimension of rings. We can define it in three main ways:

- The maximal length of a chain of prime ideals.
- In a geometric context in terms of growth rates.
- The transcendence degree of a field of fractions.

For commutative rings, all three give the same answer. There is in fact a fourth method, using homological algebra, which in the case of “nice” noetherian rings also gives the same answer.

Most of this theory dates back to 1920-1950. Rings of dimension 0 are called *artinian* rings, and in dimension 1 there are special properties which are important in number theory, particularly in the study of algebraic curves.

1 Noetherian Rings: Definitions and Examples

Throughout this section, R is a commutative ring with a 1.

Lemma 1.1. *Let M be a (left) R -module. The following are equivalent:*

1. *All submodules of M (including M itself) are finitely generated.*
2. *The ascending chain condition (ACC) holds: there are no strictly increasing infinite chains of submodules.*
3. *The maximum condition of submodules holds: any nonempty set S of submodules of M has a maximal element L , i.e. $L \subseteq L', L' \in S \implies L = L'$.*

Proof.

1. \implies 2. Suppose there is a strictly increasing chain $N_1 \subsetneq N_2 \subsetneq \dots$, and let $N = \bigcup_{i=1}^{\infty} N_i$. By 1 N is finitely generated, say by m_1, \dots, m_r . Each m_i lies in some N_{n_i} . Then let $n = \max_i n_i$, so that $m_i \in N_n$. Then $N_n = M$, contradicting strict ascent.

2. \implies 3. Assume ACC. Pick $M_1 \in S$. If it is the maximal member then we're done. If not, there is $M_2 \supsetneq M_1$. If M_2 is maximal, then we're done, otherwise there is some $M_3 \supsetneq M_2$, and so on. By ACC this process terminates, and we get a maximal element.

3. \implies 1. Let $N \triangleleft M$, and let S be the collection of all finitely generated submodules of N . Then $S \neq \emptyset$ since it contains the 0 submodule. So S contains a maximal member, say L . We then claim $N = L$. If $x \in N$ then $L + Rx \in S$, and by maximality of L , $x \in L$. \square

Definition 1.2. *An R -module satisfying 1, 2, 3 is **noetherian**.*

Lemma 1.3. *Let $N \triangleleft M$. Then M is noetherian if and only if N and M/N are noetherian.*

Proof.

\implies Let M be noetherian, so that all its submodules are finitely generated. This property is inherited by N . Also, the submodules of M/N are all of the form Q/N with $Q \triangleleft M$ containing N . If M is noetherian, then Q is finitely generated, say by x_1, \dots, x_r . Then $x_1 + N, \dots, x_r + N$ generates Q/N .

\Leftarrow Let $N, M/N$ be noetherian, and let $L_1 \subset L_2 \subset L_3 \subset \dots$ be a strictly increasing chain of submodules of M . Set $Q_i/N = (L_i + N)/N$, and $N_i = L_i \cap N$. These give ascending chains of submodules of M/N and N respectively. By ACC there are r, s with $Q_i/N = Q_r/N$ for $i \geq r$, $N_i = N_s$ for $i \geq s$. Let $k = \max\{r, s\}$. Then we claim $L_i = L_k$ for $i \geq k$. Pick $\ell \in L_i$, $i \geq k$. Then $\ell + N \in Q_k/N$, and so there is some $\ell' \in L_k$ such that $\ell - \ell' \in N \cap L_i = N \cap L_k$. So $\ell \in L_k$, and the claim is proved. Hence our original ascending chain was not strictly increasing, \nmid . \square

Lemma 1.4. 1. *If M, N are R -modules, then $M \oplus N$ is noetherian iff M and N are noetherian.*

2. *If M_1, \dots, M_n are R -modules then $M_1 \oplus \dots \oplus M_n$ is noetherian iff each M_i is noetherian.*
3. *If M is noetherian then every homomorphic image of M is noetherian.*
4. *Suppose M can be expressed as a sum of finitely many submodules (not necessarily as a direct sum) $M = M_1 + \dots + M_n$. Then M is noetherian iff each M_i is.*

Proof. 1. $M \cong N/N$, so this follows by 1.3.

2. Apply 1 and induction on n .

3. If $\theta : M \rightarrow N$ then $\text{im } \theta \cong M/\ker \theta$, so apply 1.3.

4. The forwards direction follows as $M_i \triangleleft M$. For the reverse, there is a map from $M_1 \oplus \dots \oplus M_n \rightarrow M$, $(m_1, \dots, m_n) \mapsto m_1 + \dots + m_n$, and then apply 2 and 3.

□

Definition 1.5. A ring R is **noetherian** if it is noetherian as a (left) R -module

Remark: Submodules of R as an R -module are the same as ideals of R as a ring, and so the ACC for modules gives us the ACC for ideals.

Lemma 1.6. Let R be a noetherian ring. Then any finitely generated R -module M is noetherian.

Proof. Suppose $M = Rm_1 + \dots + Rm_n$. There exist R -module epimorphisms:

$$\begin{aligned} R &\rightarrow Rm_i \\ r &\mapsto rm_i \end{aligned}$$

R is noetherian, so Rm_i is as the homomorphic image of R . Then, by 1.4 (4), so is M . □

Theorem 1.7 (Hilbert Basis Theorem). Let R be a noetherian ring. Then the polynomial ring $R[x]$ is noetherian.

Proof. We show that every ideal of $R[x]$ is finitely generated. Let I be an ideal. We define $I(n) = \{f \in I : \deg f \leq n\}$. Then $I(n) \neq \emptyset$ as $0 \in I(n)$, and $I(0) \subseteq I(1) \subseteq I(2) \subseteq \dots$

Let $R(n) = \{\text{Coefficient of } x^n \text{ in } f : f \in I(n)\} \subseteq R$. We claim $R(n) \triangleleft R$, and $R(n) \subseteq R(n+1)$.

To see this, suppose $a, b \in R(n)$. Then there are polynomials $f(x) = ax^n + \dots$, $g(x) = bx^n + \dots$ in I , where \dots indicates lower order terms. Since $I \triangleleft R$, $f \pm g \in I$, $rf \in I$ for all $r \in R$, and $xf \in I$.

Hence $a \pm b \in R(n)$, $ra \in R(n)$, and $a \in R(n+1)$, and the claim is proved.

So then we have a chain $R(0) \subseteq R(1) \subseteq R(2) \subseteq \dots$ terminates, so we may say $R(n) = R(N) \forall n \geq N$. Each of $R(0), \dots, R(N)$ is a finitely generated ideal of R , say $R(j) = (a_{j,1}, \dots, a_{j,k_j})$.

Then by definition of $R(j)$, we may take polynomials $f_{j,1}, \dots, f_{j,k_j}$ in $I(j)$ which have the $a_{j,i}$ as their leading coefficients.

Clearly $I \supseteq (f_{j,k} : 0 \leq j \leq N, 1 \leq k \leq k_j) =: J$ - it remains to show that equality holds, then we will have found a finite generating set of I . So pick $f \in I$, then we claim $f \in J$, and prove this by induction on the degree of f .

If $\deg f = 0$, then $f(x) = a$, say. But then $a \in R(0)$, and so $a = \sum_i r_i a_{0,i}$ for some $r_i \in R$. Since $f_{0,i}$ has $a_{0,i}$ as its leading coefficient and has degree zero, $f_{0,i}(x) = a_{0,i}$, and $f = \sum_i r_i f_{0,i} \in J$.

If instead $\deg f = n$, with $0 < n \leq N$, and the claim holds for all g with $\deg g < n$, then write $f(x) = ax^n + \dots$ $a \in R(n)$ then by definition, so $a = \sum_i r_{n,i} a_{n,i}$ for some $r_{n,i} \in R$. Then define $g(x) = f(x) - \sum_i r_{n,i} f_{n,i}(x)$. $g(x)$ has degree $\leq n$, and the coefficient of x^n is $a - a = 0$, hence $\deg g < n$. Since $f_{n,i} \in I$, we have $g \in I$, and hence by induction $g \in J$. But $f_{n,i} \in J$ as well, so $f \in J$.

Finally if $\deg f = n$, with $n > N$, and the claim holds for all g with $\deg g < n$, again write $f(x) = ax^n + \dots$. Then $a \in R(n) = R(N)$, so $a = \sum r_{N,j} a_{N,j}$ for $r_{N,j} \in R$. We may then define $g(x) = f(x) - \sum_i x^{n-N} r_{N,i} f_{N,i}(x)$, and use the same argument as in the previous paragraph to deduce that $f \in J$.

Hence $I \subseteq J$, and so $I = J$ and I is finitely generated. But I was an arbitrary ideal of $R[x]$, so $R[x]$ is noetherian. \square

In practice, one uses *Gröbner bases* for ideals - these are generating sets with extra properties that make algorithms more efficient.

Examples:

- Fields are noetherian.
- Principle Ideal Domains (PIDs) are noetherian.
- $\{q \in Q : q = \frac{m}{n}, m, n \in \mathbb{Z}, p \nmid n \text{ for some fixed prime } p\}$, an example of a *localisation* of \mathbb{Z} . All localisations of noetherian rings are noetherian - we will see this later.
- $k[x_1, x_2, \dots]$ is not noetherian: $(x_1) \subsetneq (x_1, x_2) \subsetneq \dots$ is an infinite strictly increasing chain.
- $k[x_1, x_2, \dots, x_n]$ is noetherian - this follows by induction using the Hilbert basis theorem.
- $\mathbb{Z}[x_1, x_2, \dots, x_n]$ is noetherian, so any finitely generated commutative ring is noetherian: if R is generated by r_1, \dots, r_n , then there is an epimorphism $\mathbb{Z}[x_1, \dots, x_n] \rightarrow R$ given by $x_i \mapsto r_i$, and R is the homomorphic image of a noetherian ring.
- If A is a free abelian group, write $\mathbb{Z}A$ for its group algebra, which is the set of formal linear combinations of elements of A , i.e. terms of the form $\sum_{\alpha \in A} \lambda_\alpha \alpha$ where $\lambda_\alpha \in \mathbb{Z}$ and only finitely many of the λ_α are nonzero.
If A is generated as a group by g_1, \dots, g_n , then its group algebra is generated as a ring by $g_1, g_1^{-1}, \dots, g_n, g_n^{-1}$.
- $k[[x]]$, the ring of formal power series with coefficients in k , is noetherian.

There are also some non-commutative examples that are both left and right noetherian:

- Enveloping algebras of a finite dimensional Lie algebra.
- Iwasawa algebras of compact p -adic groups.

Theorem 1.8. *If R is noetherian, then $R[[x]]$ is noetherian.*

Proof 1. As in 1.7, consider $R(n)$ = the set of trailing coefficients a_n , for elements $a_n x^n + \text{higher order terms}$, and mimic the proof. This is on example sheet 1. \square

We will give a second proof, which uses

Theorem 1.9 (Cohen's Theorem). *If every prime ideal in a ring R is finitely generated, then R is noetherian.*

Proof. If R is not noetherian, then there is a family of non-finitely generated ideals. Call it \mathcal{S} . By assumption, $\mathcal{S} \neq \emptyset$. Partially order \mathcal{S} by inclusion.

Suppose $I_1 \subseteq I_2 \subseteq \dots$ is a chain of non-finitely generated ideals. Then we claim $\bigcup_i I_i$ is also non-finitely generated.

If it were, say by (a_1, \dots, a_k) , then $a_i \in I_{n(i)}$ for some finite integer $n(i)$, and so, if $N = \max\{n(i) : 1 \leq i \leq k\}$, N is also finite and $a_i \in I_N$ for all i . But then $I_N = I_n$ for all $n \geq N$, and in particular I_N is finitely generated \nmid .

So \mathcal{S} has upper bounds to its chains, and so we may apply Zorn's lemma to get a maximal element of \mathcal{S} , say I , so that I is not finitely generated but any ideal containing I is finitely generated.

We now claim I must be prime. Suppose $a \notin I, b \notin I$, but $ab \in I$. Then $I + (a) \supsetneq I$, so $I + (a)$ is finitely generated, say by $i_1 + r_1a, \dots, i_n + r_na$. Define $J = \{s \in R : sa \in I\} \supseteq I + (b) \supsetneq I$. Again, J is finitely generated.

Take $t \in I \subset I + (a)$, so $t = u_1(i_1 + r_1a) + \dots + u_n(i_n + r_na)$ for some $u_i \in R$. So $t = u_1i_1 + \dots + u_ni_n + (u_1r_1 + \dots + u_nr_n)a \in (i_1) + (i_2) + \dots + (i_n) + Ja$.

Hence $I \subseteq (i_1) + \dots + (i_n) + Ja$, so $I = (i_1) + \dots + (i_n) + Ja$, so I is finitely generated \nmid .

So I must be prime, but then by our hypothesis I is still finitely generated \nmid . So R must be noetherian. \square

We will also use the following lemma:

Lemma 1.10. *Let P be a prime ideal of $R[[x]]$ and $\theta : R[[x]] \rightarrow R, x \mapsto 0$. Then P is finitely generated if and only if $\theta(P)$ is a finitely generated ideal of R .*

Proof. Clearly if P is finitely generated then $\theta(P)$ is.

Conversely, suppose $\theta(P) = Ra_1 + \dots + Ra_n$.

If $x \in P$, then $P = (a_1, \dots, a_n, x)$.

This is immediate - if $g \in P$, $g = a + \text{higher order terms}$. Now $a \in (a_1, \dots, a_n)$, so $g = \sum_i r_i a_i + xg'$ as required.

If $x \notin P$, then let f_1, \dots, f_n be power series with constant terms a_1, \dots, a_n respectively. Then $P = (f_1, \dots, f_n)$.

Take $g \in P$, say $g = b + \text{higher terms}$, with b the constant term. Then $b = \sum b_i a_i$, so $g - \sum b_i f_i = g_1 x$ for some g_1 . Note that $g_1 x \in P$, P is prime, and $x \notin P$, so $g_1 \in P$. Similarly, $g_1 = \sum c_i f_i + g_2 x$, and $g_2 \in P$. Continuing, we get $h_1, \dots, h_n \in R[[x]]$, where $h_i = b_i + c_i x + \dots$ with $g = h_1 f_1 + \dots + h_n f_n$. \square

We are now ready to give the second proof the R noetherian implies $R[[x]]$ noetherian:

Proof 2. Suppose P is a prime ideal of $R[[x]]$. Then P is finitely generated iff $\theta(P)$ is. But R is noetherian, so $\theta(P)$ is finitely generated, so P was finitely generated. Then we apply Cohen's theorem to get $R[[x]]$ noetherian. \square

1.1 Ideal Structure

Here, we assume R is a commutative ring with a 1, not necessarily noetherian.

Lemma 1.11. *The set $N(R)$ of all nilpotent¹ elements of R is an ideal, and $R/N(R)$ has no nonzero nilpotent elements.*

Proof. If $x \in N(R)$, then $x^m = 0$ for some m . Hence $(rx)^m = 0$ for all $r \in R$, and so $rx \in N(R)$.

If $x, y \in N(R)$, then $x^n = 0, y^m = 0$ for some n, m . Then $(x+y)^{n+m-1}$ expands to give terms $\lambda x^s y^t$ where $s+t = n+m-1$. So either $s \geq n$ or $t \geq m$, so all the terms are zero, and $x+y \in N(R)$.

¹An element x of a ring is called nilpotent if there is some integer m such that $x^m = 0$.

So $N(R) \triangleleft R$.

Finally, if $s \in R/N(R)$ then $s = x + N(R)$. Note that $s^n = x^n + N(R)$ for all n . If $x + N(R)$ is nilpotent then $(x + N(R))^m = N(R)$ for some m , and hence $x^m \in N(R)$. So x^m is nilpotent, and $(x^m)^n = x^{mn} = 0$ for some n . But then x is nilpotent, so $x + N(R) = 0 + N(R)$. \square

Definition 1.12. $N(R)$ is called the **nilradical** of R .

Theorem 1.13 (Krull). $N(R)$ is the intersection of all prime ideals of R .

Proof. Let $I = \bigcap_{P \text{ prime}} P$. If $x \in R$ is nilpotent then $x^n = 0 \in P \forall P$. So $x \in P \forall P \implies x \in I$, so $N(R) \subseteq I$.

Suppose x is not nilpotent. Let \mathcal{S} be the family of ideals J such that for $n > 0$, $x^n \notin J$. Then $(0) \in \mathcal{S}$, so $\mathcal{S} \neq \emptyset$, and a union of a chain of ideals in \mathcal{S} is also in \mathcal{S} . We apply Zorn's lemma to get a maximal element J_1 .

We claim J_1 is prime - suppose $yz \in J_1$, but $y, z \notin J_1$. So the ideals $J_1 + Ry, J_1 + Rz$ strictly contain J_1 , and so $x^m \in J_1 + Ry$ and $x^n \in J_1 + Rz$. But then $x^{m+n} \in J_1 + Ryz = J_1$. ∇

So J_1 is prime, so contains I , and hence $x \notin I$, so $I \supsetneq N(R)$. Thus $I = N(R)$. \square

Definition 1.14. The **radical** \sqrt{I} of an ideal I is defined by $\{r \in R : \exists k \in \mathbb{N} \text{ s.t. } r^k \in I\}$.

Note that $\sqrt{I}/I = N(R/I)$, and $\sqrt{I} = \bigcap_{\text{prime } P \supset I} P$. We say an ideal I is radical if $I = \sqrt{I}$.

Definition 1.15. The **Jacobson radical** $J(R)$ of R is the intersection of all the maximal ideals of R (so $N(R) \subseteq J(R)$).

Theorem 1.16 (Nakayama's Lemma). If M is a finitely generated R -module with $MJ = M$, where $J = J(R)$, then $M = 0$.

Proof. ² If $M \neq 0$ and is a finitely generated R -module, then by Zorn's lemma there are maximal proper submodules.

Take M_1 maximal in M . Then M/M_1 is irreducible (or simple), hence generated by $m + M_1$ say.

Then, considering the map $R \rightarrow M/M_1; r \mapsto rm + M_1$, which is an R -module homomorphism with kernel a maximal ideal, we see that $M/M_1 \cong R/I$, where I is a maximal ideal of R , so $MI \leq M_1$.

Finally, $J \leq I$, then $MJ \leq MI \leq M_1 \leq M$, so if $M \neq 0$, $MJ \leq M$. \square

For a commutative ring R , $N(R) \leq J(R)$. These need not be equal - for example, take $R = \{\frac{m}{n} \in \mathbb{Q} : p \nmid n\} = \mathbb{Z}_{(p)}$. This has unique maximal ideal $P = \{\frac{m}{n} \in \mathbb{Q} : p|m, p \nmid n\}$. It is an integral domain, so has no nonzero nilpotent elements, so $N(R) = (0)$, and $J(R) = P$.

For rings $R = k[x_1, \dots, x_n]/I$ with k algebraically closed and I any ideal, we do have $N(R) = J(R)$ - this is the Nullstellensatz - see later on.

Example: A commutative ring is **artinian** if it doesn't contain an infinite strictly descending chain of ideals (or equivalently if every nonempty set of ideals has a minimal member). An

²Note - this is not the usual Atiyah-Macdonald proof, but this one can be adapted to the case of non-commutative rings.

R -module is **artinian** if it satisfies the analogous properties for submodules. As an exercise (on the first example sheet), prove that artinian rings are noetherian.

For example, $\mathbb{Z}/p\mathbb{Z}, k[x]/(f)$. $k[x]$ is not artinian ($(x) > (x^2) > \dots$).

Recall that I is prime if and only if one following three equivalent properties holds:

$$\begin{aligned} ab \in I &\implies a \in I \text{ or } b \in I \\ R/I &\text{ is an integral domain} \\ I_1 I_2 \subseteq I &\implies I_1 \subseteq I \text{ or } I_2 \subseteq I \end{aligned}$$

Claim: $J(R) = N(R)$ for artinian rings R

This follows if we can show that R artinian \implies every prime ideal is maximal.

Proof. Let P be prime, $x \notin P$. By the descending chain condition, $(x) \supseteq (x^2) \subseteq \dots$ is not strict, so $(x^n) = (x^{n+1}) = \dots$ for some n . Hence $x^n = yx^{n+1}$ for some y . Then $x^n(1 - xy) = 0 \in P$. But $x^n \notin P$, and P is prime, so $1 - xy \in P$. Thus $y + P$ is the inverse of $x + P$ in R/P , and so R/P is a field, and P is maximal. \square

Lemma 1.17 (Artin-Tate). *Suppose we have commutative rings $R \leq S \leq T$. Suppose R is noetherian and T is generated as a ring by R and finitely many elements t_1, \dots, t_n . Suppose that T is a finitely generated S -module. Then S is generated by R and finitely many elements as an R -algebra.*

Proof. T is generated by $x_1, \dots, x_m \in T$ as an S -module, so $T = Sx_1 + \dots + Sx_m$. Then:

$$t_i = \sum_j s_{ij} x_j, \quad s_{ij} \in S \quad (1)$$

$$x_i x_j = \sum_k s_{ijk} x_k, \quad s_{ijk} \in S \quad (2)$$

Let S_0 be the ring generated by R , the s_{ij} and the s_{ijk} , so that $R \leq S_0 \leq S$.

Any element of T is polynomial in the t_i with coefficients in R . In (1), (2), each element is a linear combination of the x_j with coefficients in S_0 . Thus T is a finitely generated S_0 -module. But S_0 is noetherian, being generated as a ring by R and finitely many elements. T is noetherian as an S_0 -module, and S is an S_0 -submodule of T , hence is finitely generated as an S_0 -module.

But S_0 is generated by R and finitely many elements, so S is generated by R and finitely many elements. \square

Lemma 1.18 (Zariski). *Let k be a field, and R a finitely generated k -algebra. If R itself is a field, then it is a finite algebraic extension of k , i.e. a finitely generated k -space.*

Proof. Suppose R is generated by k and x_1, \dots, x_n , and is a field. If R is not a finite algebraic extension over k , then we can reorder the x_1, \dots, x_n so that x_1, \dots, x_m are algebraically independent, i.e. the ring generated by k and x_1, \dots, x_m is a polynomial algebra $k[x_1, \dots, x_m]$, and x_{m+1}, \dots, x_n are algebraic over the field of fractions $F = k(x_1, \dots, x_m)$. Because R is not finite algebraic over k , $m \geq 1$.

Hence R is a finite algebraic extension over F , and R is a finitely generated F -module, (i.e. vector space). Apply Artin-Tate (1.17) for $k \leq F \leq R$, it follows that F is a finitely generated k -algebra by k and q_1, \dots, q_t say, with each $q_i = f_i/g_i$, where $f_i, g_i \in k[x_1, \dots, x_m]$, $g_i \neq 0$.

Now there is a polynomial h which is prime to each of the g_i s, e.g. $g_1 \dots g_m + 1$, and the element $1/h$ cannot be in the ring generated by k and q_1, \dots, q_t . This is a contradiction, and hence $m = 0$, and R was indeed algebraic over k . \square

Theorem 1.19 (Weak Nullstellensatz). *Let k be a field, T a finitely generated k -algebra. Let P be a maximal ideal of T . Then T/P is a finite algebraic extension of k . In particular, if k is algebraically closed and T is the polynomial algebra, then the maximal ideals are of the form $(x_1 - a_1, \dots, x_n - a_n)$.*

Proof. See later. \square

Theorem 1.20 (Strong Nullstellensatz). *Let k be an algebraically closed field, and R a finitely generated k -algebra. Then $N(R) = J(R)$. Thus, if I is a radical ideal of $k[x_1, \dots, x_n]$ and $R = k[x_1, \dots, x_n]/I$, then the intersection of the maximal ideals of R is 0.*

Furthermore, any radical ideal is the intersection of the maximal ideals containing it.

Proof. Deferred until chapter 2. \square

Proof of 1.19. Let P be the maximal ideal of the finitely generated k -algebra T . Put $R = T/P$. By Zariski's lemma, T/P over k is a finite algebraic extension. If k is closed, then $k = T/P$. Set $\pi : T \rightarrow k$ with kernel P .

We then claim that $\ker \pi = (x_1 - \pi(x_1), \dots, x_n - \pi(x_n))$.

Now π fixes elements of k , so the RHS is in the kernel. Conversely, $T/(x_1 - \pi(x_1), \dots, x_n - \pi(x_n))$ is a 1-dimensional k -space, so the kernel is contained in the RHS, and so they are equal.

Recall the bijection proposed earlier between radical ideals in $\mathbb{C}[x]$ and affine algebraic sets in \mathbb{C}^n .

Rephrase this by defining $Q_{(a_1, \dots, a_n)} = (x_1 - a_1, \dots, x_n - a_n)$. We claim there is a bijection:

$$\begin{aligned} \{\text{radical ideals}\} &\leftrightarrow \{\text{algebraic subsets}\} \\ I &\mapsto \{(a_1, \dots, a_n) : I \subseteq Q_{(a_1, \dots, a_n)}\} \\ \bigcap_{(a_1, \dots, a_n) \in S} Q_{(a_1, \dots, a_n)} &\leftarrow S \end{aligned}$$

\square

1.2 Minimal and Associated Primes

Lemma 1.21. *If R is noetherian, then any ideal I contains a power of its radical \sqrt{I} . In particular, $N(R)$ is nilpotent, i.e. $N(R)^m = (0)$ for some m , as $N(R) = \sqrt{(0)}$.*

Proof. Suppose $x_1, \dots, x_m \in \sqrt{I}$ generate \sqrt{I} as an ideal. Then $x_i^{n_i} \in I$ for some n_i . Then, if n is sufficiently large (e.g. $n \geq \sum (n_i - 1) + 1$). Then \sqrt{I}^n is generated by $x_1^{r_1}, \dots, x_m^{r_m}$ with $\sum r_i = n$. We must thus have some $r_i \geq n_i$, and so $\sqrt{I}^n \subseteq I$. \square

Lemma 1.22. *If R is noetherian, then a radical ideal is the intersection of finitely many prime ideals.*

Proof. Suppose not for contradiction, and take a maximal element I from the set of radical ideals not of this form (using Zorn's lemma). We then claim that I is prime, yielding a contradiction.

If not, there is $J_1, J_2 \not\subseteq I$ with $J_1 J_2 \subseteq I$. If necessary, replace J_i by $J_i + I$, we can assume $I \subsetneq J_1, J_2$.

Then by the maximality of I , $\sqrt{J_1} = Q_1 \cap \dots \cap Q_m$; $\sqrt{J_2} = Q'_1 \cap \dots \cap Q'_n$ as prime intersections.

Set $J = \sqrt{J_1} \cap \sqrt{J_2} = Q_1 \cap \dots \cap Q_m \cap Q'_1 \cap \dots \cap Q'_n$. So $J^{n_1} \subseteq J_1, J^{n_2} \subseteq J_2$ for some n_1, n_2 . Hence $J^{n_1+n_2} \subseteq J_1 J_2 \subseteq I$. But I is radical, so $J \subseteq I$. Now all Q_i, Q'_j contain I , so $J \supseteq I$. Thus $J = I$. \square

Now suppose by the previous lemma that any radical ideal $\sqrt{I} = P_1 \cap \dots \cap P_m$ is an intersection of finitely many primes. We can remove P_i from the list if it contains any of the others, so wlog we may assume that $P_i \not\subseteq P_j$ for any $i \neq j$. If P is prime with $\sqrt{I} \subseteq P$, then $P_1 \dots P_m \subseteq \bigcap_i P_i = \sqrt{I} \subseteq P$, and so some $P_i \subseteq P$.

Definition 1.23. The **minimal primes** P over an ideal I of a noetherian ring are those such that, if P' is prime with $I \subseteq P' \subseteq P$, then $P' = P$.

Clearly the P_i mentioned above are minimal primes over I . In fact:

Lemma 1.24. Let I be an ideal in a noetherian ring. Then \sqrt{I} is the intersection of the minimal primes over I , and I contains a finite product of the minimal primes over I .

Proof. Each minimal prime over I contains \sqrt{I} . So the primes minimal over I are precisely the minimal ones over \sqrt{I} . We know \sqrt{I} is the intersection of these, and thus their product lies in \sqrt{I} , and 1.21 gives the last part. \square

Example: The Nullstellensatz bijection between radical ideals of $\mathbb{C}[x_1, \dots, x_n]$ and algebraic subsets of \mathbb{C}^n .

Suppose (a_1, \dots, a_n) is a common zero of all $f \in I$, a radical ideal. Then $I \subseteq (x_1 - a_1, \dots, x_n - a_n)$. This latter ideal is maximal as it is the kernel of $\mathbb{C}[x_1, \dots, x_n] \rightarrow \mathbb{C}; x_i \mapsto a_i$.

Now consider

$$\bigcap_{\substack{(a_1, \dots, a_n) \\ \text{common zeros} \\ \text{of all } f \in I}} (x_1 - a_1, \dots, x_n - a_n)$$

This ideal is radical, and the bijection in the Nullstellensatz implies that this radical ideal is the same as I . Thus I is an intersection of maximal ideals, and moreover all maximal ideals are of the form $(x_1 - a_1, \dots, x_n - a_n)$. Also, for any ideal J_1 of $\mathbb{C}[x_1, \dots, x_n]$, we have $N(\mathbb{C}[x_1, \dots, x_n]/J_1) = J(\mathbb{C}[x_1, \dots, x_n]/J_1)$.

1.3 Annihilators and Associated Primes

Definition 1.25. Let M be a finitely generated R -module, where R is noetherian. The **annihilator** of m , $\text{ann}(m) = \{r \in R : rm = 0\}$. A prime ideal P is an **associated prime** of M if it is the annihilator of an element of M .

We call the set of associated primes $\text{Ass}(M)$.

For example, $\text{Ass}(R/P) = \{P\}$ for P prime.

Definition 1.26. A submodule N of M is **p-primary** (or just **primary**) if $\text{Ass}(M/N) = \{p\}$ for a prime ideal p . An ideal is **p-primary** if I is p -primary as a submodule of R .

Lemma 1.27. If $\text{ann}(M) = P$ for a prime ideal P , then $P \in \text{Ass}(M)$.

Proof. Suppose that M is generated by m_1, \dots, m_k . Let $I_j = \text{ann}(m_j)$. Then the product $\prod I_j$ annihilates each m_j , so $\prod I_j \leq \text{ann}(M) = P$. So $I_j = P$ for some j as P prime, and so $P \in \text{Ass}(M)$. \square

Lemma 1.28. Let Q be maximal amongst all annihilators of nonzero elements. Then Q is a prime ideal and so $Q \in \text{Ass}(M)$.

Proof. Let $Q = \text{ann}(m)$ and $r_1 r_2 \in Q, r_2 \notin Q$. We show that $r_1 \in Q$.

Now $r_1 r_2 \in Q \implies r_1 r_2 m = 0$, so $r_1 \in \text{ann}(r_2 m)$.

And $r_2 \notin Q \implies r_2 m \neq 0$. But $Q \leq \text{ann}(r_2 m)$, and hence Q and r_2 lie in $\text{ann}(r_2 m)$. By maximality, $Q = \text{ann}(r_2 m)$, and so $r_1 \in Q$. \square

Lemma 1.29. For finitely generated nonzero R -module M , where R is noetherian, there is a chain

$$0 \leq M_1 \leq M_2 \leq \dots \leq M_t = M$$

of submodules with $M_i/M_{i-1} \cong R/P_i$ for some prime ideal P_i .

Proof. By 1.28, there is $0 \neq m_1 \in M$ with $\text{ann}(m_1)$ prime, say P_1 . Set $M_1 = Rm_1$. Hence $M_1 \cong R/P_1$. Repeat for M/M_1 to find $M_2/M_1 \cong R/P_2$ for some prime P_2 . Continue - the noetherian property forces the process to terminate. \square

Lemma 1.30. $N \subseteq M \implies \text{Ass}(M) \subseteq \text{Ass}(N) \cup \text{Ass}(M/N)$.

Proof. Take $P \in \text{Ass}(M)$, so that $P = \text{ann}(m)$ for some $m \in M$, and P is prime.

Let $M_1 = Rm = R/P$. For any $0 \neq m_1 \in M_1$, we have $\text{ann}(m_1) = P$, since P is prime. If $M_1 \cap N \neq 0$, then there is $x \in M_1 \cap N$ with $\text{ann}(x) = P$, and so $P \in \text{Ass}(N)$. Otherwise, $M_1 \cap N = 0$, and the image of M_1 in M/N is isomorphic to R/P , and hence $P \in \text{Ass}(M/N)$. \square

Lemma 1.31. $\text{Ass}(M)$ is finite for any finitely generated R -module, where R is noetherian.

Proof. Apply 1.30 inductively to the chain in 1.29 recalling that $\text{Ass}(R/P_i) = \{P_i\}$. We thus conclude $\text{Ass}(M) \subset \{P_1, \dots, P_t\}$ is finite. \square

Proposition 1.32. Each minimal prime over an ideal I is an associated prime, i.e.:

$$\{\text{minimal primes over } I\} \subseteq \text{Ass}(R/I)$$

Proof. By 1.24, there is a product of minimal primes over I , possibly with repetitions, contained in I , say $p_1^{s_1} \dots p_n^{s_n} \leq I$ with $p_i \neq p_j$ for $i \neq j$.

Let $J = \text{ann}(\underbrace{(p_2^{s_2} \dots p_n^{s_n} + I)}_M / I)$. Now $J \geq p_1^{s_1}$, and also $J p_2^{s_2} \dots p_n^{s_n} \leq I \leq p_1$. Since p_1 prime, we have $J \leq p_1$, and so $J \neq R \implies M \neq 0$.

By 1.29, there is a chain of submodules in M , say $0 \leq M_1 \leq \dots \leq M_t = M$ such that each factor is isomorphic to R/q_j for some primes q_j .

But $p_1^{s_1}$ annihilates M , and hence each M_j/M_{j-1} , and the primeness of q_j ensures that $p_1 \leq q_j$ for each j . Not all of the $q_j \geq p_1$ since $\prod q_j \leq J \leq p_1$, and hence some $q_j \leq p_1$, so $q_j = p_1$.

Now pick j minimal such that $q_j = p_1$. Then $\prod_{k < j} q_k \not\leq p_1$. We show that $p_1 \in \text{Ass}(M)$.

Take $x \in M_j \setminus M_{j-1}$. If $j = 1$, then $\text{ann}(x) = p_1$, and so $p_1 \in \text{Ass}(R/I)$. If $j > 1$, take $r \in (\prod_{k < j} q_k) \setminus p_1$. Note that $r(sx) = 0$ for any $s \in p_1 = q_j$. So $s(rx) = 0$, so $p_1 \leq \text{ann}(rx)$. However, $rx \notin M_{j-1}$ since $M_j/M_{j-1} = R/q_j = R/p_1$.

So $\text{ann}(rx) \subseteq p_1$, and hence is equal to, and we've shown that $p_1 \in \text{Ass}(M) \subseteq \text{Ass}(R/I)$. \square

Example 1.33. The converse is false. An example where $p \in \text{Ass}(R/I)$ with p is not minimal over I is as follows:

Take $R = k[x, y]$, $p = (x, y) > q = (x)$, and $I = pq = (x^2, xy)$, so that $\sqrt{I} = (x) = q$.

Then $\text{Ass}(R/I) = \{p, q\}$. The only minimal prime over I is q , since $\sqrt{I} = q$. Now I is not primary as there are two primes in $\text{Ass}(R/I)$. However, we can write $I = (x^2, xy, y^2) \cap (x)$, with $(x^2, xy, y^2) = (x, y)^2$, is p -primary, and (x) is q -primary. This is an example of **primary decomposition**:

Definition 1.34. Let M be a finitely generated R -module with R noetherian, and $N \subset M$ a submodule. Then there are submodules N_1, \dots, N_s of M containing N such that N_i is p_i -primary with p_i distinct, and $N = \bigcap_{i=1}^s N_i$, so that $M/N \hookrightarrow \bigoplus_i M/N_i$.

The primary decomposition is not necessarily unique, although in §4 of Atiyah-MacDonald proves two uniqueness theorems for finitely generated modules over noetherian rings:

- The p_i occurring in the primary decomposition are unique, and are precisely $\text{Ass}(M/N)$.
- If the p_j are minimal among all occurring p_i s, then the corresponding N_j are unique. If p_j are not minimal (which we call **embedded**), then the N_j can vary.

In 1.33, q is minimal and p is embedded. Hence the ideal (x) is unique and $\text{Ass}(R/I) = \{p, q\}$.

2 Localisation

As always, all rings are commutative with a 1.

Let S be a **multiplicatively closed** subset of R - i.e. S is closed under multiplication and $1 \in S$. Define a relation on $R \times S$ via

$$(r_1, s_1) \equiv (r_2, s_2) \iff (r_1 s_2 - r_2 s_1)x = 0 \text{ for some } x \in S$$

This is reflexive, symmetric, and transitive. Reflexivity and symmetry are easy - for transitivity, if $(r_1, s_1) \equiv (r_2, s_2) \equiv (r_3, s_3)$, then we have $(r_1 s_2 - r_2 s_1)x = 0 = (r_2 s_3 - r_3 s_2)y$.

Then $s_3 y \cdot \text{LHS} - s_1 x \cdot \text{RHS} = 0$, and so $(r_1 s_3 - r_3 s_1)s_2 x y = 0$, and $s_2 x y \in S$ since S is multiplicatively closed.

Denote the equivalence class of (r_1, s_1) as $\frac{r_1}{s_1}$, and denote the set of equivalence classes by $S^{-1}R$. Then $S^{-1}R$ is a ring, and there is a ring homomorphism $\theta : R \rightarrow S^{-1}R$ via $r \mapsto \frac{r}{1}$.

We also have the following universal property:

Lemma 2.1. *Let $\varphi : R \rightarrow T$ be a ring homomorphism with $\varphi(s)$ a unit in T for all $s \in S$. Then there is a unique ring homomorphism $\alpha : S^{-1}R \rightarrow T$ such that following diagram commutes:*

$$\begin{array}{ccc} R & \xrightarrow{\theta} & S^{-1}R \\ & \searrow \varphi & \swarrow \alpha \\ & T & \end{array}$$

Proof. For uniqueness, suppose that α exists, then $\alpha : S^{-1}R \rightarrow T$ such that $\alpha\theta = \varphi$.

Then

$$\begin{aligned} \alpha(r/1) &= \alpha(\theta(r)) = \varphi(r) \quad \forall r \in R \\ \alpha(1/s) &= \alpha((s/1)^{-1}) = (\alpha(s/1))^{-1} = \varphi(s)^{-1} \quad \forall s \in S \end{aligned}$$

Thus $\alpha(r/s) = \varphi(r)\varphi(s)^{-1}$, and α is uniquely determined.

For existence, let $\alpha(r/s) = \varphi(r)\varphi(s)^{-1}$. We need to show this is well defined.

Suppose $(r_1/s_1) = (r_2/s_2)$. Then there is $x \in S$ with $(r_1s_2 - r_2s_1)x = 0$.

So $(\varphi(r_1)\varphi(s_2) - \varphi(r_2)\varphi(s_1))\varphi(x) = 0$.

Since $\varphi(x)$ is a unit, we must have $\varphi(r_1)\varphi(s_2) = \varphi(r_2)\varphi(s_1)$, and thus $\alpha(r_1/s_1) = \alpha(r_2/s_2)$. \square

Examples:

- The field of fractions of an integral domain R . Put $S = R \setminus \{0\}$.
- $S^{-1}R = (0) \iff 0 \in S$.
- If $I \triangleleft R$, then take $S = 1 + I$ is multiplicatively closed.
- If p is a prime ideal, then let $S = R \setminus p$, and this is multiplicatively closed since p is prime. In this case, we write R_p for $S^{-1}R$ in this case.

This process of passing from R to R_p is called **localisation at p** . The elements $\frac{r}{s}$ with $r \in p$ form an ideal of R_p , and in fact is the unique maximal ideal in R_p - if $\frac{r}{s}$ is such that $r \notin p$, then $r \in S$, and hence $\frac{s}{r} \in R_p$ is its inverse.

Definition 2.2. *A ring with a unique maximal ideal is called **local**.*³

Examples:

- $\mathbb{R} = \mathbb{Z}, p = (p)$ for p prime. Then $R_p = \{\frac{m}{n} : p \nmid n\} \subset \mathbb{Q}$, with unique maximal ideal $\{\frac{m}{n} : p \mid m, p \nmid n\}$
- $R = k[x_1, \dots, x_n], p = (x_1 - \alpha_1, \dots, x_n - \alpha_n)$. Then $R_p \leq k(x_1, \dots, x_n)$, and is those functions that are defined at $(\alpha_1, \dots, \alpha_n) \in k^n$. The unique maximal ideal consists of those rational functions which are zero at $(\alpha_1, \dots, \alpha_n)$.

³Some authors require local rings to be noetherian - we will be explicit here when we need this.

2.1 Modules

Given a (left) R -module M , for a multiplicatively closed set $S \subseteq R$, define a relation on $M \times S$ by $(m_1, s_1) \equiv (m_2, s_2) \iff x(m_1 s_2 - m_2 s_1) = 0$ for some $x \in S$. Again, this is an equivalence relation, with $\frac{m}{s}$ denoting an equivalence class, and we write for the set of equivalence classes $S^{-1}M$. Now $S^{-1}M$ is an $S^{-1}R$ -module.

Write M_p in the case where $S = R \setminus p$ for some prime ideal p . If $\theta : M_1 \rightarrow M_2$ is an R -homomorphism, then $S^{-1}\theta : S^{-1}M_1 \rightarrow S^{-1}M_2$ is an $S^{-1}R$ -homomorphism, defined by $S^{-1}\theta : \frac{m_1}{s} \mapsto \frac{\theta(m_1)}{s}$.

If $\varphi : M_2 \rightarrow M_3$, then $S^{-1}(\varphi \circ \theta) = S^{-1}\varphi \circ S^{-1}\theta$.

A sequence of R -modules

$$M_0 \rightarrow M_1 \rightarrow M_2 \rightarrow \dots \xrightarrow{\theta} M_i \xrightarrow{\varphi} \dots \rightarrow M_t$$

is *exact* at M_i if $\text{im } \theta = \ker \varphi$. A *short exact sequence* is one of the form

$$0 \rightarrow M_1 \xrightarrow{\theta} M \xrightarrow{\varphi} M_2 \rightarrow 0$$

with exactness at M_1, M, M_2 . Then θ is injective, φ is surjective, and $\text{im } \theta = \ker \varphi$.

Lemma 2.3. If M_1, M, M_2 are R -modules and $M_1 \xrightarrow{\theta} M \xrightarrow{\varphi} M_2$ is exact at M , then

$$S^{-1}M_1 \xrightarrow{S^{-1}\theta} S^{-1}M \xrightarrow{S^{-1}\varphi} S^{-1}M_2$$

is exact at $S^{-1}M$.

Proof. Since $\ker \varphi = \text{im } \theta$, we have $\varphi \circ \theta = 0$. So $(S^{-1}\varphi) \circ (S^{-1}\theta) = S^{-1}(\varphi \circ \theta) = S^{-1}(0) = 0$, and hence $\text{im } S^{-1}\theta \leq \ker S^{-1}\varphi$.

Now suppose that $\frac{m}{s} \in \ker S^{-1}\varphi \subseteq S^{-1}M$. Then $\frac{\varphi(m)}{s} = 0$ in $S^{-1}M_2$ and there is some $t \in S$ with $t\varphi(m) = 0$ in M_2 . But $t\varphi(m) = \varphi(tm)$, since φ is a R -module homomorphism. Thus $tm \in \ker \varphi = \text{im } \theta$, and so $tm = \theta(m_1)$ for some $m_1 \in M_1$.

Hence in $S^{-1}M$, $\frac{m}{s} = \frac{\theta(m_1)}{ts} = (S^{-1}\theta) \left(\frac{m_1}{ts} \right) \in \text{im } S^{-1}\theta$, and so $\ker S^{-1}\varphi \leq \text{im } S^{-1}\theta$.

Thus $\ker S^{-1}\varphi = \text{im } S^{-1}\theta$, and the sequence is exact. \square

Lemma 2.4. Let $N \leq M$. Then $S^{-1}(M/N) \cong S^{-1}M/S^{-1}N$.

Proof. Apply 2.3 to the short exact sequence $0 \rightarrow N \xhookrightarrow{\iota} M \xrightarrow{\epsilon} M/N \rightarrow 0$ to get that

$$0 \rightarrow S^{-1}N \xhookrightarrow{S^{-1}\iota} S^{-1}M \xrightarrow{S^{-1}\epsilon} S^{-1}(M/N) \rightarrow 0$$

is a short exact sequence. Hence $S^{-1}(M/N) \cong S^{-1}M/S^{-1}N$. \square

If R is a ring with a multiplicatively closed subset S , and $I \trianglelefteq R$, then $S^{-1}I$ is an ideal of $S^{-1}R$. If $N \subset M$, then $S^{-1}N$ can be regarded as a submodule of $S^{-1}M$.

Lemma 2.5.

1. Every ideal in $S^{-1}R$ is of the form $S^{-1}I$ for some $I \trianglelefteq R$.
2. The prime ideals of $S^{-1}R$ are in one-to-one correspondence with the prime ideals of R that don't intersect S .

Proof.

1. Let $J \trianglelefteq S^{-1}R$, and $I = \{r \in R : \frac{r}{1} \in J\} \subseteq R$. Then if $\frac{r}{s} \in J$, we have $\frac{r}{1} \in J$, and hence $r \in I$, so $J \subseteq S^{-1}I$.

Conversely, if $r \in I$ then $\frac{r}{1} \in J$, and so $\frac{r}{s} \in J$ for all s , so $S^{-1}I \subseteq J$.

2. Let q be a prime ideal in $S^{-1}R$. Set $p := \{r \in R : \frac{r}{1} \in q\}$.

p is prime: if $xy \in p$, then $\frac{xy}{1} \in q$, and so either $\frac{x}{1}$ or $\frac{y}{1}$ is in q , so $x \in p$ or $y \in p$.

$p \cap S = \emptyset$: Suppose $r \in S \cap p$. Then $\frac{r}{1} \in q$ and $\frac{1}{r} \in q$, and so $\frac{1}{r} \cdot \frac{r}{1} = \frac{1}{1} \in q$.

Conversely, if $\frac{r}{s}, \frac{x}{y} \in S^{-1}p$, then $\frac{rx}{sy} \in S^{-1}p$. So $z(rx) \in p$ for some $z \in S$. Hence $rx \in p$ since p is prime and $z \notin p$. Thus $r \in p$ or $x \in p$, and so $\frac{r}{s} \in S^{-1}p$ or $\frac{x}{y} \in S^{-1}p$.

□

Lemma 2.6. If R is noetherian, then so is $S^{-1}R$.

Proof. Any chain of ideals in $S^{-1}R$ is of the form $J_1 \leq J_2 \leq \dots = S^{-1}I_1 \leq S^{-1}I_2 \leq \dots$

Note that, by the construction given in the proof above, if $S^{-1}I_1 \leq S^{-1}I_2$, then we must have $I_1 \leq I_2$, as $r \in I_1 \implies \frac{r}{1} \in S^{-1}I_1 \implies \frac{r}{1} \in S^{-1}I_2 \implies r \in I_2$.

Then we have a chain of ideals in R given by $I_1 \leq I_2 \leq \dots$, which must terminate as R is noetherian. Then $I_t = I_{t+k}$ for some t and all $k \in \mathbb{N}$, and thus $J_t = J_{t+k}$ for some t and all $k \in \mathbb{N}$, and so our original sequence terminated. □

Definition 2.7. A property \mathcal{P} of a ring R (or R -module M) is **local** if R (or M) has \mathcal{P} if and only if R_p (or M_p) has \mathcal{P} for every prime ideal $p \subseteq R$.

We have just shown that “being noetherian” is a local property. We will now go on to prove that another property is local:

Lemma 2.8. The following are all equivalent:

1. $M = 0$
2. $M_p = 0$ for all prime p
3. $M_q = 0$ for all maximal q

Proof. 1. \implies 2. \implies 3. is immediate. Now we show 3. \implies 1.

Suppose $M_q = 0$ for all maximal q , but $M \neq 0$. Then take $0 \neq m \in M$. The annihilator of m is a proper ideal of R , and so it is contained in a maximal ideal q by Zorn. Consider $\frac{m}{1} \in M_q$. By assumption, $M_q = 0$, and hence $\frac{m}{1} = 0$. So $sm = 0$ for some $s \in S := R \setminus q$. But we already had q containing the annihilator of m , and so we have a contradiction. □

So “being 0” is a local property!

Lemma 2.9. Let $\varphi : M \rightarrow N$ be a R -module homomorphism. Then the following are equivalent:

1. φ is injective.
2. $\varphi_p : M_p \rightarrow N_p$ is injective for all primes p of R .
3. $\varphi_q : M_q \rightarrow N_q$ is injective for all maximal q of R .

Proof. 1. \implies 2. by exactness of localisation, and 2. \implies 3. is clear.

For 3. \implies 1., let $M_1 = \ker \varphi$. Then $0 \rightarrow M_1 \rightarrow M \xrightarrow{\varphi} N \rightarrow 0$ is exact at M_1 and M . So 2.3 gives us $0 \rightarrow (M_1)_q \rightarrow M_q \xrightarrow{\varphi_q} N_q \rightarrow 0$ is exact at $(M_1)_q$ and M_q for maximal q , and hence $(M_1)_q = \ker \varphi_q = 0$, and so $(M_1)_q = 0$. Hence $M_1 = 0$ by 2.8. \square

Lemma 2.10. Let p be a prime of R and S be a multiplicatively closed subset of R , with $S \cap p = \emptyset$. By 2.5, we know $S^{-1}p$ is a prime of $S^{-1}R$. In fact, we have:

$$(S^{-1}R)_{S^{-1}p} \cong R_p$$

In particular, if q is prime with $p \leq q$, then $(R_q)_{p_q} \cong R_p$, by taking $S = R \setminus q$.

Proof. On example sheet 2. \square

3 Tensor Products

Let R be a commutative ring with a 1, and L, M, N, T all R -modules.

Definition 3.1. A function $\varphi : M \times N \rightarrow L$ is R -bilinear if

- $\varphi(r_1m_1 + r_2m_2, n) = r_1\varphi(m_1, n) + r_2\varphi(m_2, n)$
- $\varphi(m, r_1n_1 + r_2n_2) = r_1\varphi(m, n_1) + r_2\varphi(m, n_2)$

The idea of this section is reduce the study of bilinear maps to the study of linear maps.

If $\varphi : M \times N \rightarrow T$ is bilinear and $\theta : T \rightarrow L$ is linear, then the composition $\theta \circ \varphi$ is bilinear. So composition with φ gives a well-defined function φ^* from

$$\{R\text{-module maps } T \rightarrow L\} \rightarrow \{\text{bilinear maps } M \times N \rightarrow L\}$$

We say φ is **universal** if φ^* is a 1-1 correspondence for all L . If this happens, the study of bilinear maps $M \times N \rightarrow L$ is reduced to the study of linear maps $T \rightarrow L$.

Lemma 3.2.

1. Given R -modules M, N , there is an R -module T and a universal map $\varphi : M \times N \rightarrow T$
2. Given two such maps $\varphi_1 : M \times N \rightarrow T_1, \varphi_2 : M \times N \rightarrow T_2$, there is a unique isomorphism $\beta : T_1 \rightarrow T_2$ with $\varphi_2 = \beta \circ \varphi_1$.

Proof.

1. Let F be the free R -module on generators $e_{(m,n)}$, indexed by pairs $m, n \in M \times N$. Then let X be the R -submodule generated by all elements of the forms:

$$\begin{aligned} e_{(r_1 m_1 + r_2 m_2, n)} - r_1 e_{(m_1, n)} - r_2 e_{(m_2, n)} \\ e_{(m, r_1 n_1 + r_2 n_2)} - r_1 e_{(m, n_1)} - r_2 e_{(m, n_2)} \end{aligned}$$

Now set $T = F/X$, and write $m \otimes n$ for the image of the basis element $e_{(m,n)}$ in T . Then T is generated as an R -module by elements of the form $m \otimes n$, and

$$\begin{aligned} (r_1 m_1 + r_2 m_2) \otimes n &= r_1 (m_1 \otimes n) + r_2 (m_2 \otimes n) \\ m \otimes (r_1 n_1 + r_2 n_2) &= r_1 (m \otimes n_1) + r_2 (m \otimes n_2) \end{aligned}$$

Define $\varphi : M \times N \rightarrow T; (m, n) \mapsto m \otimes n$.

Then any map $\alpha : M \times N \rightarrow L$ extends to an R -module map $\bar{\alpha} : F \rightarrow L; e_{(m,n)} \mapsto \alpha(m, n)$. If α is bilinear, then $\bar{\alpha}$ is zero on all the generators of X , hence on all of X . So $\bar{\alpha}$ induces an R -module map $\alpha' : T \rightarrow L$ with $\alpha'(m \otimes n) = \alpha(m, n)$, and α' is uniquely defined by this.

2. Suppose there are universal maps $\varphi_i : M \times N \rightarrow T_i$ for $i = 1, 2$. Since φ_1 is universal, there is a unique R -module map $\beta_1 : T_1 \rightarrow T_2$ with $\varphi_2 = \beta_1 \circ \varphi_1$. Similarly, there is unique $\beta_2 : T_2 \rightarrow T_1; \varphi_1 = \beta_2 \circ \varphi_2$.

Then $(\beta_2 \circ \beta_1) \circ \varphi_1 = \beta_2 \circ \varphi_2 = \varphi_1 = \text{id} \circ \varphi_1$. But φ^* is a bijection, and hence $\beta_2 \circ \beta_1 = \text{id}_{T_1}$, and similarly $\beta_1 \circ \beta_2 = \text{id}_{T_2}$. Hence β_1 is the required isomorphism.

□

Definition 3.3. T in the above proof is written $M \otimes_R N$, the **tensor product** of M and N over R . We often drop the subscript R if it's clear what ring we're working over.

Note: not all elements of $M \otimes_R N$ are of the form $m \otimes n$. A generated element is a finite sum of $\sum_{i=1}^r m_i \otimes n_i$.

If $R = k$ is a field, and M, N are finite dimensional k -vector spaces, of dimension s, t respectively, then $((a_1, \dots, a_s), (b_1, \dots, b_t)) \mapsto (a_i b_j)_{\substack{1 \leq i \leq s \\ 1 \leq j \leq t}}$ is a universal map, and $M \otimes_k N = k^{st}$.

We may also define the tensor product over non-commutative rings, where M is a right R -module and N is a left R -module. Then $M \otimes N$ is only an abelian group, and not necessarily an R -module. If M is an (R, S) -bimodule, and N an (S, T) -bimodule, then $M \otimes N$ is an (R, T) -bimodule.

As an simple example, you can check that $\mathbb{Z}/r\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/s\mathbb{Z} \cong \mathbb{Z}/\text{gcd}(r, s)\mathbb{Z}$.

It's also worth noting that we can do the same with trilinear maps $L \times M \times N \rightarrow T$, giving us $L \otimes M \otimes N$.

Lemma 3.4. *There are unique isomorphisms:*

1. $M \otimes N \rightarrow N \otimes M; m \otimes n \mapsto n \otimes m$.
2. $(M \otimes N) \otimes L \rightarrow M \otimes (N \otimes L) \rightarrow M \otimes N \otimes L; (m \otimes n) \otimes \ell \mapsto m \otimes (n \otimes \ell) \mapsto m \otimes n \otimes \ell$
3. $(M \oplus N) \otimes L \rightarrow (M \otimes L) \oplus (N \otimes L); (m, n) \otimes \ell \mapsto ((m \otimes \ell), (n \otimes \ell))$
4. $R \otimes_R M \rightarrow M; r \otimes m \mapsto rm$

Proof.

1. The map $M \times N \rightarrow N \otimes M; (m, n) \rightarrow n \otimes m$ is bilinear. So by universality there is a map $M \otimes N \rightarrow N \otimes M; m \otimes n \mapsto n \otimes m$, which clearly has an inverse.
2. Exercise on sheet 2.
3. There is a bilinear map $\phi : (M \oplus N) \times L \rightarrow (M \otimes L) \oplus (N \otimes L); ((m, n), \ell) \mapsto (m \otimes \ell, n \otimes \ell)$.

Using the universal property, we have a unique linear map as described in the statement, so we need to produce an inverse, and then it is an isomorphism. We have the following diagram:

$$\begin{array}{ccccc}
 & & (M \otimes L) \oplus (N \otimes L) & & \\
 & \swarrow \pi_1 & & \searrow \pi_2 & \\
 M \otimes L & & & & N \otimes L \\
 & \searrow \psi_1 & & \swarrow \psi_2 & \\
 & & (M \oplus N) \otimes L & &
 \end{array}$$

Where π_1, π_2 are the projections, and ψ_1, ψ_2 are the obvious linear inclusions.

Set $\psi = \psi_1 \pi_1 + \psi_2 \pi_2$ to give the linear map:

$$\begin{aligned}
 \psi : (M \otimes L) \oplus (N \otimes L) &\rightarrow (M \oplus N) \otimes L \\
 ((m \otimes \ell_1), (n \otimes \ell_2)) &\mapsto (m, 0) \otimes \ell_1 + (0, n) \otimes \ell_2
 \end{aligned}$$

Then ψ is the required inverse of ϕ .

4. Left as an exercise. See Atiyah-Macdonald 2.14.

□

Example: $\text{Hom}(M \otimes N, L) \cong \text{Hom}(M, \text{Hom}(N, L))$. Given a bilinear map $\varphi : M \times N \rightarrow L$, we get $\theta : M \rightarrow \text{Hom}(N, L)$ where $m \mapsto (\theta_m : N \rightarrow L, n \mapsto \varphi(m, n))$.

Conversely, given a linear map $\theta : M \rightarrow \text{Hom}(N, L)$, we get a bilinear map $M \times N \rightarrow L, (m, n) \mapsto \theta(m)(n)$. Hence we have a 1-1 correspondence

$$\{\text{bilinear maps } M \times N \rightarrow L\} \leftrightarrow \{\text{linear maps } M \rightarrow \text{Hom}(N, L)\}$$

But the LHS corresponds to the linear maps $M \otimes N \rightarrow L$.