

# Global optimization method with dual Lipschitz constant estimates for problems with non-convex constraints

Roman Strongin, Konstantin Barkalov, Semen Bevzuk

*Lobachevsky State University of Nizhni Novgorod, Nizhni Novgorod, Russia*

---

## Abstract

In the present work, the constrained global optimization problems, in which the functions are of the “black box” type and satisfy the Lipschitz condition, are considered. The algorithms for solving the problems of this class require the use of the adequate estimates of the a priori unknown Lipschitz constants for the problem functions. A novel approach presented in this paper is based on a simultaneous use of two estimates of the Lipschitz constant: an overestimated and the underestimated ones. The upper estimate provides the global convergence whereas the lower one reduces the number of trials necessary to find the global optimizer with required precision. At that, the considered algorithm of solving the constrained problems doesn’t use the ideas of the penalty function method; each constraint of the problem is accounted for separately. The convergence conditions of the proposed algorithm are formulated according to corresponding theorem. The results of the numerical experiments on the series of the multiextremal problems with the non-convex constraints demonstrating the efficiency of the proposed scheme of dual Lipschitz constant estimates are presented.

*Keywords:* Global optimization, Multiextremal problems, Non-convex constraints, Lipschitz constant estimates

---

## 1. Introduction

## 2. Index Method of Accounting for the Constraints and Dimensionality Reduction

$$\min \{ \varphi(y) : y \in D, g_j(y) \leq 0, 1 \leq j \leq m \}, \quad (1)$$

$$D = \{ y \in R^N : a_i \leq y_i \leq b_i, 1 \leq i \leq N \}. \quad (2)$$

$$|g_j(y') - g_j(y'')| \leq L_j \|y' - y''\|, y', y'' \in D, 1 \leq j \leq m+1. \quad (3)$$

$$\varphi(y_\epsilon) = \min \{ \varphi(y) : y \in D, g_j(y) \leq -\epsilon_j, 1 \leq j \leq m \}, \quad (4)$$

$$Y_\epsilon = \{ y \in D, g_j(y) \leq 0, 1 \leq j \leq m, \varphi(y) \leq \varphi(y_\epsilon) \}. \quad (5)$$

$$D_1 = D, D_{j+1} = \{ y \in D_j : g_j(y) \leq 0 \}, 1 \leq j \leq m. \quad (6)$$

$$\varphi(y(x^*)) = \min \{ \varphi(y(x)) : x \in [0, 1], g_j(y(x)) \leq 0, 1 \leq j \leq m \}, \quad (7)$$

$$|g_j(y(x')) - g_j(y(x''))| \leq K_j |x' - x''|^{1/N}, x', x'' \in [0, 1], 1 \leq j \leq m+1,$$

$$Q_1 = [0, 1], Q_{j+1} = \{ x \in Q_j : g_j(y(x)) \leq 0 \}, 1 \leq j \leq m. \quad (8)$$

$$g_j(y(x)) \leq 0, 1 \leq j \leq \nu - 1, g_\nu(y(x)) > 0,$$

$$x^k, z^k = g_\nu(y(x^k)), \nu = \nu(x^k), \quad (9)$$

$$\psi(x^*) = \min \{ \psi(x) : x \in [0, 1] \}, \quad (10)$$

$$\psi(x) = \frac{g_\nu(y(x))}{K_\nu} - \begin{cases} 0, & \nu \leq m, \\ \frac{\varphi^*}{K_{m+1}}, & \nu = m+1. \end{cases} \quad (11)$$

$$\psi(x) = \frac{\varphi(y(x)) - \varphi^*}{K_{m+1}}$$

### 3. Index Algorithm of Global Search

$$0 = x_0 < x_1 < \dots < x_k < x_{k+1} = 1,$$

$$I_\nu = \{i : 1 \leq i \leq k, \nu = \nu(x_i)\}, \quad 1 \leq \nu \leq m+1,$$

$$M = \max \{ \nu = \nu(x_i), \quad 1 \leq i \leq k \}.$$

$$\mu_\nu = \max \left\{ \frac{|z_i - z_j|}{(x_i - x_j)^{1/N}}, \quad i, j \in I_\nu, \quad i > j \right\}, \quad (12)$$

$$z_\nu^* = \begin{cases} -\epsilon_\nu, & \nu < M, \\ \min \{ z_i : i \in I_\nu \}, & \nu = M, \end{cases} \quad (13)$$

$$R(i) = \Delta_i + \frac{(z_i - z_{i-1})^2}{r_\nu^2 \mu_\nu^2 \Delta_i} - 2 \frac{z_i + z_{i-1} - 2z_\nu^*}{r_\nu \mu_\nu}, \quad \nu = \nu(x_i) = \nu(x_{i-1}), \quad (14)$$

$$R(i) = 2\Delta_i - 4 \frac{z_i - z_\nu^*}{r_\nu \mu_\nu}, \quad \nu = \nu(x_i) > \nu(x_{i-1}), \quad (15)$$

$$R(i) = 2\Delta_i - 4 \frac{z_{i-1} - z_\nu^*}{r_\nu \mu_\nu}, \quad \nu = \nu(x_{i-1}) > \nu(x_i), \quad (16)$$

$$R(t) = \max \{ R(i) : 1 \leq i \leq k+1 \}. \quad (17)$$

$$x^{k+1} = \frac{x_t + x_{t-1}}{2}, \quad \nu(x_{t-1}) \neq \nu(x_t).$$

$$x^{k+1} = \frac{x_t + x_{t-1}}{2} + \frac{\text{sign}(z_t - z_{t-1})}{2r_\nu} \left[ \frac{|z_t - z_{t-1}|}{\mu_\nu} \right]^N, \quad \nu = \nu(x_{t-1}) = \nu(x_t).$$

$$g_j(y(x)) = G_j(y(x)), \quad x \in Q_j, \quad 1 \leq j \leq m+1,$$

$$r_\nu \mu_\nu > 2^{3-\frac{1}{N}} L_\nu \sqrt{N+3}, \quad 1 \leq \nu \leq m+1, \quad (18)$$

$$\varphi(\bar{y}) = \inf \{ \varphi(y^q) : g_j(y^q) \leq 0, 1 \leq j \leq m, q = 1, 2, \dots \} \leq \varphi(y_\epsilon).$$

$$\epsilon_\nu = \mu_\nu \delta, \quad 1 \leq \nu \leq m, \quad (19)$$

#### 4. Index Algorithm with Dual Lipschitz Constant Estimates

$$\frac{r_\nu \mu_\nu}{2^{3-\frac{1}{N}} L_\nu \sqrt{N+3}}, \quad 1 \leq \nu \leq m+1, \quad (20)$$

$$R(i) = R_{glob}(i) = R_{loc}(i) = 2\Delta_i > 0. \quad (21)$$

$$z_i + z_{i-1} - 2z_M^* = |z_i - z_{i-1}| \leq \mu_M \Delta_i$$

$$\begin{aligned} R(i) &= \Delta_i + \frac{(z_i - z_{i-1})^2}{r_M^2 \mu_M^2 \Delta_i} - 2 \frac{z_i + z_{i-1} - 2z_M^*}{r_M \mu_M} \geq \Delta_i + \frac{(z_i - z_{i-1})^2}{r_M^2 \mu_M^2 \Delta_i} - 2 \frac{\mu_M \Delta_i}{r_M \mu_M} \\ &\geq \Delta_i + \frac{(z_i - z_{i-1})^2}{r_M^2 \mu_M^2 \Delta_i} - 2 \frac{\Delta_i}{r_M} = \Delta_i + \Delta_i \frac{(z_i - z_{i-1})^2 / \Delta_i^2}{r_M^2 \mu_M^2 \Delta_i^2} - 2 \frac{\Delta_i}{r_M} \\ &\geq \Delta_i + \frac{\Delta_i}{r_M^2} - 2 \frac{\Delta_i}{r_M} = \Delta_i \left( 1 - \frac{1}{r_M} \right)^2 > 0 \end{aligned}$$

$$R_{glob}(i) = \Delta_i \left(1 - \frac{1}{r_M^{glob}}\right)^2 > \Delta_i \left(1 - \frac{1}{r_M^{loc}}\right)^2 = R_{loc}(i) \quad (22)$$

$$R_{glob}(i) = \rho R_{loc}(i) \quad (23)$$

$$R(i) = \max \{R_{glob}(i), \rho R_{loc}(i)\} \quad (24)$$

$$R(t) = \max \{R(i) : 1 \leq i \leq k+1\} \quad (25)$$

$$x^{k+1} = \frac{x_t + x_{t-1}}{2}, \quad \nu(x_{t-1}) \neq \nu(x_t).$$

$$x^{k+1} = \frac{x_t + x_{t-1}}{2} + \frac{\text{sign}(z_t - z_{t-1})}{2r_\nu} \left[ \frac{|z_t - z_{t-1}|}{\mu_\nu} \right]^N, \quad \nu = \nu(x_{t-1}) = \nu(x_t).$$

$$\begin{aligned} \varphi(y_1, y_2) = & -1.5y_1^2 \exp \{1 - y_1^2 - 20.25(y_1 - y_2)^2\} - \\ & - (0.5(y_1 - 1)(y_2 - 1))^4 \exp \left\{ 2 - (0.5(y_1 - 1))^4 - (y_2 - 1)^4 \right\} \end{aligned}$$

$$\begin{aligned} g_1(y_1, y_2) &= 0.01 \left( (y_1 - 2.2)^2 + (y_2 - 1.2)^2 - 2.25 \right) \leq 0, \\ g_2(y_1, y_2) &= 100 \left( 1 - (y_1 - 2)^2/1.44 + (0.5y_2)^2 \right) \leq 0, \\ g_3(y_1, y_2) &= 10 \left( y_2 - 1.5 - 1.5 \sin(6.283(y_1 - 1.75)) \right) \leq 0 \end{aligned}$$

$$\varphi(\bar{y}) = \inf \{ \varphi(y^k) : g_j(y^k) \leq 0, 1 \leq j \leq m, k = 1, 2, \dots \} \leq \varphi(y_\epsilon). \quad (26)$$

$$R_{glob}(t) < \rho R_{loc}(t)$$

$$z_j = g_\nu(y(x_j)) \leq g_\nu(y(\bar{x})) + 2L_\nu \sqrt{N+3}(x_j - \bar{x})^{1/N}, \quad \nu = \nu(x_j)$$

$$z_{j-1} = g_\nu(y(x_{j-1})) \leq g_\nu(y(\bar{x})) + 2L_\nu\sqrt{N+3}(\bar{x} - x_{j-1})^{1/N}, \quad \nu = \nu(x_{j-1})$$

$$g_\nu(y(\bar{x})) \leq -\epsilon_\nu, \quad 1 \leq \nu \leq m.$$

$$g_{m+1}(y(\bar{x})) \leq z_{m+1}^*$$

$$\begin{aligned} R(j) &= \Delta_i + \frac{(z_j - z_{j-1})^2}{r_\nu^2 \mu_\nu^2 \Delta_i} - 2 \frac{z_j + z_{j-1} - 2z_\nu^*}{r_\nu \mu_\nu} \\ &\geq \Delta_i + 4 \frac{z_\nu^* - \left( g_\nu(y(\bar{x})) + L_\nu\sqrt{N+3} \left( (x_j - \bar{x})^{\frac{1}{N}} + (\bar{x} - x_{j-1})^{\frac{1}{N}} \right) \right)}{r_\nu \mu_\nu} \\ &= \Delta_i - 4 \frac{L_\nu\sqrt{N+3} \left( (x_j - \bar{x})^{\frac{1}{N}} + (\bar{x} - x_{j-1})^{\frac{1}{N}} \right)}{r_\nu \mu_\nu} + 4 \frac{z_\nu^* - g_\nu(y(\bar{x}))}{r_\nu \mu_\nu} \\ &= \Delta_i - 4 \frac{L_\nu\sqrt{N+3} \left( \alpha^{\frac{1}{N}} + (1-\alpha)^{\frac{1}{N}} \right)}{r_\nu \mu_\nu} + 4 \frac{z_\nu^* - g_\nu(y(\bar{x}))}{r_\nu \mu_\nu} \\ &\geq \Delta_i \left( 1 - 4 \frac{L_\nu\sqrt{N+3} \max_{0 \leq \alpha \leq 1} \left( \alpha^{\frac{1}{N}} + (1-\alpha)^{\frac{1}{N}} \right)}{r_\nu \mu_\nu} \right) + 4 \frac{z_\nu^* - g_\nu(y(\bar{x}))}{r_\nu \mu_\nu} \\ &= \Delta_i \left( 1 - 4 \frac{2^{3-1/N} L_\nu\sqrt{N+3}}{r_\nu \mu_\nu} \right) + 4 \frac{z_\nu^* - g_\nu(y(\bar{x}))}{r_\nu \mu_\nu} \geq 0 \end{aligned}$$

$$\begin{aligned} R(j) &= 2\Delta_i - 4 \frac{z_j - z_\nu^*}{r_\nu \mu_\nu} \geq 2\Delta_i + 4 \frac{z_\nu^* - \left( g_\nu(y(\bar{x})) + 2L_\nu\sqrt{N+3}(x_j - \bar{x})^{\frac{1}{N}} \right)}{r_\nu \mu_\nu} \\ &= 2\Delta_i - 8 \frac{L_\nu\sqrt{N+3}(x_j - \bar{x})^{\frac{1}{N}}}{r_\nu \mu_\nu} + 4 \frac{z_\nu^* - g_\nu(y(\bar{x}))}{r_\nu \mu_\nu} \\ &\geq 2\Delta_i \left( 1 - 4 \frac{L_\nu\sqrt{N+3}}{r_\nu \mu_\nu} \right) + 4 \frac{z_\nu^* - g_\nu(y(\bar{x}))}{r_\nu \mu_\nu} > 0 \end{aligned}$$

$$R_{glob}(j) > \rho R_{loc}(t) \tag{27}$$

## 5. TEST

There are various bibliography styles available. You can select the style of  
10 your choice in the preamble of this document. These styles are Elsevier styles  
based on standard styles like Harvard and Vancouver. Please use BibTeX to  
generate your bibliography and include DOIs whenever available.

Here are two sample references:

[1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23,  
15 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36].

## References

- [1] Y. Evtushenko, Numerical methods for finding global extreme (case of a  
non-uniform mesh), *Comput. Maths. Math. Phys.* 11 (6) (1971) 38–54.
- [2] S. Piyavskii, An algorithm for finding the absolute extremum of a function,  
20 *Comput. Maths. Math. Phys.* 12 (4) (1972) 57–67.
- [3] B. Shubert, A sequential method seeking the global maximum of a function,  
*SIAM Journal on numerical analysis* 9 (3) (1972) 379–388.
- [4] R. Strongin, On a certain global-minimization algorithm, *Radiophysics and  
Quantum Electronics* 13 (4) (1970) 419–424.
- 25 [5] Y. Evtushenko, V. Malkova, A. A. Stanevichyus, Parallel global optimiza-  
tion of functions of several variables, *Comput. Math. Math. Phys.* 49 (2)  
(2009) 246–260.
- [6] Y. Evtushenko, M. Posypkin, A deterministic approach to global box-  
constrained optimization, *Optim. Lett.* 7 (2013) 819–829.
- 30 [7] Strongin R.G., Sergeyev Ya.D., Global optimization with non-convex con-  
straints. Sequential and parallel algorithms, Kluwer Academic Publishers,  
Dordrecht, 2000.

- [8] Sergeyev, Y.D., Strongin, R.G., Lera, D., Introduction to global optimization exploiting space-filling curves, Springer Briefs in Optimization, Springer, New York, 2013.
- [9] J. Pinter, Global optimization in action (Continuous and Lipschitz Optimization: Algorithms, Implementations and Applications), Kluwer Academic Publishers, Dordrecht, 1996.
- [10] Jones, D.R., The DIRECT global optimization algorithm, in: The Encyclopedia of Optimization, Springer, Heidelberg, 2009, pp. 725–735.
- [11] G. Wood, Multidimensional bisection applied to global optimisation, Comput. Math. Appl. 21 (6-7) (1991) 161–172.
- [12] C. Meewella, D. Mayne, An algorithm for global optimization of lipschitz continuous functions, J. Optim. Theory Appl. 57 (2) (1988) 307–322.
- [13] R. Mladineo, An algorithm for finding the global maximum of a multimodal multivariate function, Math. Program. 34 (2) (1986) 188–200.
- [14] Vaz, A. I.F. and Vicente, L.N., PSwarm: a hybrid solver for linearly constrained global derivative-free optimization, Optimization Methods and Software 24 (4-5) (2009) 669–685.
- [15] R. Stripinis, R. Paulavičius, J. Žilinskas, Penalty functions and two-step selection procedure based DIRECT-type algorithm for constrained global optimization, J. Struct Multidisc Optim 59 (6) (2019) 2155.
- [16] R. Paulavičius, J. Žilinskas, Advantages of simplicial partitioning for Lipschitz optimization problems with linear constraints, Optim. Lett. 10 (2) (2016) 237–246.
- [17] G. Di Pillo, S. Lucidi, F. Rinaldi, An approach to constrained global optimization based on exact penalty functions, J Glob Optim 54 (2012) 251.
- [18] G. Di Pillo, G. Liuzzi, S. Lucidi, A DIRECT-type approach for derivative-free constrained global optimization, Comput Optim Appl 65 (2016) 361.



- 60 [19] K. Barkalov, I. Lebedev, Comparing two approaches for solving constrained global optimization problems, *Lecture Notes in Computer Science* 10556 (2017) 301–306.
- [20] K. Barkalov, I. Lebedev, Parallel algorithm for solving constrained global optimization problems, *Lecture Notes in Computer Science* 10421 (2017) 396–404.
- 65 [21] Sergeyev, Y.D., Kvasov, D.E., Global search based on efficient diagonal partitions and a set of Lipschitz constants, *SIAM J. Optim.* 16 (3) (2006) 910–937.
- [22] J. Žilinskas, Branch and bound with simplicial partitions for global optimization, *Math. Model. Anal.* 13 (1) (2008) 145–159.
- 70 [23] V. Sovrasov, Comparison of several stochastic and deterministic derivative-free global optimization algorithms, *Lecture Notes in Computer Science* 11548 (2019) 70–81.
- [24] Kvasov, D.E., Pizzuti, C., Sergeyev, Y.D., Local tuning and partition strategies for diagonal GO methods, *Numer. Math.* 94 (1) (2003) 93–106.
- 75 [25] Lera, D., Sergeyev, Ya.D., An information global minimization algorithm using the local improvement technique, *J. Glob. Optim.* 48 (1) (2010) 99–112.
- [26] Sergeyev, Y.D., Mukhametzhanov, M.S., Kvasov, D.E., Lera, D., Derivative-free local tuning and local improvement techniques embedded in the univariate global optimization, *J. Optim. Theory Appl.* 171 (1) (2016) 186–208.
- 80 [27] R. Horst, H. Tuy, *Global Optimization – Deterministic Approaches*, Springer-Verlag, Berlin, 1996.
- [28] Gablonsky, J.M., Kelley, C.T., A locally-biased form of the DIRECT algorithm, *J. Glob. Optim.* 21 (1) (2001) 27–37.
- 85

- [29] D. Jones, C. Perttunen, B. Stuckman, Lipschitzian optimization without the Lipschitz constant, *J. Optim. Theory Appl.* 79 (1) (1993) 157–181.
- [30] Gaviano, M., Kvasov, D.E., Lera, D., Sergeyev, Ya.D., Software for generation of classes of test functions with known local and global minima for global optimization, *ACM Transactions on Mathematical Software* 29 (4) (2003) 469–480.
- [31] K. Barkalov, R. Strongin, Solving a set of global optimization problems by the parallel technique with uniform convergence, *J. Glob. Optim.* 71 (1) (2018) 21–36.
- [32] Paulavičius, R., Sergeyev, Y.D., Kvasov, D.E., Žilinskas, J., Globally-biased DISIMPL algorithm for expensive global optimization, *J. Glob. Optim.* 59 (2-3) (2014) 545–567.
- [33] Sergeyev, Y.D., Kvasov, D.E., A deterministic global optimization using smooth diagonal auxiliary functions, *Commun. Nonlinear. Sci. Numer. Simulat.* 21 (1-3) (2015) 99–111.
- [34] R. Strongin, V. Gergel, K. Barkalov, A. Sysoyev, Generalized parallel computational schemes for time-consuming global optimization, *Lobachevskii Journal of Mathematics* 39 (4) (2018) 576–586.
- [35] V. Gergel, An approach for generating test problems of constrained global optimization, *Lecture Notes in Computer Science* 10556 (2017) 314–319.
- [36] V. Gergel, K. Barkalov, I. Lebedev, M. Rachinskaya, A. Sysoyev, A flexible generator of constrained global optimization test problems, *AIP Conference Proceedings* 2070 (2019) 20009.