Coláiste na Tríonóide, Baile Átha Cliath
Trinity College Dublin
Ollscoil Átha Cliath | The University of Dublin

School of Social Sciences and Philosophy

Assignment Submission Form

Student Name:	Alexander	Student ID Number:	24364960
Programme Title:	PhD Political Science	Module Title:	Quantiative Methods I
Date Submitted:	22.10.2026	Lecturer(s):	Dr. Jeffrey Ziegler
Assessment Title:	Homework 2		

PLAGIARISM DECLARATION

☑ I have read and understand the plagiarism provisions in the General Regulations (http://www.tcd.ie/calendar)

☑ I have completed the Online Tutorial 'Ready, Steady, Write' (https://libguides.tcd.ie/academic-integrity/ready-steady-write)

GENERATIVE AI USE DECLARATION

☑ I have read Trinity's Statement on Generative AI and understand my responsibilities (https://www.tcd.ie/media/tcd/academic-affairs/pdfs/College_Statement_on_GenAI_v2_2026.pdf)

DISCLOSURE OF GenAI USE

Choose all options that apply to your use of Generative AI tools (e.g., ChatGPT, Claude, Gemini, Copilot, Grammarly AI, etc.) in preparing this work:

☐ **NO GenAI USED**
I did not use any generative AI tools in the preparation of this work.

☐ **BRAINSTORMING**
I used GenAI to help explore topics or brainstorm approaches.

☐ **RESEARCH/LEARNING ASSISTANCE**
I used GenAI to help locate sources, understand concepts, or clarify information.

☐ **WRITING ASSISTANCE**
I used GenAI to help draft sections of my work, but I substantially revised, fact-checked, and edited all content.

☐ **DIRECT AI CONTENT INCLUSION**
I have included AI-generated content in my work with proper citation and quotation marks as required by Trinity guidelines. (https://libguides.tcd.ie/gen-ai/acknowledging-referencing)

☑ **OTHER USE**
(Please describe below any other way you used GenAI that doesn't fit the categories above.)


Checked the spelling with Grammarly.
Used Gemini to understand the different assumptions, that effects the p-value of a bivariate regession compared with a t-test (not really a task of the homework assignment). Furthermore, I used it to solve some LaTeX errors.

ACADEMIC INTEGRITY COMMITMENT

I understand that:

- This submission is my own work. I have not plagiarised, I have properly cited all sources, I have not used unauthorised assistance, and I comply with Trinity's academic integrity policies.
- Proper GenAI disclosure demonstrates academic integrity and AI literacy.
- Undisclosed use of GenAI constitutes plagiarism under Trinity policy.
- I am responsible for fact-checking all AI-generated information.
- **I have familiarised myself with my module's specific policy on the use of GenAI.**
- Transparency in AI use is valued and supports my academic development.

I declare that the information provided above is complete and accurate.

Signed:  Date: 22.10.2026

Problem Set 2

Applied Stats/Quant Methods 1

Due: October 22, 2026

Instructions

- Please show your work! You may lose points by simply writing in the answer. If the problem requires you to execute commands in `R`, please include the code you used to get your answers. Please also include the `.R` file that contains your code. If you are not sure if work needs to be shown for a particular problem, please ask.
- Your homework should be submitted electronically on GitHub.
- This problem set is due before 23:59 on Thursday October 22, 2026. No late assignments will be accepted.

Question 1: Political Science

The following table was created using the data from a study run in a major Latin American city.[1] As part of the experimental treatment in the study, one employee of the research team was chosen to make illegal left turns across traffic to draw the attention of the police officers on shift. Two employee drivers were upper class, two were lower class drivers, and the identity of the driver was randomly assigned per encounter. The researchers were interested in whether officers were more or less likely to solicit a bribe from drivers depending on their class (officers use phrases like, "We can solve this the easy way" to draw a bribe). The table below shows the resulting data.

[1] Fried, Lagunes, and Venkataramani (2010). "Corruption and Inequality at the Crossroad: A Multimethod Study of Bribery and Discrimination in Latin America. *Latin American Research Review.* 45 (1): 76-97.

	Not Stopped	Bribe requested	Stopped/given warning
Upper class	14	6	7
Lower class	7	7	1

(a) Calculate the χ^2 test statistic by hand/manually (even better if you can do "by hand" in R).

As we have two categorical variables, we use the $chi - test$ for **statistical independence**. The test statistic is calculated with the following formula:

$$\chi^2 = \sum_{i=1}^{k} \frac{(f_o - f_e)^2}{f_e}$$

Where $f_{e,i}$ is calculated with the formula:

$$f_e = \frac{\text{Row Total} \cdot \text{Column Total}}{\text{Total Observations}} \tag{1}$$

I did it partly manually in R, using matrices and the way R handles mathematical operations with this data structure. First, I created a matrix containing the values of our contingency table.

```
experiment <- matrix(c(14, 6, 7, 7, 7, 1), nrow = 2, byrow = TRUE)
```

By combining indexing with R's sum() function, we can now calculate the expected value for each cell. Since there are only 6 cells, I write out the formula explicitly.

```
> expected_frequency <- matrix(c(
+   sum(experiment[1, ]) * sum(experiment[, 1])/sum(experiment),
+   sum(experiment[1, ]) * sum(experiment[, 2])/sum(experiment),
+   sum(experiment[1, ]) * sum(experiment[, 3])/sum(experiment),
+   sum(experiment[2, ]) * sum(experiment[, 1])/sum(experiment),
+   sum(experiment[2, ]) * sum(experiment[, 2])/sum(experiment),
+   sum(experiment[2, ]) * sum(experiment[, 3])/sum(experiment)
+ ), nrow = 2, byrow = TRUE)
> expected_frequency
[,1]     [,2]     [,3]
[1,] 13.5 8.357143 5.142857
[2,]  7.5 4.642857 2.857143
```

Next, I calculated the χ^2 for each cell and summed them. I did this for two reasons: first, I discussed the problem set results with fellow students, which let us compare them more easily. Second, I thought this might help calculate the standardized residuals. However, I ended up using the full formula for the standardized residuals because I wanted to make sure that my probability calculations for rows and columns worked correctly.

```
     [,1]     [,2]     [,3]
[1,] 13.5 8.357143 5.142857
[2,]  7.5 4.642857 2.857143
> chi_squared <- (experiment - expected_frequency)**2 / expected_frequency
> chi_squared
[,1]      [,2]      [,3]
[1,] 0.01851852 0.6648352 0.6706349
[2,] 0.03333333 1.1967033 1.2071429
> sum(chi_squared) # Chi-squared has a value of 3.791168
[1] 3.791168
> chi_sum <- sum(chi_squared)
> chi_sum
[1] 3.791168
```

(b) Now calculate the p-value from the test statistic you just created (in `R`).[2] What do you conclude if $\alpha = 0.1$?

Using the `chi_sum` object from the previous task, I used the approriate R function:

```
pchisq(chi_sum,
       df = (nrow(experiment)-1)*(ncol(experiment)-1),
       lower.tail=FALSE)
```

The probability of a χ^2 value that is as large as our empirical χ^2 or larger is 0.1502. Therefore, we cannot reject H0 at an alpha level of 0.1, as $0.1502 > 0.10$. Based on our test the variables are **statistically independent**.

[2]Remember frequency should be > 5 for all cells, but let's calculate the p-value here anyway.

(c) Calculate the standardized residuals for each cell and put them in the table below.

I again used my matrix object to calculate the row and column probabilities. Afterward, I used cbind and rbind to create matrices that contain the values in a format to calculate with, without running into problems based on R's recycling of shorter vectors.

```
> p_row = rowSums(experiment) / sum(experiment)
> p_column = colSums(experiment) / sum(experiment)
>
> p_row_matrix <- cbind(p_row, p_row, p_row)
> p_column_matrix <- rbind(p_column, p_column)
> p_row_matrix
p_row     p_row     p_row
[1,] 0.6428571 0.6428571 0.6428571
[2,] 0.3571429 0.3571429 0.3571429
> p_column_matrix
[,1]      [,2]      [,3]
p_column  0.5 0.3095238 0.1904762
p_column  0.5 0.3095238 0.1904762
```

I plugged this matrices in the formula and received:

```
> pearson_res <- (experiment - expected_frequency) / sqrt(expected_frequency * (1
> pearson_res
p_row     p_row     p_row
[1,]  0.3220306 -1.641957  1.523026
[2,] -0.3220306  1.641957 -1.523026
```

The results could be used to fill the table.

	Not Stopped	Bribe requested	Stopped/given warning
Upper class	0.3220306	-1.641957	1.523026
Lower class	-0.3220306	1.641957	-1.523026

(d) How might the standardized residuals help you interpret the results?

Generally, the *not-stopped* category is very close to the expected value, while the *bribe accepted"* and *warned* categories show somewhat larger adjusted residuals.
This may indicate that our experimental treatment did not affect whether someone was stopped in the first place, but rather whether the police officer accepted a bribe after they stopped someone. One confounding reason could be that police officers decided to stop cars based on factors outside our experimental treatment (e.g., a desire to make a break). Therefore, it may be reasonable to recalculate the test statistic without the not-stopped category.

Question 2: Economics

Chattopadhyay and Duflo were interested in whether women promote different policies than men.[3] Answering this question with observational data is pretty difficult due to potential confounding problems (e.g. the districts that choose female politicians are likely to systematically differ in other aspects too). Hence, they exploit a randomized policy experiment in India, where since the mid-1990s, $\frac{1}{3}$ of village council heads have been randomly reserved for women. A subset of the data from West Bengal can be found at the following link: `https://raw.githubusercontent.com/kosukeimai/qss/master/PREDICTION/women.csv`

```
df_india <- readr::read_csv("https://raw.githubusercontent.com/kosukeimai/qss/
    master/PREDICTION/women.csv")
```

Each observation in the data set represents a village and there are two villages associated with one GP (i.e. a level of government is called "GP"). Figure 1 below shows the names and descriptions of the variables in the dataset. The authors hypothesize that female politicians are more likely to support policies female voters want. Researchers found that more women complain about the quality of drinking water than men. You need to estimate the effect of the reservation policy on the number of new or repaired drinking water facilities in the villages.

Figure 1: Names and description of variables from Chattopadhyay and Duflo (2004).

Name	Description
`GP`	An identifier for the Gram Panchayat (GP)
`village`	identifier for each village
`reserved`	binary variable indicating whether the GP was reserved for women leaders or not
`female`	binary variable indicating whether the GP had a female leader or not
`irrigation`	variable measuring the number of new or repaired irrigation facilities in the village since the reserve policy started
`water`	variable measuring the number of new or repaired drinking-water facilities in the village since the reserve policy started

[3] Chattopadhyay and Duflo. (2004). "Women as Policy Makers: Evidence from a Randomized Policy Experiment in India. *Econometrica*. 72 (5), 1409-1443.

(a) State a null and alternative (two-tailed) hypothesis.

H_0: The number of water facilities is not different between villages with reserved seats at the GP level and villages without reserved seats at the GP level.

H_1: The number of water facilities is different between villages with reserved seats at the GP level and villages without reserved seats at the GP level.

(b) Run a bivariate regression to test this hypothesis in `R` (include your code!).

```
> model_india <- lm(data = df_india,
+                   water ~ as.factor(reserved))
> summary(model_india)

Call:
lm(formula = water ~ as.factor(reserved), data = df_india)

Residuals:
Min      1Q  Median      3Q     Max
-23.991 -14.738  -7.865   2.262 316.009

Coefficients:
Estimate Std. Error t value Pr(>|t|)
(Intercept)            14.738      2.286   6.446 4.22e-10 ***
as.factor(reserved)1    9.252      3.948   2.344   0.0197 *
---
Signif. codes:  0 ‘***’ 0.001 ‘**’ 0.01 ‘*’ 0.05 ‘.’ 0.1 ‘ ’ 1

Residual standard error: 33.45 on 320 degrees of freedom
Multiple R-squared:  0.01688, Adjusted R-squared:  0.0138
```

```
F-statistic: 5.493 on 1 and 320 DF,  p-value: 0.0197
```

The p-value of $0.0197 < 0.05$. Therefore, we can reject the $H0$. In other words, there is a difference in repaired or newly installed water facilities between villages with and without a reservation policy at the gp level. However, we should be aware that the observations are not really independent, as villages from the same GP will very likely deviate from our estimate in the same way.

(c) Interpret the coefficient estimate for reservation policy.

Without a categorical (binary) covariate, the intercept estimates the value for the reference category ($reservationpolicy = 0$), while the slope captures the difference between villages with and without a reservation policy. Specifically, villages in GPs without a reservation policy saw 14.73 facilities repaired or newly installed, while villages with a reservation policy saw an additional 9.25 facilities repaired or newly installed, for a total of about 24 repaired or newly installed facilities.