



**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

**INSPIRING CREATIVE AND INNOVATIVE MINDS**

[www.utm.my](http://www.utm.my)

# Chapter 4

# Graph Theory


## (Part 1)

- A graph  $G$  is a triple  $(V, E, f)$ , where
  - $V$  is a finite nonempty set, called the set of vertices
  - $E$  is a finite set (may be empty), called the set of edges
  - $f$  is a function, called an incidence function, that assign to each edge,  $e \in E$ , a one-element subset  $\{v\}$  or a two-element subset  $\{v, w\}$ , where  $v$  and  $w$  are vertices.
- We can write  $G$  as  $(V, E, f)$  or  $(V, E)$  or simply as  $G$ .

[www.utm.my](http://www.utm.my)

- Let,
  - $V = \{v_1, v_2, v_3, v_4, v_5, v_6, v_7\}$
  - $E = \{e_1, e_2, e_3, e_4, e_5, e_6, e_7\}$
- and  $f$  be defined by
  - $f(e_1) = f(e_2) = \{v_1, v_2\}$
  - $f(e_3) = \{v_4, v_3\}$
  - $f(e_4) = f(e_6) = f(e_7) = \{v_6, v_3\}$
  - $f(e_5) = \{v_2, v_4\}$
- Then  $G = (V, E, f)$  is a graph

## example



prepared by Razana Alwee


## example

- Let  $V = \{v_1, v_2, v_3, v_4\}$ ,  $E = \{e_1, e_2, e_3\}$   
and
  - $f(e_1) = \{v_1, v_2\}$
  - $f(e_2) = \{v_3, v_3\}$
  - $f(e_3) = \{v_3, v_4\}$
- Then  $G = (V, E, f)$  is a graph





## example



prepared by Razana Alwee



**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

[www.utm.my](http://www.utm.my)

# Characteristics of Graph




# Adjacent Vertices

- An edge  $e$  in a graph that is associated with the pair of vertices  $v$  and  $w$  is said to be **incident** on  $v$  and  $w$ , and  $v$  and  $w$  are said to be incident on  $e$  and to be **adjacent vertices**.
- A vertex that is an endpoint of a loop is said to be adjacent to itself.



# example

$v_4$  and  $v_3$  are adjacent



prepared by Razana Alwee




# Isolated Vertex

- Let  $G$  be a graph and  $v$  be a vertex in  $G$ .
- We say that  $v$  is an isolated vertex if it is not incident with any edge.

## example

- $v_5$  and  $v_7$  are isolated vertices.




prepared by Razana Alwee

# Loop

- An edge incident on a single vertex is called a loop.

**Example:**  $e_2$  is a loop




prepared by Razana Alwee

## Parallel Edges

Two or more distinct edges with the same set of endpoints are said to be parallel.


- $e_1$  and  $e_2$  are **parallel**.

Example



## Example

Given a graph as shown below,



- Write a vertex set and the edge set, and give a table showing the edge-endpoint function.
- Find all edges that are incident on a, all vertices that are adjacent to a, all edges that are adjacent to  $e_2$ , all loops, all parallel edges, all vertices that are adjacent to themselves and all isolated vertices.



# Example 1 - Solution

**Solution:**

- a) Vertex set,  $V = \{a, b, c, d, e, f, i, h\}$  and  
the set of edges,  $E = \{e_1, e_2, e_3, e_4, e_5, e_6, e_7, e_8, e_9, e_{10}\}$

Edge	Endpoints
$e_1$	$\{a, b\}$
$e_2$	$\{a, c\}$
$e_3$	$\{b, e\}$
$e_4$	$\{c, d\}$
$e_5$	$\{d, e\}$
$e_6$	$\{a, b\}$
$e_7$	$\{d\}$
$e_8$	$\{i, h\}$
$e_9$	$\{h\}$
$e_{10}$	$\{i\}$



b)

incident on a,	e1, e2, e6
adjacent to a,	c, b
adjacent to $e_2$ ,	e1, e4, e6
loops,	e7, e9, e10
parallel edges,	e1, e6
adjacent to themselves,	i, h, d
isolated vertices,	f



**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

[www.utm.my](http://www.utm.my)

# The Concept of Degree




## Degree of a vertex

- Let  $G$  be a graph and  $v$  be a vertex of  $G$ .
- The degree of  $v$ , written  $\deg(v)$  or  $d(v)$  is the number of edges incident with  $v$ .
- Each loop on a vertex  $v$  contributes **2** to the degree of  $v$ .

## example


- $\deg(v_1) = 1$ ;  $\deg(v_2) = 2$ ;  $\deg(v_3) = 3$ ;  $\deg(v_4) = 2$



prepared by Razana Alwee

## exercise

- Find the degree of each vertex in the graph.




prepared by Razana Alwee



## solution

- Find the degree of each vertex in the graph.



**Solution:**  $\deg(v_1) = 1$ ;  $\deg(v_2) = 4$ ;  $\deg(v_3) = 4$ ;  $\deg(v_4) = 5$

prepared by Razana Alwee




**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

[www.utm.my](http://www.utm.my)

# Types of Graphs

## Simple Graphs


- A graph  $G$  is called a **simple graph** if  $G$  does not contain any parallel edges and any loops.
- **Example**



prepared by Razana Alwee

## Connected Graph


- A graph  $G$  is connected if given any vertices  $v$  and  $w$  in  $G$ , there is a path from  $v$  to  $w$ .
- **Example:**



prepared by Razana Alwee

## example

- not connected



prepared by Razana Alwee





# Regular Graphs

- Let  $G$  be a graph and  $k$  be a nonnegative integer.
- $G$  is called a  $k$ -regular graph if the degree of each vertex of  $G$  is  $k$ .






# example



prepared by Razana Alwee

## Complete Graph

- A simple graph with  $n$  vertices in which there is an edge between every pair of distinct vertices is called a complete graph on  $n$  vertices.
- This is denoted by  $K_n$ .
- **Example**






prepared by Razana Alwee



# Subgraph

- Let  $G=(V,E)$  be a graph.
- $H=(U,D)$  is a subgraph of  $G$  if
  - $U \subseteq V$  and  $D \subseteq E$
  - for every edge  $e \in D$ , if  $e$  is incident on  $v$  and  $w$ , then  $v, w \in V$ .

# example





**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

[www.utm.my](http://www.utm.my)

# Graph Representation



# Matrix Representation of a Graph


- To write programs that process and manipulate graphs, the graphs must be stored, that is, represented in computer memory.
- A graph can be represented (in computer memory) in several ways.
- 2-dimensional array: adjacency matrix and incidence matrix.



# Adjacency Matrices

- Let  $G$  be a graph with  $n$  vertices.
- The adjacency matrix,  $A_G$  is an  $n \times n$  matrix  $[a_{ij}]$  such that,  
 $a_{ij}$  = the number of edges from  $v_i$  to  $v_j$ , {undirected  $G$ }  
or,  
 $a_{ij}$  = the number of arrows from  $v_i$  to  $v_j$ , {directed  $G$ }  
for all  $i, j = 1, 2, \dots, n$ .


## example



$$A_G = \begin{bmatrix} 0 & 1 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 1 & 0 \end{bmatrix}$$

prepared by Razana Alwee

## example



$$A_G = \begin{bmatrix} 0 & 2 & 1 & 0 \\ 2 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$


prepared by Razana Alwee



## example

$$A_G = \begin{bmatrix} 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 2 & 0 & 1 \\ 1 & 2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 \end{bmatrix}$$

## example



prepared by Razana Alwee


# Adjacency Matrices

- Notice that the matrix  $A_G$  is a **symmetric matrix** if it is representing an undirected graph, where

$$a_{ij} = a_{ji}$$

- If  $G$  is a directed graph (**digraph**), then  $A_G$  need not be a symmetric matrix.

## example



$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 0 & 0 \end{bmatrix}$$

prepared by Razana Alwee


# Incidence Matrices

- Let  $G$  be a graph with  $n$  vertices and  $m$  edges.
- The incidence matrix  $I_G$  is an  $n \times m$  matrix  $[a_{ij}]$  such that,

$$a_{ij} = \begin{cases} 0 & \text{if } v_i \text{ is not an end vertex of } e_j, \\ 1 & \text{if } v_i \text{ is an end vertex of } e_j, \text{ but } e_j \text{ is not a loop} \\ 2 & \text{if } e_j \text{ is a loop at } v_i \end{cases}$$



## example




$$\begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 0 & 1 \\ 1 & 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \end{matrix}$$

Notice that the sum of the  $i$ th row is the degree of  $v_i$


## exercise

- Find the adjacency matrix and the incidence matrix of the graph.



prepared by Razana Alwee

# Exercise - Solution



Adjacency matrix

Incidence matrix

$$A_G = \begin{matrix} & \begin{matrix} v_1 & v_2 & v_3 & v_4 \end{matrix} \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{bmatrix} \end{matrix}$$

$$I_G = \begin{matrix} & \begin{matrix} e_1 & e_2 & e_3 & e_4 \end{matrix} \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} \end{matrix}$$



## Exercise Past Year 2015/2016


A cat show is being judged from pictures of the cats. The judges would like to see pictures of the following pairs of cats next to each other for their final decision: Fifi and Putih, Fifi and Suri, Fifi and Bob, Bob and Cheta, Bob and Didi, Bob and Suri, Cheta and Didi, Didi and Suri, Didi and Putih, Suri and Putih, Putih and Jeep, Jeep and Didi.

Draw a graph modeling this situation.

(3 marks)


# Exercise Solution

## Past Year 2015/2016



# Exercise Past Year 2015/2016

Given a graph as shown in Figure 1.



**Figure 1**

- Find the incidence matrix of the graph. (4 marks)
- Find the adjacency matrix of the graph. (3 marks)

## Exercise Solution Past Year 2015/2016

(i) Incident Matrix,  $I_G =$

	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$i$	$k$
1	1	1	0	1	0	0	0	0	0	0
2	1	0	1	0	0	0	0	0	0	0
3	0	1	0	0	1	1	1	0	0	0
4	0	0	1	1	1	0	0	1	0	1
5	0	0	0	0	0	1	0	0	0	0
6	0	0	0	0	0	0	1	1	2	1

(ii) Adjacency Matrix,  $A_G =$

	1	2	3	4	5	6
1	0	1	1	1	0	0
2	1	0	0	1	0	0
3	1	0	0	1	1	1
4	1	1	1	0	0	2
5	0	0	1	0	0	0
6	0	0	1	2	0	1




**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

[www.utm.my](http://www.utm.my)


# Isomorphisms



# Isomorphism



$G_1$



$G_2$

- Are these 2 graphs the same?
  
- When we say that 2 graphs are the same mean they are isomorphic to each other.



# Isomorphism

- Graphs  $G_1$  and  $G_2$  are isomorphic if there is  
a one-to-one, onto function  $f$  from the vertices of  $G_1$  to  
the vertices of  $G_2$   
and  
a one-to-one, onto function  $g$  from the edges of  $G_1$  to  
the edges of  $G_2$

## Isomorphism

- An edge  $e$  is incident on  $v$  and  $w$  in  $G_1$  if and only if the edge  $g(e)$  is incident on  $f(v)$  and  $f(w)$  in  $G_2$ .
- The pair of functions  $f$  and  $g$  is called an isomorphism of  $G_1$  onto  $G_2$ .
- Graphs  $G_1$  and  $G_2$  are isomorphic if and only if for some ordering of their vertices, their adjacency matrices are equal.

## Definition



Let  $G = \{V, E\}$  and  $G' = \{V', E'\}$  be graphs.  $G$  and  $G'$  are said to be isomorphic if there exist a pair of functions  $f : V \rightarrow V'$  and  $g : E \rightarrow E'$  such that  $f$  associates each element in  $V$  with exactly one element in  $V'$  and vice versa;  $g$  associates each element in  $E$  with exactly one element in  $E'$  and vice versa, and for each  $v \in V$ , and each  $e \in E$ , if  $v$  is an endpoint of the edge  $e$ , then  $f(v)$  is an endpoint of the edge  $g(e)$ .



- ◆ If two graphs is isomorphic, they must have:
  - the same number of vertices and edges,
  - the same degrees for corresponding vertices,
  - the same number of connected components,
  - the same number of loops and parallel edges,
  - both graphs are connected or both graph are not connected,
  - pairs of connected vertices must have the corresponding pair of vertices connected.
  
- ◆ In general, it is easier to prove two graphs are not isomorphic by proving that one of the above properties fails.



## example

- Determine whether  $G$  is isomorphic to  $H$ .





## example

- Both graphs are simple and have the **same number of vertices** and the **same number of edges**.



## example

- All the vertices of both graphs have degree 2.







## example

- Define  $f: U \rightarrow V$ , where  $U = \{u_1, u_2, u_3, u_4\}$  and  $V = \{v_1, v_2, v_3, v_4\}$

$$f(u_1) = v_1, \quad f(u_2) = v_4, \quad f(u_3) = v_3, \quad f(u_4) = v_2$$



prepared by Razana Alwee



- To verify whether  $G$  and  $H$  are isomorphic, we examine the adjacency matrix  $A_G$  with rows and columns labeled in the order  $u_1, u_2, u_3, u_4$  and the adjacency matrix  $A_H$  with rows and columns labeled in the order  $v_1, v_4, v_3, v_2$ .

## example



- $A_G$  and  $A_H$  are the same,  $G$  and  $H$  are isomorphic.

$$A_G = \begin{matrix} & \begin{matrix} u_1 & u_2 & u_3 & u_4 \end{matrix} \\ \begin{matrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix} \end{matrix} \quad A_H = \begin{matrix} & \begin{matrix} v_1 & v_4 & v_3 & v_2 \end{matrix} \\ \begin{matrix} v_1 \\ v_4 \\ v_3 \\ v_2 \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix} \end{matrix}$$



prepared by Razana Alwee

## Exercise

**Q: Show that the following two graphs are isomorphic.**



# Exercise Solution




- Both have 5 vertices and 6 edges
- Both are connected and simple graph
- Both have 2 vertices with 3 degree and 3 vertices with 2 degree
- $f(A_{G_1}) = A_{G_2}$        $f(B_{G_1}) = B_{G_2}$
- $f(C_{G_1}) = C_{G_2}$        $f(D_{G_1}) = D_{G_2}$
- $f(E_{G_1}) = E_{G_2}$
- $\therefore G_1$  and  $G_2$  are isomorphic


## Exercise

**Q: Is these two graphs are isomorphic?**

$G :$




$H :$




# Exercise Solution

$G :$




$H :$




- Both have 8 vertices and 10 edges
  - Both are connected and simple graph
  - Both have 4 vertices with 3 degree and 4 vertices with 2 degree
  - However  $f: G \rightarrow H$  cannot be defined
- $\therefore G$  and  $H$  are not isomorphic

## Exercise Past Year 2015/2016

Determine whether the graphs in Figure 2 (*A* and *B*) are isomorphic. If the graphs are isomorphic, find their adjacency matrices; otherwise, give an invariant that the graphs do not share. (6 marks)




(A)



(B)



# Exercise Solution Past Year 2015/2016



Adjacency Matrix,  $A_A =$

$$\begin{matrix} & \begin{matrix} a & b & c & d & e & f \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \\ e \\ f \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$

Adjacency Matrix,  $A_B =$

$$\begin{matrix} & \begin{matrix} x & w & u & y & t & v \end{matrix} \\ \begin{matrix} x \\ w \\ u \\ y \\ t \\ v \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$

- Both have 6 vertices and 8 edges
- Both are connected and simple graph
- Both have 1 vertex with 4 degree, 2 vertices with 3 degree and 3 vertices with 2 degree
- $f(a_A) = x_B$                        $f(b_A) = w_B$   
 $f(c_A) = u_B$                        $f(d_A) = y_B$   
 $f(e_A) = t_B$                        $f(f_A) = v_B$
- ∴  $A$  and  $B$  are isomorphic



**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA

[www.utm.my](http://www.utm.my)

# Trails, Paths & Circuits

# Term and Description

- A **walk** from  $v$  to  $w$  is a finite alternating sequence of adjacent vertices and edges of  $G$ . Thus a walk has the form

$$(v_0, e_1, v_1, e_2, v_2, \dots, v_{n-1}, e_n, v_n)$$

where the  $v$ 's represent vertices, the  $e$ 's represent edges,  $v = v_0$ ,  $w = v_n$ , and for  $i = 1, 2, \dots, n$ .  $v_{i-1}$  and  $v_i$  are the endpoints of  $e_i$ .

- A **trivial walk** from  $v$  to  $w$  consist of the single vertex  $v$ . The walk contains zero edges (has length zero)
- The **length of a walk** is the number of edges it has.




## Term and Description (cont.)

- A **trail** from  $v$  to  $w$  is a walk from  $v$  to  $w$  that does not contain a repeated edge.
- A **path** from  $v$  to  $w$  is a trail from  $v$  to  $w$  that does not contain a repeated vertex.
- A **closed walk** is a walk that start and ends at the same vertex.
- A **circuit/cycle** is a closed walk that contains at least one edge and does not contain a repeated edge.
- A **simple circuit** is a circuit that does not have any other repeated vertex except the first and the last.

## example


- $(1, a, 2, b, 3, c, 4, d, 2)$  is a walk of length 4 from vertex 1 to vertex 2.



prepared by Razana Alwee

## example

- $(1, a, 2, b, 3, c, 4, d, 2, e, 5)$  is a trail.




**Note:**

Trail: No repeated edge (can repeat vertex).

prepared by Razana Alwee

## example

- (6, g, 5, e, 2, d, 4) is a path.




**Note:**

Path: No repeated vertex and edge.

## example

- $(2, f, 6, g, 5, e, 2, d, 4, c, 3, b, 2)$  is a circuit/cycle.




**Note:** circuit → start and end at same vertex, no repeated edge.

prepared by Razana Alwee



## example


- $(5, g, 6, f, 2, e, 5)$  is a simple circuit.



**Note:** Simple circuit → start and end at same vertex, no repeated edge or vertex except for the start and end vertex.

## exercise

- Tell whether the following is either a walk, trail, path, circuit, simple circuit, closed walk or none of these.
  - $(v_1, e_1, v_2)$
  - $(v_2, e_2, v_3, e_3, v_4, e_4, v_3)$
  - $(v_4, e_7, v_5, e_6, v_1, e_1, v_2, e_2, v_3, e_3, v_4)$
  - $(v_4, e_4, v_3, e_3, v_4, e_5, v_2, e_1, v_1, e_6, v_5, e_7, v_4)$



## Exercise solution

- Tell whether the following is either a walk, trail, path, circuit, simple circuit, closed walk or none of these.

■  $(v_1, e_1, v_2)$

**Path**

■  $(v_2, e_2, v_3, e_3, v_4, e_4, v_3)$


**Trail**

■  $(v_4, e_7, v_5, e_6, v_1, e_1, v_2, e_2, v_3, e_3, v_4)$

**Simple cycle**

■  $(v_4, e_4, v_3, e_3, v_4, e_5, v_2, e_1, v_1, e_6, v_5, e_7, v_4)$

**Cycle**





**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA


[www.utm.my](http://www.utm.my)

# Euler Trail & Circuit

## Euler Circuits

- A circuit in a graph that includes all the edges of the graph is called an Euler circuit.
- Let  $G$  be a graph. An **Euler circuit** for  $G$  is **a circuit that contains every vertex and every edges of  $G$** . That is, an Euler circuit for  $G$  is a sequence of adjacent vertices and edges in  $G$  that has at least one edges, starts and ends at the same vertex, **uses every vertex of  $G$  at least once**, and **uses every edge of  $G$  exactly once**.

# example




(1, a, 2, c, 5, e, 5, d, 4, f, 3, g, 2, b, 1)  
is an Euler circuit

prepared by Razana Alwee

## Euler Trail

- A trail from  $v$  to  $w$  ( $v \neq w$ ) with no repeated edges is called an Euler trail if it **contains all the edges and all the vertices**.
- Let  $G$  be a graph, and let  $v$  and  $w$  be two distinct vertices of  $G$ . An **Euler trail** from  $v$  to  $w$  is a sequence of adjacent vertices and edges that **starts at  $v$**  and **ends at  $w$** , **passes through every vertex of  $G$  at least once**, and **traverses every edge of  $G$  exactly once**.

## example



(1, a, 2, c, 5, e, 5, d, 4, f, 3, b, 2)  
is an Euler trail

prepared by Razana Alwee






## Theorem

- If  $G$  is a connected graph and **every vertex has even degree**, then  $G$  has an **Euler circuit**.
- A graph has an **Euler trail** from  $v$  to  $w$  ( $v \neq w$ ) if and only if it is connected and  **$v$  and  $w$  are the only vertices having odd degree**.


# Königsberg Bridge Problem



Starting at one land area, is it possible to walk across all of the bridges exactly once and return to the starting land area?

## Königsberg Bridge Problem

- Graph of the Königsberg Bridge Problem




no Euler circuit

It is not possible to walk across all of the bridges exactly once and return to the starting land area

prepared by Razana Alwee


## Königsberg Bridge Problem

- Since 1736, two additional bridges have been constructed on the Pregel river.



## example

Vertex	1	2	3	4	5	6	7
Degree	2	4	4	4	2	2	4




This graph has an Euler circuit

prepared by Razana Alwee

## example

Vertex	1	2	3	4	5	6
Degree	2	4	2	4	2	2




This graph has an Euler circuit

prepared by Razana Alwee

## example

Vertex	1	2	3	4	5	6
Degree	2	3	3	2	2	2




This graph has an Euler trail

prepared by Razana Alwee

## exercise

- Decide whether the graph has an Euler circuit. If the graph has an Euler circuit, exhibit one.




prepared by Razana Alwee



## Exercise solution



- Decide whether the graph has an Euler circuit. If the graph has an Euler circuit, exhibit one.





prepared by Razana Alwee

## exercise

Q: Which of the following graphs has Euler circuit?  
Justify your answer.



# Exercise Solution




**Q: Which of the following graphs has Euler circuit?  
Justify your answer.**

- $G_1$  has Euler circuit since all vertices have even degree
- $G_2$  does not has Euler circuit since 2 vertices (vertex  $a$  &  $b$ ) have odd degree


## Exercise Past Year 2015/2016

Determine whether the graph in Figure 3 has an Euler cycle or Euler path. If the graph has an Euler cycle or Euler path, exhibit one; otherwise, give an argument that shows there is no Euler path. (4 marks)



**Figure 3**

# Exercise Solution Past Year 2015/2016



- The graph does not has Euler circuit because 2 vertices (vertex 1 & 5) have odd degree
- Hence, the graph has Euler path
- **1-b-2-a-3-c-1-d-4-f-4-e-3-k-5-g-4-h-6-i-5**



**UTM**  
UNIVERSITI TEKNOLOGI MALAYSIA


[www.utm.my](http://www.utm.my)

# Hamilton Circuits

## Hamiltonian Circuit

- A circuit in a graph  $G$  is called a Hamiltonian circuit if it contains each vertex of  $G$ .
- Given a graph  $G$ , a **Hamiltonian circuit** for  $G$  is **a simple circuit that includes every vertex of  $G$**  (but doesn't need to include all edges). That is, a Hamiltonian circuit for  $G$  is a sequence of adjacent vertices and distinct edges in which **every vertex of  $G$  appears exactly once, except for the first and the last, which are the same.**

# Around the world game




Sir William Rowan Hamilton marketed a puzzle in the mid-1800s in the form of dodecahedron

prepared by Razana Alwee




# Around the world game



Each corner bore  
the name of a city

prepared by Razana Alwee


# Around the world game



The problem was to  
start at any city,  
travel along the  
edges,  
visit each city exactly  
one time  
and return to the  
initial city


prepared by Razana Alwee

# The graph



prepared by Razana Alwee


# Hamiltonian Circuit



**a-f-g-p-q-r-s-t-o-n-m-l-k-j-i-h-b-c-d-e-a**

## example


This graph has a Hamiltonian circuit



**1-a-2-b-3-f-5-e-4-k-6-l-7-m-1**

prepared by Razana Alwee


# example



**1-j-6-k-4-e-5-f-3-b-2-i-7-m-1**

prepared by Razana Alwee

## example




no Hamiltonian circuit

prepared by Razana Alwee

## example

This graph has a Hamiltonian circuit




**1-2-6-4-3-5-7-1**

prepared by Razana Alwee




## example



**1-6-4-2-3-5-7-1**

prepared by Razana Alwee

## example




no Hamiltonian circuit

prepared by Razana Alwee

## example


This graph has a Hamiltonian circuit



**1-2-3-4-5-1**

prepared by Razana Alwee

## example




**3-5-4-2-1-3**

prepared by Razana Alwee

## Hamiltonian Path


- A path in a graph  $G$  is called a Hamiltonian path if it contains each vertex of  $G$ .
- Example:



prepared by Razana Alwee

## exercise


- Find a Hamiltonian circuit in this graph.



prepared by Razana Alwee

## Exercise solution

- Find a Hamiltonian circuit in this graph.




**1-b-6-g-5-i-4-h-3-d-2-a-1**

prepared by Razana Alwee

# Exercise Past Year 2015/2016

Determine whether the graph in Figure 4 has an Hamiltonian cycle. If yes, exhibit one.


(3 marks)



**Figure 4**



# Exercise Solution Past Year 2015/2016



- The graph has Hamiltonian cycle
- ***7-f-3-c-4-d-1-a-2-e-6-k-5-m-8-h-7***