


Chapter 5: Trigonometry Reference

Right triangle trigonometry



A *right triangle* is defined as having one angle precisely equal to 90° (a *right angle*).

Trigonometric identities

[{#Trigonometric identities}]

$$\sin x = \frac{O}{H} \quad \cos x = \frac{A}{H} \quad \tan x = \frac{O}{A} \quad \tan x = \frac{\sin x}{\cos x}$$

$$\csc x = \frac{H}{O} \quad \sec x = \frac{H}{A} \quad \cot x = \frac{A}{O} \quad \cot x = \frac{\cos x}{\sin x}$$

H is the *Hypotenuse*, always being opposite the right angle. Relative to angle x , O is the *Opposite* and A is the *Adjacent*.

"Arc" functions such as "arcsin", "arccos", and "arctan" are the complements of normal trigonometric functions. These functions return an angle for a ratio input. For example, if the tangent of 45° is equal to 1, then the "arctangent" (arctan) of


1 is 45° . "Arc" functions are useful for finding angles in a right triangle if the side lengths are known.

The Pythagorean theorem

$\{\#\text{Pythagorean Theorem}\}$

$$H^2 = A^2 + O^2$$

Non-right triangle trigonometry



The Law of Sines (for *any* triangle)

$\{\#\text{Law of sines}\}$ $\{\#\text{Sines, law of}\}$

$$\frac{\sin a}{A} = \frac{\sin b}{B} = \frac{\sin c}{C}$$

The Law of Cosines (for *any* triangle)

[] {#Law of cosines} [] {#Cosines, law of}

$$A^2 = B^2 + C^2 - (2BC)(\cos a)$$

$$B^2 = A^2 + C^2 - (2AC)(\cos b)$$

$$C^2 = A^2 + B^2 - (2AB)(\cos c)$$

Trigonometric equivalencies

[] {#Trigonometric equivalencies}

$$\sin -x = -\sin x$$

$$\cos -x = \cos x$$

$$\tan -t = -\tan t$$

$$\csc -t = -\csc t$$

$$\sec -t = \sec t$$

$$\cot -t = -\cot t$$

$$\sin 2x = 2(\sin x)(\cos x)$$

$$\cos 2x = (\cos^2 x) - (\sin^2 x)$$

$$\tan 2t = \frac{2(\tan x)}{1 - \tan^2 x}$$

$$\sin^2 x = \frac{1}{2} - \frac{\cos 2x}{2}$$

$$\cos^2 x = \frac{1}{2} + \frac{\cos 2x}{2}$$

Hyperbolic functions

[] {#Hyperbolic functions} [] {#Unit, radian}

$$\sinh x = \frac{e^x - e^{-x}}{2}$$

$$\cosh x = \frac{e^x + e^{-x}}{2}$$

$$\tanh x = \frac{\sinh x}{\cosh x}$$

Note: all angles (x) must be expressed in units of *radians* for these hyperbolic functions. There are 2π radians in a circle (360°).

Contributors

Contributors to this chapter are listed in chronological order of their contributions, from most recent to first. See Appendix 2 (Contributor List) for dates and contact information.

Harvey Lew (??? 2003): Corrected typographical error: "tangent" should have been "cotangent".