TEM_UW_APP

Technical description

Lehigh TEM Felzer Lab

November 2026

Contents

1 Overview **2**

2 What files are what **2**

3 Setup **2**

4 Running a case **2**

5 Configuration **3**

6 Input data **3**

7 Test **4**

8 Documentation **4**

9 Limits **4**

10 Global input data **4**
10.1 Sources 5
10.2 Ozone 6
10.3 Nitrogen deposition 6
10.4 Land cover 6
10.5 References 7

11 Converting old files **7**
11.1 Control and parameter files 7
11.2 ASCII inputs 7

1 Overview

Prepares and runs a case of TEM_2026. The app regrids gridded NetCDF data to the model grid, writes the NetCDF inputs and `tem_config.yaml` of the model, and runs or submits `xtem`. It also builds the global input files and converts the files of the older TEM versions. The earlier CSV/XML workflow is not supported; its last version is the commit before this one.

2 What files are what

Path	Content
`case_submit.sh`	entry point, run from the case directory
`config/run_config.yaml`	all keys with defaults and comments
`src/driver.py`, `tem_climate.py`, `tem_config.py`	case builder: regridding, units, `tem_config.yaml`
`src/make_global_inputs.py`	downloads the global datasets and builds `co2.nc`, `static.nc`, `lulc.nc`, `flash.nc` and `popdens.nc` for any region
`src/xml2yaml.py`, `src/ascii2nc.py`	convert old XML, parameter and ASCII files
`launchers/`	local and SLURM launchers
`examples/west/`	western United States example
`docs/`	`global_data.md` (sources of the input data), `converting.md`
`tools/make_docs.py`	builds the two PDFs

3 Setup

```
conda env create -f env.yaml && conda activate tem_uw_app
git clone https://github.com/Lehigh-TEM-Felzer-Lab/TEM_2026
make -C TEM_2026 -j4                  # serial: build/xtem
make -C TEM_2026 mpi -j4              # MPI, for n_cpus > 1: build_mpi/xtem
export TEM_UW_APP_ROOT=$PWD
```

4 Running a case

1. Copy `examples/west/run_config.yaml` to an empty directory.
2. Set `tem_src` (the TEM_2026 checkout), the climate files, the box and the years. For `n_cpus > 1` set `tem_exe: "{tem_src}/build_mpi/xtem"`.
3. Run `$TEM_UW_APP_ROOT/case_submit.sh` from that directory. With `run_mode: slurm` it submits with sbatch; with `run_model: false` it only builds the case.

```
run_config.yaml   tem_config.yaml
INPUT/            climate.nc static.nc lulc.nc co2.nc (lightning.nc population.nc)
OUTPUT/           tem.nc, written by the model
logs/
```

5 Configuration

`config/run_config.yaml` lists every key; keys you do not set take its values and unknown keys are an error.

Key	Description
`tem_src`, `tem_exe`	TEM_2026 checkout and executable (default `{tem_src}/build/xtem`)
`netcdf_climate_in_file`, `netcdf_climate_in_frequency`	climate files (path, glob or list) and their time step (hourly, daily, monthly)
`netcdf_static_in_file`, `netcdf_lulc_in_file`, `netcdf_co2_in_file`	static, land-cover and CO2 files in the layout of TEM_2026 `docs/io_structure.md`
`netcdf_fire_in_file`, `netcdf_population_in_file`	lightning flashes and population density, needed when `veg_fire: 1`
`varmap`	`<model variable>: <variable in your files>`; the model name is used first, the map only when the file lacks it
`horizontal_resolution`, `lon_min` to `lat_max`, `regrid_method`	model grid and regridding (`box_mean`, `bilinear`, `nearest`)
`spinflag`, `veg_fire`, `xnfertflag`, ...	the options of `tem_config.yaml`, one key each
`cmntecd`, `vegecd`, ...	parameter files, by default `{tem_src}/params/biogeography/`
`clmoutvars`, `temoutvars`	variables written to `OUTPUT/tem.nc`
`run_mode`, `n_nodes`, `n_cpus`, `walltime`, `partition`, `mem`, `sbatch_options`	sbatch settings as in pyUFS; the cells are shared between the `n_cpus` MPI tasks

Climate units are read from the `units` attribute and converted. `tmin` and `tmax` can replace `tair` and `dtr`, and `dewpoint` or `huss` can replace `vpr`. Daily and hourly data are averaged to months.

6 Input data

`climate.nc` is made by the app from your climate files. The other inputs come from `src/make_global_inputs.py`:

```
A="--out INPUT --cache downloads --lon-min -125 --lon-max -110 --lat-min 32 --lat-max 49 --res 0.5"
python src/make_global_inputs.py $A co2 --start 2015 --end 2024
python src/make_global_inputs.py $A soil
python src/make_global_inputs.py $A elev
python src/make_global_inputs.py $A wind
python src/make_global_inputs.py $A static
python src/make_global_inputs.py $A lulc --luh2 states.nc --luh2-static staticData_quarterdeg.nc --year 2015
```

Ozone needs a CEDA token (`CEDA_TOKEN`); all other files are downloaded by their subcommand (`lightning`, `population`, `ndep`, ...). Sources, download links, units and the land-cover assumptions are in docs/global_data.md. Files of older TEM versions are converted as in docs/converting.md. Then set the `netcdf_*_in_file` keys of `run_config.yaml` to the files.

7 Test

`python tools/run_test.py` is one file that clones TEM_2026 and TEM_UW_APP, builds the model, downloads ERA5 (needs `~/.cdsapirc` and `cdsapi`) and the other input data, makes a global 1 degree case for one year and runs it monthly and daily. Flags such as `--competition`, `--fire`, `--all` add cases; `python tools/run_test.py -h` lists them and the docstring describes the work directory, the archive and the placeholders for ozone and N deposition.

8 Documentation

File	Content
run_config_Nov2026.pdf	every key of `run_config.yaml` with default and meaning
TEM_UW_APP-Technical-Description.pdf	this README and the documents in `docs/`

`python tools/make_docs.py` rebuilds both (needs pandoc and xelatex).

9 Limits

- The app writes monthly or daily forcing (`run_dt_units` is `month` or `day`; daily needs daily forcing, `netcdf_climate_in_frequency` set to `daily`, and a TEM_2026 build from `make daily`). Daily physics is not validated.
- Tested on the western United States example only; the inputs the app writes for it are identical to those shipped with TEM_2026.
- The global land-cover builder is not validated against the lab's earlier inputs.

10 Global input data

`src/make_global_inputs.py` writes the input files of TEM_2026 for any box and resolution (run it from the root of TEM_UW_APP). The commands below are for a 0.5 degree box over the western United States. Needs `pip install xarray netcdf4 pandas scipy rioxarray` and `wget`.

```
A="--out INPUT --cache downloads --lon-min -125 --lon-max -110 --lat-min 32 --lat-max 49 --res 0.5"
python src/make_global_inputs.py $A co2 --start 2015 --end 2024
python src/make_global_inputs.py $A soil
python src/make_global_inputs.py $A elev
python src/make_global_inputs.py $A wind
python src/make_global_inputs.py $A lightning
python src/make_global_inputs.py $A population
python src/make_global_inputs.py $A ndep --start 2005 --end 2014
CEDA_TOKEN=<token> python src/make_global_inputs.py $A ozone
python src/make_global_inputs.py $A static
python src/make_global_inputs.py $A lulc --luh2 states.nc --luh2-static staticData_quarterdeg.nc --year 2015
```

Every step downloads its source into `--cache` once. `tools/run_test.py` runs all of them for the globe at 1 degree. `static` merges the elevation, soil, wind, ozone and N deposition files into `static.nc`. `climate.nc` is made by the app from ERA5, MACA, gridMET or CMIP output. The regridding is the mean of the source cells in a target cell, so the target should be coarser than the source.

10.1 Sources

File	Variable (unit)	Source
`co2.nc`	`co2` (ppmv)	NOAA GML Mauna Loa annual mean; SSP scenarios for later years (Meinshausen et al. 2020)
`flash.nc`	`lightning_flashes` (km-2 month-1)	LIS/OTD 0.5 degree monthly climatology HRMC v2.3.2015 (Cecil et al. 2014), flash rate per day times the days of the month
`popdens.nc`	`population_density` (km-2)	WorldPop 2020 persons per 1 km pixel (WorldPop 2018), summed in each cell and divided by the cell area
`static.nc`	`elev` (m)	ETOPO 2022, 60 arc-second ice surface (NOAA NCEI 2022)
	`sand`, `silt`, `clay` (percent)	SoilGrids 2.0 at 5 km (Poggio et al. 2021), layers 0-5, 5-15 and 15-30 cm weighted by thickness, g kg-1 divided by 10
	`wind` (m s-1, by month)	WorldClim 2.1 mean wind 1970-2000 (Fick and Hijmans 2017); the height is 2 m, not 10 m
	`ozone` (AOT40, by month)	Gridded surface ozone metrics v2.7 from CEDA (Evans and Sofen 2016; Sofen et al. 2016), needs a token
	`ndep` (mg N m-2 yr-1)	ISIMIP3b NHx plus NOy deposition, mean of 2005 to 2014 (Yang et al. 2026), g N m-2 month-1 times 1000
`lulc.nc`	cohorts of each cell	LUH2 v2h (Hurtt et al. 2020); potential vegetation of Ramankutty and Foley (1999)

Sources, which the script downloads itself except LUH2 (`tools/run_test.py` fetches that with wget):

```
wget https://gml.noaa.gov/webdata/ccgg/trends/co2/co2_annmean_mlo.txt
S=https://files.isric.org/soilgrids/latest/data_aggregated/5000m
wget $S/sand/sand_0-5cm_mean_5000.tif      # also silt, clay; 5-15 and 15-30 cm
N=https://www.ngdc.noaa.gov/thredds/fileServer/global/ETOPO2022/60s/60s_surface_elev_netcdf
wget $N/ETOPO_2022_v1_60s_N90W180_surface.nc
wget https://geodata.ucdavis.edu/climate/worldclim/2_1/base/wc2.1_10m_wind.zip
wget https://sage-public-files.s3.amazonaws.com/global-potential-vegetation/potveg_nc.tar.gz
wget https://ghrc.nsstc.nasa.gov/pub/lis/climatology/LIS-OTD/HRMC/data/nc/LISOTD_HRMC_V2.3.2015.nc
wget https://data.worldpop.org/GIS/Population/Global_2000_2020/2020/0_Mosaicked/ppp_2020_1km_Aggregated.tif
I=https://files.isimip.org/ISIMIP3b/InputData/socioeconomic/n-deposition/histsoc
wget $I/ndep-nhx_histsoc_monthly_1850_2014.nc    # also ndep-noy
wget https://luh.umd.edu/LUH2/LUH2_v2h/states.nc                    # 5.8 GB
wget https://luh.umd.edu/LUH2/LUH2_v2h/staticData_quarterdeg.nc
```

The `ndep` step (downloaded file, units and the 1 degree result), the `lightning` step on a synthetic file and the `population` step on a synthetic raster were tested, as were the CO2, soil, elevation and wind steps. The LIS/OTD file could not be reached from the test machine, so its variable name (`HRMC_COM_FR`), its dimensions and its unit (flashes km-2 day-1) are assumed from the dataset description; if the step fails, download the file and pass it with `--file` (and `--var`). The WorldPop step was not run on the full 830 MB raster. The LUH2 step was tested on a synthetic file, not on the downloaded `states.nc`. Scenario CO2 after the observed years is not downloaded; pass a file with one `year ppmv` pair per line to `co2 --scenario` (see https://esgf-node.llnl.gov/projects/input4mips/).

10.2 Ozone

The ozone file needs a CEDA account (https://help.ceda.ac.uk/article/39-ceda-account) and an access token (https://services.ceda.ac.uk/account/token/; valid for 3 days). With the token in `CEDA_TOKEN`, `ozone` downloads the monthly 1 degree file of the dataset record https://catalogue.ceda.ac.uk/uuid/c0455ab814224a05ac74642d3d44b73e/ (archive directory http://data.ceda.ac.uk/badc/proc-earth-model/data/ozone-metrics/v2.7) and uses the variable whose name contains `aot40`. Pass `--file` and `--var` for a file you downloaded. The file name, variable and unit of AOT40 were not checked, because the download needs a login. The script averages a time axis into a 12-month climatology; check the result. The data come from monitoring stations, so cells without stations are missing and TEM skips cells with missing data.

10.3 Nitrogen deposition

`ndep` adds the NHx and NOy files of ISIMIP3b (https://doi.org/10.48364/ISIMIP.600567.1, unit g N m-2 month-1, 1850 to 2014), sums the months of `--start` to `--end` (default 2005 to 2014), averages the years and multiplies by 1000 to get mg N m-2 yr-1. The file carries the unit per N, so no further conversion is needed. For other data pass `--file`, `--var` and `--scale`. Another source is the ACCMIP multi-model deposition of Lamarque et al. (2013) (https://doi.org/10.5194/acp-13-7997-2013).

10.4 Land cover

The script writes up to three cohorts per cell from the LUH2 fractions of one year and the cell area:

Cohort	LUH2 states	`currentveg`	`agstate`	`standage`
1 natural	`primf`, `primn`, `secdf`, `secdn`, `range`	potential vegetation of the cell	0	3000
2 crops	`c3ann`, `c4ann`, `c3per`, `c4per`, `c3nfx`	50 (Crops)	1	1
3 pasture	`pastr`	51 (Pasture)	1	1

`chrtarea` is the area in km2, `potveg` the TEM type of the potential vegetation, `subtype` equals `currentveg` and `FRI` is 2000. A cohort exists where its area exceeds 0.001 km2. The model counts the cohorts of each cell from the cells where `standage` has data. `cohort`, `potveg`, `currentveg` and `agstate` carry `flag_values` and `flag_meanings` that list the TEM vegetation types by name. Urban land is not simulated.

The 15 classes of Ramankutty and Foley map to TEM types (`params/biogeography/tcomm50_lulc.yaml` of TEM_2026) as 1 to 16, 2 to 18, 3 to 33, 4 to 9, 5 to 10, 6 and 7 to 4, 8 to 8, 9 to 14, 10 to 13, 11 and 12 to 15, 13 to 3, 14 to 21 and 15 to 2. This crosswalk, the `subtype` rule, `standage` of the managed cohorts and `agstate` of pasture are assumptions of the script; the model developers should confirm them before a production run. Management layers (`tillflag`, `fertflag`, `irrgflag`) and land-cover change in time are not written, so a run uses `tlulcflag: 0`.

`lulc.nc` has `cohort` as dimension and as int64 coordinate; `lat` and `lon` are float32, categorical fields int64 with `_FillValue -9999`, other fields float32. The layouts of all files are in `docs/io_structure.md` of TEM_2026.

10.5 References

- Cecil, D. J., Buechler, D. E. and Blakeslee, R. J. (2014) Atmos. Res. 135-136, 404-414. https://doi.org/10.1016/j.atmosres.2012.06.028
- Evans, M. J. and Sofen, E. D. (2016) Gridded Global Surface Ozone Metrics data (1971-2015), version 2.7. CEDA. https://doi.org/10.5285/897a3958-5bfc-4311-9bb0-01134bf6aefa
- Fick, S. E. and Hijmans, R. J. (2017) Int. J. Climatol. 37, 4302-4315. https://doi.org/10.1002/joc.5086
- Hurtt, G. C. et al. (2020) Geosci. Model Dev. 13, 5425-5464. https://doi.org/10.5194/gmd-13-5425-2020
- Lamarque, J.-F. et al. (2013) Atmos. Chem. Phys. 13, 7997-8018. https://doi.org/10.5194/acp-13-7997-2013
- Meinshausen, M. et al. (2020) Geosci. Model Dev. 13, 3571-3605. https://doi.org/10.5194/gmd-13-3571-2020
- NOAA NCEI (2022) ETOPO 2022 15 Arc-Second Global Relief Model. https://doi.org/10.25921/fd45-gt74
- Poggio, L. et al. (2021) SOIL 7, 217-240. https://doi.org/10.5194/soil-7-217-2021
- Ramankutty, N. and Foley, J. A. (1999) Global Biogeochem. Cycles 13, 997-1027. https://doi.org/10.1029/1999GB900046
- Sofen, E. D. et al. (2016) Earth Syst. Sci. Data 8, 41-59. https://doi.org/10.5194/essd-8-41-2016
- WorldPop (www.worldpop.org) and CIESIN, Columbia University (2018) Global High Resolution Population Denominators Project. University of Southampton. https://doi.org/10.5258/SOTON/WP00647
- Yang, J., Tian, H. and Büchner, M. (2026) ISIMIP3b Simulated Historical Nitrogen Deposition Input Data, v1.1. ISIMIP Repository. https://doi.org/10.48364/ISIMIP.600567.1

11 Converting old files

For runs set up with the older TEM versions (TEM_Monthly, TEM_Biogeography, GDSTEM and others).

11.1 Control and parameter files

```
python src/xml2yaml.py us_tem45_lulc.xml -o tem_config.yaml
python src/xml2yaml.py tcomm45_lulc.ecd -o tcomm50_lulc.yaml
```

`xml2yaml.py` turns an XML control file or an `.ecd`/`.dat` parameter file into YAML. Every tag becomes a key and the XML comments become YAML comments. It repairs the known typos of the old files (a mismatched closing tag, an unclosed quote) and lists the ASCII input names it finds, since TEM_2026 reads NetCDF through the `netcdf_*_in_file` keys. The other keys keep their old names; compare the result with `templates/tem_config.yaml` of TEM_2026.

11.2 ASCII inputs

```
python src/ascii2nc.py clm tair.csv prec.csv dtr.csv nirr.csv vpr.csv -o climate.nc
python src/ascii2nc.py clm wind.csv aot40.csv -o static.nc --grid climate.nc
python src/ascii2nc.py elev elev.csv -o static.nc --grid climate.nc
python src/ascii2nc.py soil soil.csv -o static.nc --grid climate.nc
python src/ascii2nc.py ndep ndep.csv -o static.nc --grid climate.nc
python src/ascii2nc.py co2 co2.csv -o co2.nc
python src/ascii2nc.py lulc lulc.csv -o lulc.nc --grid climate.nc
```

Repeated calls with the same `-o` add to the file. The kinds and the record layouts are listed at the top of `src/ascii2nc.py`:

Kind	Record
`clm`	lon, lat, VAR, area, year, total, max, ave, min, jan to dec, region (year -999: climatology by month)
`daily`	lon, lat, VAR, area, year, month, day, value, region
`elev`	lon, lat, ELEV, area, elev, region
`soil`	lon, lat, SOIL, area, sand, silt, clay, wsoil, source, region
`ndep`	lon, lat, year, value
`co2`	year, ppmv

Kind	Record
`lulc`	lon, lat, VAR, year, cohort, srccohort, standage, chrtarea, potveg, currentveg, subtype, agstate, agprevstate, tillflag, fertflag, irrgflag, disturbflag, disturbmonth, FRI, slashpar, vconvert, prod10par, prod100par, vrespar, sconvert, region

Variables are renamed as TEM_2026 expects (`CLDINESS` to `clouds`, `DTAIRRNG` to `dtr`, `VPRPRESS` to `vpr`, `AOT40` to `ozone`). The model counts the cohorts from `lulc.nc`, so the old cohort-count file (`MXCOHRTS`) is not needed. The old files list only cells with data: `--grid` takes a NetCDF file of the full grid, usually the climate file, and restores the empty cells.

Checked by writing the example inputs of TEM_2026 as ASCII, converting them back (equal to the originals within the ASCII rounding) and running the model on the result (maximum GPP 423.98 against 424.01). The `daily` and `ndep` kinds were not tested. The old ASCII fire inputs and the old output files are not converted; `tem_utils` converts old HIST output.