

```

> restart: unprotect('D'):

m := 1;

twist := 0;

d := 6;                      degree := d + 1:

##

SS := x^(d+1) + 1^(d+1) + 0*x^6:
HSS := X^(d+1) + T^(d+1) + 0*X^6:

#SS := x^(d+1) + 1^(d+1) + 1*x^6:
#HSS := X^(d+1) + T^(d+1) + 1*T^(d+1-6)*X^6:

##

Jet_xR := expand(
+ (1/Ry^(m-0*2)) * add(AA||j * x1^(m-0*3-j) * (R1/RR)^j *
Delta_xR^0, j=0..m-0*3)
+ (1/Ry^(m-1*2)) * add(BB||j * x1^(m-1*3-j) * (R1/RR)^j *
Delta_xR^1, j=0..m-1*3)
+ (1/Ry^(m-2*2)) * add(CC||j * x1^(m-2*3-j) * (R1/RR)^j *
Delta_xR^2, j=0..m-2*3)
+ (1/Ry^(m-3*2)) * add(DD||j * x1^(m-3*3-j) * (R1/RR)^j *
Delta_xR^3, j=0..m-3*3)
+ (1/Ry^(m-4*2)) * add(EE||j * x1^(m-4*3-j) * (R1/RR)^j *
Delta_xR^4, j=0..m-4*3) ):

##

for l from 0 to m - 0*3 do dA[l] := (m-0*2)*d - m + l*1 + 0*1 +
twist: od:

for l from 0 to m - 1*3 do dB[l] := (m-1*2)*d - m + l*1 + 1*1 +
twist: od:

for l from 0 to m - 2*3 do dC[l] := (m-2*2)*d - m + l*1 + 2*1 +
twist: od:

for l from 0 to m - 3*3 do dD[l] := (m-3*2)*d - m + l*1 + 3*1 +
twist: od:

for l from 0 to m - 4*3 do dE[l] := (m-4*2)*d - m + l*1 + 4*1 +
twist: od:

##

```

```

Delta_xR := x1 * (R2/RR - R1*R1/RR^2) - x2 * (R1/RR):
Solx1 := - y1*Ry/Rx + R1/Rx:
Solx2 := - Rxx*y1^2*Ry^2/Rx^3 + 2*Rxx*y1*Ry*R1/Rx^3 - Rxx*
R1^2/Rx^3 + 2*y1^2*Rxy*Ry/Rx^2
          - 2*y1*Rxy*R1/Rx^2 - y1^2*Ryy/Rx - y2*Ry/Rx + R2/Rx:
##
Jet_Ry := sort(mttaylor(
          expand(subs(R2=-(RR^2*Delta_Ry-RR*R1*y2-
R1^2*y1)/(RR*y1),
          expand(subs({x1=Solx1,x2=Solx2},
Jet_xR))),
          [R1,y1,Delta_Ry], 100), [R1,y1,Delta_Ry]):
##
printlevel := 2:
for k from 0 to m/3 do
for j from 0 to m-3*k do
EEq[m-j-3*k,j,k] := factor(
  Ry^(m-j-3*k)*RR^(m-j-3*k)*Rx^(m-2*k)
*
  coeftayl(Jet_Ry, [R1,y1,Delta_Ry]=[0,0,0], [m-j-3*k,j,k]) ):
od: od:
## EQUATION RR EXPLICITE
RR := y^(d+1) + SS;
HRR := Y^(d+1) + HSS;
RY := diff(HRR, Y):
Rx := diff(RR,x):
Ry := diff(RR,y):
Rxx := diff(RR,x,x):
Rxy := diff(RR,x,y):
Ryy := diff(RR,y,y):
##
printlevel := 2:
for k from 0 to m/3 do
for j from 0 to m-3*k do
Eq[m-j-3*k,j,k] := mttaylor(EEq[m-j-3*k,j,k], y, (m-j-3*k)*d):
Var[m-j-3*k,j,k] := indets(Eq[m-j-3*k,j,k]) minus {x,y}:

```

od: od:

##

$m := 1$

$twist := 0$

$d := 6$

$RR := x^7 + y^7 + 1$

$HRR := T^7 + X^7 + Y^7$

(1)

> ## EXPAND ALL UNKNOWNNS UP TO $y^{(1*d)}$

for l from 0 to m - 0*3 do
AA[l] := add(add(A[l][i,j]*xⁱ*y^j, i=0..dA[l]-j), j=0..min(dA[l], 1*d-1)): od:

for l from 0 to m - 1*3 do
BB[l] := add(add(B[l][i,j]*xⁱ*y^j, i=0..dB[l]-j), j=0..min(dB[l], 1*d-1)): od:

for l from 0 to m - 2*3 do
CC[l] := add(add(C[l][i,j]*xⁱ*y^j, i=0..dC[l]-j), j=0..min(dC[l], 1*d-1)): od:

for l from 0 to m - 3*3 do
DD[l] := add(add(D[l][i,j]*xⁱ*y^j, i=0..dD[l]-j), j=0..min(dD[l], 1*d-1)): od:

for l from 0 to m - 4*3 do
EE[l] := add(add(E[l][i,j]*xⁱ*y^j, i=0..dE[l]-j), j=0..min(dE[l], 1*d-1)): od:

SYSTEM FOR DIVISIBILITY BY $y^{(1*d)}$

Systemey1d := {}:

for k from 0 to (m-1)/3 do

Coeffs_y1d[1,m-1-3*k,k] := {coeffs(expand(mtaylor(Eq[1,m-1-3*k,k], y, 1*d)), [x,y])}:

Systemey1d := Systemey1d union Coeffs_y1d[1,m-1-3*k,k]:

od:

Systeme_y1d := Systemey1d:

VARIABLES FOR DIVISIBILITY BY $y^{(1*d)}$

Variables_y1d := indets(Systeme_y1d):

```
## RESOLUTION  $y^{(1*d)}$ 
```

```
Solutions_y1d := solve(Systeme_y1d, Variables_y1d):
```

```
assign(Solutions_y1d):
```

```
##
```

```
for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;
```

```
l := 0
```

```
1
```

```
l := 1
```

```
1
```

```
l := 2
```

```
l := 0
```

```
l := 0
```

```
l := 0
```

```
l := 0
```

(2)

```
> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(2*d)}$ 
```

```
for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],2*d-1)): od:
```

```
for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l],2*d-1)): od:
```

```
for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l],2*d-1)): od:
```

```
for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],2*d-1)): od:
```

```
for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],2*d-1)): od:
```

```
## SYSTEM FOR DIVISIBILITY BY  $y^{(2*d)}$ 
```

```

Systemey2d := {}:
for k from 0 to (m-2)/3 do
  Coeffs_y2d[2,m-2-3*k,k] := {coeffs(expand(mtaylor(Eq[2,m-2-3*k,
k], y, 2*d)), [x,y])}:
Systemey2d := Systemey2d union Coeffs_y2d[2,m-2-3*k,k]:
od:
Systeme_y2d := Systemey2d:
## VARIABLES FOR DIVISIBILITY BY y^(2*d)
Variables_y2d := indets(Systeme_y2d):
## RESOLUTION y^(2*d)
Solutions_y2d := solve(Systeme_y2d, Variables_y2d):
assign(Solutions_y2d):
##
for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

(3)

```

> ## EXPAND ALL UNKNOWNNS UP TO y^(3*d)
for l from 0 to m - 0*3 do
  AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],3*d-1)): od:
for l from 0 to m - 1*3 do
  BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB

```

```

[l], 3*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l], 3*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l], 3*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l], 3*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY y^(3*d)

Systemey3d := {}:

for k from 0 to (m-3)/3 do

Coeffs_y3d[3,m-3-3*k,k] := {coeffs(expand(mtaylor(Eq[3,m-3-3*k,
k], y, 3*d)), [x,y])}:

Systemey3d := Systemey3d union Coeffs_y3d[3,m-3-3*k,k]:

od:

Systeme_y3d := Systemey3d:

## VARIABLES FOR DIVISIBILITY BY y^(3*d)

Variables_y3d := indets(Systeme_y3d):

## RESOLUTION y^(3*d)

Solutions_y3d := solve(Systeme_y3d, Variables_y3d):

assign(Solutions_y3d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0

```

$l := 0$

$l := 0$

$l := 0$

(4)

```
> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(4*d)}$ 

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA[l],4*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB[l],4*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC[l],4*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD[l],4*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE[l],4*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY  $y^{(4*d)}$ 

Systemey4d := {}:

for k from 0 to (m-4)/3 do

Coeffs_y4d[4,m-4-3*k,k] := {coeffs(expand(mTaylor(Eq[4,m-4-3*k,
k], y, 4*d)), [x,y])}:

Systemey4d := Systemey4d union Coeffs_y4d[4,m-4-3*k,k]:

od:

Systeme_y4d := Systemey4d:

## VARIABLES FOR DIVISIBILITY BY  $y^{(4*d)}$ 

Variables_y4d := indets(Systeme_y4d):

## RESOLUTION  $y^{(4*d)}$ 

Solutions_y4d := solve(Systeme_y4d, Variables_y4d):

assign(Solutions_y4d):

##
```

```

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

```

```
l := 0
```

```
1
```

```
l := 1
```

```
2
```

```
l := 2
```

```
l := 0
```

```
l := 0
```

```
l := 0
```

```
l := 0
```

(5)

```
> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(5*d)}$ 
```

```

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA[l],5*d-1)): od:

```

```

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB[l],5*d-1)): od:

```

```

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC[l],5*d-1)): od:

```

```

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD[l],5*d-1)): od:

```

```

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE[l],5*d-1)): od:

```

```
## SYSTEM FOR DIVISIBILITY BY  $y^{(5*d)}$ 
```

```
Systemey5d := {}:
```

```
for k from 0 to (m-5)/3 do
```

```

  Coeffs_y5d[5,m-5-3*k,k] := {coeffs(expand(mTaylor(Eq[5,m-5-3*k,
k], y, 5*d)), [x,y])}:

```

```
Systemey5d := Systemey5d union Coeffs_y5d[5,m-5-3*k,k]:
```



```

od:
Systeme_y5d := Systemey5d:
## VARIABLES FOR DIVISIBILITY BY  $y^{(5*d)}$ 
Variables_y5d := indets(Systeme_y5d):
## RESOLUTION  $y^{(5*d)}$ 
Solutions_y5d := solve(Systeme_y5d, Variables_y5d):
assign(Solutions_y5d):
##
for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

```

$l := 0$

1

$l := 1$

2

$l := 2$

$l := 0$

$l := 0$

$l := 0$

$l := 0$

(6)

```

> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(6*d)}$ 
for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA[l],6*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB[l],6*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC[l],6*d-1)): od:

for l from 0 to m - 3*3 do

```

```

DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],6*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],6*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY y^(6*d)

Systemey6d := {}:

for k from 0 to (m-6)/3 do

Coeffs_y6d[6,m-6-3*k,k] := {coeffs(expand(mtaylor(Eq[6,m-6-3*k,
k], y, 6*d)), [x,y])}:

Systemey6d := Systemey6d union Coeffs_y6d[6,m-6-3*k,k]:

od:

Systeme_y6d := Systemey6d:

## VARIABLES FOR DIVISIBILITY BY y^(6*d)

Variables_y6d := indets(Systeme_y6d):

## RESOLUTION y^(6*d)

Solutions_y6d := solve(Systeme_y6d, Variables_y6d):

assign(Solutions_y6d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

```

> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(7*d)}$ 

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l], 7*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l], 7*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l], 7*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l], 7*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l], 7*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY  $y^{(7*d)}$ 

Systemey7d := {}:

for k from 0 to (m-7)/3 do

Coeffs_y7d[7,m-7-3*k,k] := {coeffs(expand(mtaylor(Eq[7,m-7-3*k,
k], y, 7*d)), [x,y])}:

Systemey7d := Systemey7d union Coeffs_y7d[7,m-7-3*k,k]:

od:

Systeme_y7d := Systemey7d:

## VARIABLES FOR DIVISIBILITY BY  $y^{(7*d)}$ 

Variables_y7d := indets(Systeme_y7d):

## RESOLUTION  $y^{(7*d)}$ 

Solutions_y7d := solve(Systeme_y7d, Variables_y7d):

assign(Solutions_y7d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

```

$l := 0$

```

1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

(8)

```

>
## EXPAND ALL UNKNOWNNS UP TO y^(8*d)

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],8*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l],8*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l],8*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],8*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],8*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY y^(8*d)

Systemey8d := {}:

for k from 0 to (m-8)/3 do

Coeffs_y8d[8,m-8-3*k,k] := {coeffs(expand(mtaylor(Eq[8,m-8-3*k,
k], y, 8*d)), [x,y])}:

Systemey8d := Systemey8d union Coeffs_y8d[8,m-8-3*k,k]:

od:

Systeme_y8d := Systemey8d:

## VARIABLES FOR DIVISIBILITY BY y^(8*d)

Variables_y8d := indets(Systeme_y8d):

```

```

## RESOLUTION  $y^{(8*d)}$ 

Solutions_y8d := solve(Systeme_y8d, Variables_y8d):

assign(Solutions_y8d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

(9)

```

> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(9*d)}$ 

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA[l],9*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB[l],9*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC[l],9*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD[l],9*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE[l],9*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY  $y^{(9*d)}$ 

Systemey9d := {}:

```

```

for k from 0 to (m-9)/3 do
  Coeffs_y9d[9,m-9-3*k,k] := {coeffs(expand(mtaylor(Eq[9,m-9-3*k,
k], y, 9*d)), [x,y])}:
  Systemey9d := Systemey9d union Coeffs_y9d[9,m-9-3*k,k]:
od:
Systeme_y9d := Systemey9d:
## VARIABLES FOR DIVISIBILITY BY y^(9*d)
Variables_y9d := indets(Systeme_y9d):
## RESOLUTION y^(9*d)
Solutions_y9d := solve(Systeme_y9d, Variables_y9d):
assign(Solutions_y9d):
##
for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

(10)

```

> ## EXPAND ALL UNKNOWNNS UP TO y^(10*d)
for l from 0 to m - 0*3 do
  AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],10*d-1)): od:
for l from 0 to m - 1*3 do
  BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l],10*d-1)): od:

```

```

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l],10*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],10*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],10*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY y^(10*d)

Systemey10d := {}:

for k from 0 to (m-10)/3 do

Coeffs_y10d[10,m-10-3*k,k] := {coeffs(expand(mtaylor(Eq[10,m-10
-3*k,k], y, 10*d)), [x,y])}:

Systemey10d := Systemey10d union Coeffs_y10d[10,m-10-3*k,k]:

od:

Systeme_y10d := Systemey10d:

## VARIABLES FOR DIVISIBILITY BY y^(10*d)

Variables_y10d := indets(Systeme_y10d):

## RESOLUTION y^(10*d)

Solutions_y10d := solve(Systeme_y10d, Variables_y10d):

assign(Solutions_y10d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0

```

```

>
## EXPAND ALL UNKNOWNNS UP TO  $y^{(11*d)}$ 

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],11*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l],11*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l],11*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],11*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],11*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY  $y^{(11*d)}$ 

Systemey11d := {}:

for k from 0 to (m-11)/3 do

Coeffs_y11d[11,m-11-3*k,k] := {coeffs(expand(mtaylor(Eq[11,m-11
-3*k,k], y, 11*d)), [x,y])}:

Systemey11d := Systemey11d union Coeffs_y11d[11,m-11-3*k,k]:

od:

Systeme_y11d := Systemey11d:

## VARIABLES FOR DIVISIBILITY BY  $y^{(11*d)}$ 

Variables_y11d := indets(Systeme_y11d):

## RESOLUTION  $y^{(11*d)}$ 

Solutions_y11d := solve(Systeme_y11d, Variables_y11d):

assign(Solutions_y11d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od:

```



```

for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

```

```
l := 0
```

```
1
```

```
l := 1
```

```
2
```

```
l := 2
```

```
l := 0
```

```
l := 0
```

```
l := 0
```

```
l := 0
```

(12)

```
> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(12*d)}$ 
```

```

for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],12*d-1)): od:

```

```

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l],12*d-1)): od:

```

```

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l],12*d-1)): od:

```

```

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],12*d-1)): od:

```

```

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],12*d-1)): od:

```

```
## SYSTEM FOR DIVISIBILITY BY  $y^{(12*d)}$ 
```

```
Systemey12d := {}:
```

```
for k from 0 to (m-12)/3 do
```

```

Coeffs_y12d[12,m-12-3*k,k] := {coeffs(expand(mtaylor(Eq[12,m-12
-3*k,k], y, 12*d)), [x,y])}:

```

```
Systemey12d := Systemey12d union Coeffs_y12d[12,m-12-3*k,k]:
```

```
od:
```

```

Systeme_y12d := Systemey12d:
## VARIABLES FOR DIVISIBILITY BY  $y^{(12*d)}$ 
Variables_y12d := indets(Systeme_y12d):
## RESOLUTION  $y^{(12*d)}$ 
Solutions_y12d := solve(Systeme_y12d, Variables_y12d):
assign(Solutions_y12d):
##
for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

(13)

```

> ## EXPAND ALL UNKNOWNNS UP TO  $y^{(13*d)}$ 
for l from 0 to m - 0*3 do
AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA[l],13*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB[l],13*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC[l],13*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD[l],13*d-1)): od:

for l from 0 to m - 4*3 do

```

```

EE[l] := add(add(E[l][i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE[l], 13*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY y^(13*d)

Systemey13d := {}:

for k from 0 to (m-13)/3 do

Coeffs_y13d[13,m-13-3*k,k] := {coeffs(expand(mtaylor(Eq[13,m-13-3*k,k], y, 13*d)), [x,y])}:

Systemey13d := Systemey13d union Coeffs_y13d[13,m-13-3*k,k]:

od:

Systeme_y13d := Systemey13d:

## VARIABLES FOR DIVISIBILITY BY y^(13*d)

Variables_y13d := indets(Systeme_y13d):

## RESOLUTION y^(13*d)

Solutions_y13d := solve(Systeme_y13d, Variables_y13d):

assign(Solutions_y13d):

##

for l from 0 to m - 0*3 do nops(expand(AA[l])) od;
for l from 0 to m - 1*3 do nops(expand(BB[l])) od;
for l from 0 to m - 2*3 do nops(expand(CC[l])) od;
for l from 0 to m - 3*3 do nops(expand(DD[l])) od;
for l from 0 to m - 4*3 do nops(expand(EE[l])) od;

l := 0
1
l := 1
2
l := 2
l := 0
l := 0
l := 0
l := 0

```

(14)

```

> ## EXPAND ALL UNKNOWNNS UP TO y^(14*d)

for l from 0 to m - 0*3 do

```

```

AA||l := add(add(A||l[i,j]*x^i*y^j, i=0..dA[l]-j), j=0..min(dA
[l],14*d-1)): od:

for l from 0 to m - 1*3 do
BB||l := add(add(B||l[i,j]*x^i*y^j, i=0..dB[l]-j), j=0..min(dB
[l],14*d-1)): od:

for l from 0 to m - 2*3 do
CC||l := add(add(C||l[i,j]*x^i*y^j, i=0..dC[l]-j), j=0..min(dC
[l],14*d-1)): od:

for l from 0 to m - 3*3 do
DD||l := add(add(D||l[i,j]*x^i*y^j, i=0..dD[l]-j), j=0..min(dD
[l],14*d-1)): od:

for l from 0 to m - 4*3 do
EE||l := add(add(E||l[i,j]*x^i*y^j, i=0..dE[l]-j), j=0..min(dE
[l],14*d-1)): od:

## SYSTEM FOR DIVISIBILITY BY y^(14*d)

Systemey14d := {}:

for k from 0 to (m-14)/3 do

Coeffs_y14d[14,m-14-3*k,k] := {coeffs(expand(mtaylor(Eq[14,m-14
-3*k,k], y, 14*d)), [x,y])}:

Systemey14d := Systemey14d union Coeffs_y14d[14,m-14-3*k,k]:

od:

Systeme_y14d := Systemey14d:

## VARIABLES FOR DIVISIBILITY BY y^(14*d)

Variables_y14d := indets(Systeme_y14d):

## RESOLUTION y^(14*d)

Solutions_y14d := solve(Systeme_y14d, Variables_y14d):

assign(Solutions_y14d):

##

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

l := 0
1
l := 1

```

$$\begin{aligned}
& 2 \\
& l := 2 \\
& l := 0 \\
& l := 0 \\
& l := 0 \\
& l := 0
\end{aligned}$$

(15)

```

> ## HOLOMORPHICITY AT INFINITY
  for l from 0 to m - 0*3 do
    HA||l := add(add(A||l[i,j]*X^i*Y^j*T^(dA[l]-i-j), i=0..dA[l]-j),
j=0..dA[l]):
  od:
#
  for l from 0 to m - 1*3 do
    HB||l := add(add(B||l[i,j]*X^i*Y^j*T^(dB[l]-i-j), i=0..dB[l]-j),
j=0..dB[l]):
  od:
#
  for l from 0 to m - 2*3 do
    HC||l := add(add(C||l[i,j]*X^i*Y^j*T^(dC[l]-i-j), i=0..dC[l]-j),
j=0..dC[l]):
  od:
#
  for l from 0 to m - 3*3 do
    HD||l := add(add(D||l[i,j]*X^i*Y^j*T^(dD[l]-i-j), i=0..dD[l]-j),
j=0..dD[l]):
  od:

```

```

#
for l from 0 to m - 4*3 do
  HE||l := add(add(E||l[i,j]*X^i*Y^j*T^(dE[l]-i-j), i=0..dE[l]-j),
j=0..dE[l]):
od:
##
for j from 0 to m - 0*3 do
  HK[j,0] := HA||j:
od:
#
for j from 0 to m - 1*3 do
  HK[j,1] := HB||j:
od:
#
for j from 0 to m - 2*3 do
  HK[j,2] := HC||j:
od:
#
for j from 0 to m - 3*3 do
  HK[j,3] := HD||j:
od:
#
for j from 0 to m - 4*3 do
  HK[j,4] := HE||j:
od:

```

```

> ##

```

```

  printlevel := 2:

```

```

for q from 0 to m/3 do
for p from 0 to m-3*q do

```

```

HEq[p,q] := expand(mtaylor(add(add(add(

```

```

  (-1)^k * 2^(p-r) * (d+1)^(j+k-p-q) * binomial(j,r) * (k!)/((p-
r)!*q!*(k+r-p-q)!)

```

```

* X^(m-q-j-2*k) * T^(k-q) * RY^(2*k-2*q) * HK[j,k],

```

```

j=r..m-3*k),
r=max(0,p+q-k)..min(p,m-3*k)),
k=q..m/3)

```

```

, T, m-p-3*q)):

```

```

od: od;

```

$q := 0$

$HEq := \text{table}([])$

$HEq_{0,0} := 7 A_{0,6} Y^6$

$HEq_{1,0} := A_{0,6} Y^6$

$q := 1$

(16)

```

> HSysteme := {seq(seq(coeffs(HEq[p,q],[T,X,Y]),p=0..m-3*q),q=0..
m/3)} minus {0}:

```

```

HSolutions := solve(HSysteme):

```

```

assign(HSolutions):

```

```

##

```

```

for l from 0 to m - 0*3 do nops(expand(AA||l)) od;
for l from 0 to m - 1*3 do nops(expand(BB||l)) od;
for l from 0 to m - 2*3 do nops(expand(CC||l)) od;
for l from 0 to m - 3*3 do nops(expand(DD||l)) od;
for l from 0 to m - 4*3 do nops(expand(EE||l)) od;

```

$l := 0$

1

$l := 1$

1

$l := 2$

$l := 0$

[

$l := 0$
 $l := 0$
 $l := 0$

(17)

[> Jet_xR := factor(

$$\begin{aligned}
& + (1/Ry^{(m-0*2)}) * x^{(-twist)} * add(AA||j * x1^{(m-0*3-j)} * \\
& (R1/RR)^j * Delta_{xR}^0, j=0..m-0*3) \\
& + (1/Ry^{(m-1*2)}) * x^{(-twist)} * add(BB||j * x1^{(m-1*3-j)} * \\
& (R1/RR)^j * Delta_{xR}^1, j=0..m-1*3) \\
& + (1/Ry^{(m-2*2)}) * x^{(-twist)} * add(CC||j * x1^{(m-2*3-j)} * \\
& (R1/RR)^j * Delta_{xR}^2, j=0..m-2*3) \\
& + (1/Ry^{(m-3*2)}) * x^{(-twist)} * add(DD||j * x1^{(m-3*3-j)} * \\
& (R1/RR)^j * Delta_{xR}^3, j=0..m-3*3) \\
& + (1/Ry^{(m-4*2)}) * x^{(-twist)} * add(EE||j * x1^{(m-4*3-j)} * \\
& (R1/RR)^j * Delta_{xR}^4, j=0..m-4*3));
\end{aligned}$$

$$Jet_{xR} := 0 \quad (18)$$

```

> Vars := indets(Jet_xR) minus {x, x1, x2, y, R1, R2};
NbVars := nops(Vars);

```

Vars := ;
NbVars := 0

(19)

```

> Ens_Vars := {}:
for i from 1 to nops(Vars) do
  Ens_Vars := Ens_Vars union {op(i,Vars)=0}:
od:
EnsVars := Ens_Vars;

```

EnsVars := :

(20)

```

> for i from 1 to nops(Vars) do
  JetSol[i] := factor(subs(EnsVars, subs(op(i,Vars)=1, Jet_xR))):
od;

```

L

$i := 1$

(21)