

DOI: 10.1159/000499659

Received: 11/9/2018

Accepted: 3/17/2019

Published(online): 3/19/2019

-----  
Corticosteroid action in the brain: the potential of selective receptor modulation

Viho E.M.G. Buurstede J.C. Mahfouz A. Koorneef L.L. van Weert L.T.C.M. Houtman R. Hunt H.J.  
. Kroon J. Meijer O.C.

-----  
ISSN: 0028-3835 (Print), eISSN: 1423-0194 (Online)

<https://www.karger.com/NEN>

Neuroendocrinology  
-----

Disclaimer:

Accepted, unedited article not yet assigned to an issue. The statements, opinions and data contained in this publication are solely those of the individual authors and contributors and not of the publisher and the editor(s). The publisher and the editor(s) disclaim responsibility for any injury to persons or property resulting from any ideas, methods, instructions or products referred to in the content.

Copyright:

This article is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND) (<http://www.karger.com/Services/OpenAccessLicense>). Usage and distribution for commercial purposes as well as any distribution of modified material requires written permission.

The Author(s). Published by S. Karger AG, Basel  
-----

## **Corticosteroid action in the brain: the potential of selective receptor modulation**

Eva M.G. Viho <sup>1,2</sup>, Jacobus C. Buurstede <sup>1,2</sup>, Ahmed Mahfouz <sup>3,4</sup>, Lisa L. Koorneef <sup>1,2</sup>, Lisa T.C.M van Weert <sup>1,2</sup>, René Houtman <sup>5</sup>, Hazel J. Hunt <sup>6</sup>, Jan Kroon <sup>1,2</sup>, Onno C. Meijer <sup>1,2\*</sup>

<sup>1</sup> Department Internal Medicine, Division of Endocrinology, Leiden University Medical Center, Leiden, The Netherlands.

<sup>2</sup> Einthoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, the Netherlands.

<sup>3</sup> Delft Bioinformatics Laboratory, Delft University of Technology, Delft, The Netherlands.

<sup>4</sup> Leiden Computational Biology Center, Leiden University Medical Center, Leiden, Netherlands.

<sup>5</sup> PamGene BV, Den Bosch, The Netherlands.

<sup>6</sup> Corcept Therapeutics, Menlo Park, California 94025, USA.

Short Title: *Selective glucocorticoid receptor modulation in the brain*

\*Corresponding author

Onno C. Meijer

Department of Internal Medicine

Division of Endocrinology

Leiden University Medical Center

Albinusdreef 2

2333 ZA, Leiden The Netherlands

Tel.: +31715261109

E-mail: o.c.meijer@lumc.nl

Keywords: glucocorticoid receptor, learning and memory, nuclear receptor coregulators, neuropsychiatric diseases, neurodegenerative diseases

## Abstract

Glucocorticoid hormones have important effects on brain function in the context of acute and chronic stress. Many of these are mediated by the glucocorticoid receptor (GR). GR has a transcriptional activity which is highly context-specific and differs between tissues and even between cell types. The outcome of GR-mediated transcription depends on the interactome of associated coregulators. Selective Glucocorticoid Receptor Modulators (SGRMs) are a class of GR ligands that can be used to activate only a subset of GR-coregulator interactions, thereby giving the possibility to induce a unique combination of agonistic and antagonistic GR properties. We describe SGRM action in animal models of brain function and pathology, and argue for their utility as molecular filters, to characterize context-specific GR interactome and transcriptional activity that are responsible for particular glucocorticoid-driven effects in cognitive processes such as memory consolidation. The ultimate objective of this approach is to identify molecular processes that are responsible for adaptive and maladaptive effects of glucocorticoids in the brain.

## **Stress and glucocorticoids in brain function and memory**

### *Stress, homeostasis and the hypothalamic pituitary adrenal axis*

Stress is a state following a perceived threat to homeostasis. It involves the activation of several responsive systems, including the endocrine, nervous and immune systems, altogether facilitating adaptation of the organism to the stressor (1). More specifically, the stress response includes the central corticotropin-releasing hormone (CRH) system, the sympathetic nervous system, and the hypothalamic pituitary adrenal (HPA) axis. The latter constitutes the main neuroendocrine effector of the stress response, in which the stressor triggers activity of the hypothalamic paraventricular nucleus (PVN). This initiates a cascade of hormonal processes starting with CRH release from the PVN, which in turn triggers production and release of adrenocorticotrophic hormone (ACTH) from the anterior pituitary gland, ultimately resulting in the secretion of glucocorticoid (GC) hormones by the adrenal cortex into the circulation.

### *Glucocorticoid levels and rhythm: from adaptation to maladaptation*

The predominant GC in humans is cortisol while rodents exclusively produce corticosterone. Basal levels of GCs fluctuate as they display circadian and ultradian rhythms, which synchronize physiological processes and maintain appropriate stress responsiveness of the HPA axis and brain circuits (2-4). The acute GC responses to stress – or short-term medical treatment with synthetic GCs – have many context-dependent effects that either curtail the initial stress response, exemplified by anti-inflammatory actions, or support prolonged redistribution of energy stores and adaptation to future stressors (5, 6). Long-term and/or high exposure to endogenous and synthetic GCs is however associated with many adverse effects, including the development of metabolic diseases, osteoporosis, psychiatric symptoms and cognitive deficits (7).

The brain is a prominent target of GCs, and it constitutes the central structure for adaptation to stress. GC effects on the brain are pleiotropic – as they influence behavior, cognition, mood, and programming of the stress response – all functions to adapt to stressors. Shifts in GC levels are associated with complex changes in neuronal activity that differ over time and according to the brain region examined (4). At the cellular level, GCs are necessary for neuronal differentiation, integrity, growth, and synaptic and dendritic plasticity (8, 9). These cellular processes support brain functions such as decision-making, reward-based behavior,

motor control, visual information processing, learning and memory, food intake and energy regulation.

The inter-connected limbic structures that mediate effects of stress and influence the HPA axis through PVN projecting neurons are considered the main functional GC targets (10). These include the medial pre-frontal cortex (mPFC), the striatum, the hippocampus and the amygdala. For example, stress impairs the long-term potentiation in projections from the basolateral amygdala to the pre-limbic PFC and to the ventral hippocampus-mPFC connection (11, 12). Acute GC treatment changes synaptic strength and excitability within hours, while repeated GC exposure or chronic stress paradigms consolidate such changes structurally via dendritic remodeling. This includes atrophy and reduction in apical spine density in the hippocampus region CA3 (13, 14) and the medial prefrontal pyramidal cells (15-17), but also increased dendritic complexity in *e.g.* the amygdala (18).

The cellular basis for most GC effects in the brain is largely unknown. Below we will argue that targeting specific processes downstream of GC receptors activation may be a good strategy to unravel these effects. Before we address this question, we will discuss general molecular mechanisms of corticosteroid receptor signaling as well as different types of ligands.

### *The receptors: sensitivity of brain regions*

The effects of GCs are mediated by the glucocorticoid receptor (GR) and the mineralocorticoid receptor (MR). These corticosteroid receptors are closely related members of the family of nuclear steroid receptors, which act as ligand-dependent transcription factors. They differ in tissue-distribution and ligand affinity (4). MR is an aldosterone receptor in tissues that convert GCs into inactive metabolites, but in relation to the stress system, it acts as a receptor for cortisol and corticosterone. The high GC affinity of the MR results in its function as a sensor of basal GC levels, and its involvement in shaping the initial response to stress (19). GR has a lower affinity (20) and is more ubiquitously expressed throughout the human body. The difference in affinity is about 10-fold, as is apparent from the pharmacological dissociation constants ( $K_d$ ) of  $\sim 0.5$  nM for MR and 5 nM for GR. GR therefore responds to elevated levels of GCs, *i.e.* during stress. Specifically in the brain, MR is mainly restricted to the hippocampus and other limbic areas, while GR is widely expressed. Within brain structures, different cell types show both qualitative and quantitative differences in their nuclear receptor expression profile. The use of single-cell profiling with RNA

sequencing (scRNA-seq) allows large-scale comprehensive molecular classification of cell types in the brain and its sub-regions. For example, one study identified 69 different neuronal cell sub-types in the human temporal cortex (21). These recent data suggest differential expression of GR in specific cell types in the human cortex, and also in hippocampal sub-regions (21-23). They also illustrate that scRNA-seq approaches can be used to identify specific cell types and intracellular pathways required for GC action in cognitive functions such as memory consolidation.

After development of specific receptor (ant)agonists and the discovery of the two corticosteroid receptor types in the brain, it became clear GCs are essential for memory formation and each receptor type plays a distinctive role. For example, GR blockade shortly after learning interferes with long-term memory consolidation, while MR blockade interfered with response strategy in novel situations (24). Later studies using GR dimerization-deficient mice confirmed these pharmacological experiments, and established the necessity of genomic GR action in memory consolidation (25). Likely, there is also a role for early, non-genomic effects of GCs in the promotion of memory formation ('encoding'). The formation and consolidation of spatial and declarative memory heavily rely on hippocampal physiology and synaptic plasticity, for instance via the modulation of glutamatergic transmission (26-28). More recent work has also established roles of GR and MR in other – non spatial - learning tasks; for example memory retrieval is impaired by prior exposure to GCs (29). The suppressive effects on memory retrieval may be turned into use for treatment of post-traumatic stress disorder (PTSD), anxiety and phobic disorders (30, 31).

Processes that depend on GR are recapitulated – or exaggerated – by the use of synthetic GR agonists (like dexamethasone and prednisone), which are the mainstay in the treatment of inflammatory diseases, autoimmune disorders and hematologic cancers (3, 32). The often maladaptive effects associated with excessive GC exposure include increased food intake and weight gain (33), disturbance of awakening/sleeping rhythm (8), anxiety, depression (34) and impaired cognitive functions (7).

Neuropsychiatric side effects after synthetic GC treatment may be the result of GR hyperactivation but there is also evidence that implicates MR hypoactivation. GR-specific agonists suppress endogenous cortisol production through GR-mediated negative feedback on the HPA axis, which results in diminished MR activity, that can be restored by exogenous cortisol treatment (35, 36). Nevertheless, GR antagonists are widely considered to counteract

the adverse consequences of endogenous GC exposure, and may be considered for clinical use in some brain diseases, for instance in depression (37, 38), alcohol addiction (39) and neurodegenerative disease (40).

Since the classical GR antagonist RU486 also binds to progesterone and androgen receptors (PR and AR) (41), efforts have been made to develop more selective GR antagonists. ORG34850 was shown to inhibit GR-mediated negative feedback on the HPA axis, and was therefore considered to be a potential treatment for mood disorders (as dysfunctional HPA axis is known to be involved in depression) (42). More recently the selective GR antagonist CORT113176 was shown to be efficacious in a model for alcohol seeking and self-administration in alcohol-dependent rats (39).

Thus, GR agonists and antagonists may help to understand which processes involve GR, and may be used to counteract maladaptive consequences of GR overactivation. However, they will activate or antagonize all GR-mediated effects, and given the pleiotropic nature of GR signaling, this may also trigger unwanted activities. There are additional types of ligands, called ‘dissociated ligands’ or Selective GR Modulators (SGRMs) that activate only a subset of GR-dependent signals in the cell (43, 44). These may not only be advantageous in clinical settings, but they are also very promising tools to identify the cellular mechanisms underlying the many different effects of GCs in the brain.

### **Glucocorticoid Receptor-mediated transcription**

Non-genomic effects notwithstanding (45), GR is best understood as ligand-dependent transcriptional factor (TF). The receptor binds directly to DNA via its central DNA binding domain to inverted-repeat Glucocorticoid Responsive Elements (GREs) or to half site DNA sequences, which mainly leads to upregulation of gene transcription (46). This mechanism is shared with the other steroid receptors. Direct DNA occupancy of GR can also lead to the repression of target genes via so-called negative GREs (nGREs) (47, 48). DNA binding often occurs in conjunction with other transcription factors, and is typically followed by recruitment of coregulator proteins that either stimulate or repress transcription (49). Additionally, GR can indirectly mediate transcription by inducing protein-protein interactions with other TFs already bound to DNA specific loci (50). One well-known example is the repressive cross-talk between GR and pro-inflammatory transcription factor AP-1 (51, 52).

### *Glucocorticoid receptor genomic occupancy in the hippocampus*

Chromatin immunoprecipitation followed by sequencing (ChIP-seq) can be used to identify GR-occupied genomic regions *in vivo*, which can differ between species, tissues, cell types and physiological state. In the rat hippocampus, GR signaling is mostly dependent on direct DNA-binding. In 2013, Polman et al. identified almost 2500 genomic GR binding sites (GBS) with ChIP-seq in rat hippocampus. Almost all these GBS contained a GRE (53). More recently, it has been confirmed that 89.9% of hippocampal GBS contained full or half GREs (54, 55). Because ChIP-seq has only been performed on whole hippocampus homogenate, it is possible that protein-protein interactions in subsets of activated neurons have been overlooked. Nevertheless, these results show that direct DNA binding is a major mechanism contributing to GR activity in the hippocampus. The context (e.g. cell-type) dependence of GRE-driven targets is apparent from the specific association of GR binding with other transcription factors, such as Nuclear Factor-1 (NF-1) (53-55). Likely, GR can also heterodimerize with MR at a subset of binding sites (55, 56).

#### *Chromatin remodeling*

GR transcriptional activity is regulated at different levels, starting with the organization of the nuclear chromatin and the availability of GREs for binding of the receptors. Gene transcription is orchestrated by nucleosome architecture and chromatin configuration. Both levels are further coordinated by genomic and epigenomic mechanisms that give dynamics to the chromosome layout and thereby modulate its accessibility. It has been shown that cell-specific GR-DNA interaction patterns are pre-determined by cell-specific differences in chromatin accessibility. Genome-wide DNase I analysis and ChIP-seq were used to assess respectively chromatin accessibility and GR binding at high resolution before and after treatment with the synthetic GC dexamethasone. This revealed that pre-existing accessibility of the chromatin largely but not completely determines genomic occupancy of GR (57). It is of interest to identify also at which loci the GR can induce chromatin opening by itself, rather than binding at pre-existing accessible regions.

#### *Steroid receptor-associated coregulatory complexes*

Epigenetic remodelers and chromatin context have a critical role in determining the transcriptional outcome, and therefore the directionality and intensity of gene expression changes. However, gene regulation also relies on the complex formation of GR with coregulatory partners (49). GR transcription complexes typically consist of approximately 10 different coregulators, either stably or dynamically associated with each other (49, 58). Tens



of transcriptional coregulators are known to interact with GR, resulting in a large variety of transcriptional complexes that lead to highly diverse gene expression outcomes (59). For example, the members of the Steroid Receptor Coactivator (SRC) family are transcriptional coregulators that are differentially expressed in the brain, particularly the hippocampus, the cortex and the hypothalamus (60). Knockout of SRC-1 is associated with disturbed regulation of important GR targets: *Pomc* in the pituitary (61) and *Crh*, both in the hypothalamus and amygdala (62, 63). It has been shown that the absence of SRC coding genes *NCoA2* and *NCoA3* (Nuclear Coactivator 2 and 3) had opposite effects on anxiety responses. Female *NCoA2* knockout mice demonstrated decreased anxiety-like behavior while *NCoA3* knockout increased it. The latter data suggest that loss in SRC function underlies changes in behavioral phenotypes, but it is still unclear which steroid receptor pathways are involved in these effects as the coregulators affect several steroid receptors (64). The coregulators may thus be viewed as integrators of multiple steroid signals. In a recent study, region-dependent expression of 62 coregulators and co-expression with all steroid receptors was described in the brain (65) (Fig. 1). It is clear that the co-expression of GR and MR with coregulators is dependent on the brain region. Region-dependent recruitment of coregulator proteins likely underlies the region-specific effects of steroid-receptor mediated transcription.

The substantial number of distinct GR signaling pathways and the need for specific manipulation is the basis for the category of SGRMs (43, 44, 66). Historically, dissociated ligands bind GR and have higher efficacy at transrepressive protein-protein interactions than at transcription via GREs (47, 48). These types of ligands have been pursued to separate anti-inflammatory effects from unwanted metabolic side effects, but it has turned out that anti-inflammatory effects also involve GRE-dependent transcription. For example, GR activation can lead to the upregulation of I $\kappa$ B- $\alpha$  (NF- $\kappa$ B inhibitor alpha), which limits the pro-inflammatory actions of NF- $\kappa$ B (51). In addition, recent data suggest that inhibition of NF- $\kappa$ B driven proinflammatory transcription may depend on GR binding to nGREs (67). The term ‘selective modulators’ relates to ligands that stimulate interactions with only a subset of the GR coregulators that are recruited in the presence of full agonists (59, 68). Based on their selective efficacy, this class of drugs has the potential to combine agonistic and antagonistic properties in GR-mediated transcription. This may allow dissection of beneficial from adverse effects, and thus holds potential to improve current GC-based therapies. We recently discovered that the actual combination of agonism and antagonism is sometimes required to generate beneficial effects on disease outcome. To date, the best example concerns a liver

steatosis disease model, in which the SGRM CORT188335 mimicked GR agonism by stimulating lipid efflux via very low-density lipoprotein (VLDL) production, whereas it lacked agonist efficacy in stimulating fatty acids uptake by the liver. In this way the hepatic lipid flux was affected in such a way that efflux dominated over influx, and liver steatosis could be attenuated (69).

Although a substantial number of whole genome transcriptional and ChIP-seq datasets have been generated, it remains a major challenge to couple the extensive transcriptional outcome of GR activation to effects at the level of synaptic signaling and behavior. Comparing the effects of SGRMs on behavior, coregulator interaction and the transcriptional signature, may help to unravel the target genes and signaling pathways underlying particular GR effects in the brain and beyond. Below, we illustrate this approach based on experiments with two recently developed SGRMs, CORT108297 and CORT118335.

#### *CORT108297 and CORT118335 in memory, behavior and neurodegenerative diseases*

In an attempt to understand the GC effects on memory consolidation, SGRMs CORT108297 and CORT118335 were studied in animal models. CORT108297 is a high-affinity GR ligand ( $K_d = 0.9$  nM) (70), while CORT118335 has a lower affinity for GR ( $K_d$  of ~8 nM), and shows some affinity for the MR, for which it acts as an antagonist (71). CORT108297 was shown to have GR agonistic effects in an inhibitory avoidance memory task (Fig. 2A) (72), a paradigm known to be potentiated by GR (73) and usually set up to assess memory strength (74). CORT118335 had opposite effects on memory consolidation as CORT118335 injection an hour before the avoidance memory task antagonized the memory-enhancing effect of corticosterone, similarly to the classical GR antagonist RU486 (Fig. 2B) (59).

In a separate study, CORT108297 was shown to decrease immobility in a forced-swim stress paradigm, which was interpreted as GR antagonist effects on depression-like behavior (75). CORT108297 also displayed antagonist-like effects on corticosterone-induced reduction of neuronal differentiation (72), analogous to the effects of the full GR antagonist RU486 (76). In terms of gene expression, CORT108297 was shown to act both as agonist and as antagonist, depending on the target gene (72). A transcriptome analysis in the liver showed that in this tissue CORT118335 acts as a partial agonist on most GR target genes, but lacks agonism at a – functionally important – subset of targets (69). Both compounds are thus selective modulators, rather than classical agonists or antagonists.

CORT108297 has also been studied in models of neurodegeneration. There is a substantial association between HPA axis dysfunction and Alzheimer's disease (AD), as AD patients show elevated basal cortisol levels (77, 78). The GR antagonist RU486 has beneficial effects in many models of AD (40, 79, 80). In a rat model for AD, deregulation of the HPA axis is associated with cognitive impairments, apoptotic and neuroinflammatory processes, and an induction of amyloidogenic pathway. In this model, CORT108297 treatment restored synaptic markers in the hippocampus and cognitive function in spatial short-term memory (81). CORT108297 also restored hippocampal integrity and normalized neurogenesis in the dentate gyrus in mutant Wobbler mice as a model for human amyotrophic lateral sclerosis (82). These mice also show motoneuron degeneration, motor deficits, astrogliosis and microgliosis in the spinal cord, which are correlated with increased levels of corticosterone in plasma, brain and spinal cord. In these two neurodegenerative pathologic models it is likely that the antagonistic properties of selective modulators are mainly responsible for the beneficial effects. Indeed, the full GR selective antagonist CORT113176 also rescued the phenotype of mutant Wobbler mice (83).

#### *Differential glucocorticoid receptor coregulator recruitment and gene expression*

The differential agonistic and antagonistic effects of CORT118335 and CORT108297 on various processes likely relate to their differential effects on gene expression that in turn depend on distinct coregulator recruitment by the ligand-bound GR. Differential GR-coregulator interaction profiles for CORT118335 and CORT108297 were demonstrated via the Microarray Assay for Real-time Coregulator-Nuclear receptor Interaction (MARCoNI) technology (58), which measures *in vitro* interactions between the GR ligand binding domain (LBD) and peptides containing the coregulator domains that are responsible for interactions with the GR (LxxLL motif containing Nuclear Receptor (NR)-boxes) (84, 85). The GR interaction profiles in the presence of SGRMs can in this way be compared with those of full GR agonists (cortisol, dexamethasone) and antagonists (mifepristone) (72).

As an example, SRC-1 is a GR coregulator associated with HPA axis function and the regulation of specific GR target genes (62). The involvement of SRC-1 in negative feedback regulation of the HPA axis is complex due to the fact that there are two splice variants, SRC-1A and SRC-1E. They share three NR boxes, but SRC-1A has an additional NR-box in the C-terminal part of the protein (86). Moreover, SRC-1A is highly expressed in the pituitary and the hypothalamus, whereas both factors seem to be equally expressed in many other brain

regions. This implies that there is a ‘targetable’ GR-NR-box interaction that is specific for the hypothalamus and the pituitary (61).

CORT108297 differentiates GR interactions with the two SRC-1 splice variants as it preferentially induces an interaction between GR-LBD and the SRC-1A NR-box 4 in the MARCoNI assay. The full agonist dexamethasone does not show any preference for SRC-1 NR-boxes (72). These observations suggest that CORT108297 selective effects on gene expression could be based on its potential to induce specific interactions between GR and the SRC-1 coregulator NR-box 4. This notion seems to hold, as CORT108297 could differentiate between regulation of *Crh* transcription in the hypothalamus (agonism) and the central nucleus of the amygdala (no agonism) (72).

The GR coregulator interactions that are induced upon binding of CORT108297 and CORT118335 are both intermediate between those seen as full agonists (dexamethasone) and full antagonists (RU486). These SGRM-induced coregulator interaction profiles show partial overlap, but also clear differences (59, 72) (Fig. 3). Specifically, among 155 NR-boxes, 40 were shared between CORT118335, CORT108297 and dexamethasone, which likely represent shared agonistic properties. For a number of motifs, CORT118335 displayed agonist-like or antagonist-like GR interactions that were not present for CORT108297 (59).

It is an attractive hypothesis to relate functional differences to the coregulator interactions. The number of differential interactions of GR bound CORT108297 and CORT118335 provides a short-list of responsible signaling pathways. Likewise, motifs that differ between CORT108297, CORT118335, RU486 on one hand and full agonists on the other, may point to coregulators that underlie the GR-mediated aggravation of neurodegenerative processes and cognitive impairments.

Those coregulators that are differentially recruited after CORT108297 or CORT118335 binding to the GR may explain the functional differences between the compounds (59). Comparisons between SGRMs in terms of coregulator interactions with GR and the resulting functional effects may allow the linking of particular signaling pathways with more integrative consequences, especially in the brain where several cognitive and behavioral functions are regulated by GCs.

## Conclusion and perspectives

In conclusion, GR-mediated transcription depends on several parameters including chromatin accessibility, DNA-binding configuration, interaction with other transcription factors and GR coregulator interactome. These parameters are highly context-dependent and differ according to the tissue, the cell type, the physiological state and GR ligand. Ligand-related changes that are reflected in behavior – particularly in memory consolidation – could rely on variations in GR coregulator interactome in the hippocampus and the other limbic structures involved in memory. The combination of behavioral and transcriptional effects of SGRMs – with knowledge about their induced GR coregulatory interactome, and the cell specific co-expression of potentially interacting partners (65) – represents an interesting new research strategy to identify molecular pathways that are responsible for adaptive and maladaptive effects of GCs on brain function (Fig. 4).

Future work may include validation of these putative interactions by using co-immunoprecipitation (co-IP) and ChIP-seq directed towards the identified coregulators and GR, either in mixed cell populations or at the single-cell level in order to characterize the cells functionally involved in the observed changes in learning and memory. It is important to consider that the integration of the simultaneous levels of modulation represent a substantial challenge. The coregulators only represent one level of GR transcriptional modulation, and it would be of interest to also investigate the nucleosome configuration, chromatin accessibility, other transcription factors or post-translational modifications of the interacting effectors.

Beyond their fundamental input, SGRMs also hold potential therapeutic value in GC-related disorders of the nervous system and beyond. Some authors have suggested superior effects of selective modulators over pure antagonists (81). GR selectivity over other steroid receptors is the first asset of SGRMs, as it prevents side effects related to AR, PR or even MR activities (although CORT118335 does act as a low affinity MR antagonist). Furthermore, the other major advantage of SGRMs is the specificity regarding their agonistic and antagonistic properties according to the cell type or transcriptional target, which provides targeting of only a subset of processes. For neurodegenerative diseases, it appears that residual GC anti-inflammatory efficacy combined with antagonism on classical neuro-endangerment may represent the ideal SGRM properties (83). The application of the GR coregulator interactome hypothesis in these models will also allow the dissection of GR-mediated effects and the potential benefits of selective GR modulation compared to GR antagonism or agonism.

The ultimate goal for this line of research is to identify the GR-mediated transcriptional effects that are responsible for adaptive processes and for brain disorders or pathologies, and to evaluate therapeutic targeting of the latter pathways.

## Statements

### Disclosure Statement

This research was partly funded by Corcept Therapeutics who develop new GR ligands. HH is an employee of Corcept Therapeutics. RH is an employee of Pamgene BV.

### Funding Sources

NWO/ALW (Netherlands Organization for Scientific Research), NWO-ZonMw, Board of Directors of the LUMC.

### Author Contributions

Conception and design: OM, JK, JB, EV.

Study supervision: OM, JK.

Acquisition of data: RH, AM, JB, EV.

Writing of the manuscript: OM, JK, EV.

Review of the manuscript: OM, JK, JB, HH, LvW, LK, AM.

## References

1. Smith SM, Vale WW. The role of the hypothalamic-pituitary-adrenal axis in neuroendocrine responses to stress. *Dialogues in clinical neuroscience*. 2006;8(4):383-95.
2. Lightman SL, Conway-Campbell BL. The crucial role of pulsatile activity of the HPA axis for continuous dynamic equilibration. *Nat Rev Neurosci*. 2010;11(10):710-8.
3. Oakley RH, Cidlowski JA. The biology of the glucocorticoid receptor: new signaling mechanisms in health and disease. *J Allergy Clin Immunol*. 2013;132(5):1033-44.
4. Joels M. Corticosteroids and the brain. *J Endocrinol*. 2018;238(3):R121-R30.
5. Munck A, Guyre PM, Holbrook NJ. Physiological Functions of Glucocorticoids in Stress and Their Relation to Pharmacological Actions\*. *Endocrine Reviews*. 1984;5(1):25-44.
6. Sapolsky RM. Stress hormones: good and bad. *Neurobiol Dis*. 2000;7(5):540-2.
7. Marques AH, Silverman MN, Sternberg EM. Glucocorticoid dysregulations and their clinical correlates. From receptors to therapeutics. *Ann N Y Acad Sci*. 2009;1179:1-18.
8. Fietta P, Fietta P, Delsante G. Central nervous system effects of natural and synthetic glucocorticoids. *Psychiatry Clin Neurosci*. 2009;63(5):613-22.
9. Liston C, Gan WB. Glucocorticoids are critical regulators of dendritic spine development and plasticity in vivo. *Proc Natl Acad Sci U S A*. 2011;108(38):16074-9.

10. Myers B, McKlveen JM, Herman JP. Glucocorticoid actions on synapses, circuits, and behavior: implications for the energetics of stress. *Front Neuroendocrinol.* 2014;35(2):180-96.
11. Maroun M, Richter-Levin G. Exposure to Acute Stress Blocks the Induction of Long-Term Potentiation of the Amygdala–Prefrontal Cortex Pathway & In Vivo. *The Journal of Neuroscience.* 2003;23(11):4406.
12. Richter-Levin G, Maroun M. Stress and amygdala suppression of metaplasticity in the medial prefrontal cortex. *Cereb Cortex.* 2010;20(10):2433-41.
13. Magariños AMa, McEwen BS, Flügge G, Fuchs E. Chronic Psychosocial Stress Causes Apical Dendritic Atrophy of Hippocampal CA3 Pyramidal Neurons in Subordinate Tree Shrews. *The Journal of Neuroscience.* 1996;16(10):3534.
14. Watanabe Y, Gould E, McEwen BS. Stress induces atrophy of apical dendrites of hippocampal CA3 pyramidal neurons. *Brain Research.* 1992;588(2):341-5.
15. Radley JJ, Rocher AB, Miller M, Janssen WG, Liston C, Hof PR, et al. Repeated stress induces dendritic spine loss in the rat medial prefrontal cortex. *Cereb Cortex.* 2006;16(3):313-20.
16. Radley JJ, Sisti HM, Hao J, Rocher AB, McCall T, Hof PR, et al. Chronic behavioral stress induces apical dendritic reorganization in pyramidal neurons of the medial prefrontal cortex. *Neuroscience.* 2004;125(1):1-6.
17. Wellman CL. Dendritic reorganization in pyramidal neurons in medial prefrontal cortex after chronic corticosterone administration. *Journal of Neurobiology.* 2001;49(3):245-53.
18. Mitra R, Sapolsky RM. Acute corticosterone treatment is sufficient to induce anxiety and amygdaloid dendritic hypertrophy. *Proceedings of the National Academy of Sciences.* 2008;105(14):5573.
19. de Kloet ER, Joels M, Holsboer F. Stress and the brain: from adaptation to disease. *Nat Rev Neurosci.* 2005;6(6):463-75.
20. Reul JMHM, Kloet ERD. Two Receptor Systems for Corticosterone in Rat Brain: Microdistribution and Differential Occupation. *Endocrinology.* 1985;117(6):2505-11.
21. Hodge RD, Bakken TE, Miller JA, Smith KA, Barkan ER, Graybuck LT, et al. Conserved cell types with divergent features between human and mouse cortex. *bioRxiv.* 2018.
22. Cembrowski MS, Wang L, Sugino K, Shields BC, Spruston N. Hipposeq: a comprehensive RNA-seq database of gene expression in hippocampal principal neurons. *Elife.* 2016;5:e14997.
23. Saunders A, Macosko E, Wysoker A, Goldman M, Krienen F, Bien E, et al. A Single-Cell Atlas of Cell Types, States, and Other Transcriptional Patterns from Nine Regions of the Adult Mouse Brain. *bioRxiv.* 2018.
24. Oitzl MS, de Kloet ER. Selective corticosteroid antagonists modulate specific aspects of spatial orientation learning. *Behavioral Neuroscience.* 1992;106(1):62-71.
25. Oitzl MS, Reichardt HM, Joels M, de Kloet ER. Point mutation in the mouse glucocorticoid receptor preventing DNA binding impairs spatial memory. *Proceedings of the National Academy of Sciences.* 2001;98(22):12790.
26. Sarabdjitsingh RA, Jezequel J, Pasricha N, Mikasova L, Kerkhofs A, Karst H, et al. Ultradian corticosterone pulses balance glutamatergic transmission and synaptic plasticity. *Proc Natl Acad Sci U S A.* 2014;111(39):14265-70.
27. Karst H, Joels M. Corticosterone slowly enhances miniature excitatory postsynaptic current amplitude in mice CA1 hippocampal cells. *J Neurophysiol.* 2005;94(5):3479-86.
28. Joels M, Sarabdjitsingh RA, Karst H. Unraveling the time domains of corticosteroid hormone influences on brain activity: rapid, slow, and chronic modes. *Pharmacol Rev.* 2012;64(4):901-38.
29. Roozendaal B, Griffith QK, Buranday J, de Quervain DJF, McGaugh JL. The hippocampus mediates glucocorticoid-induced impairment of spatial memory retrieval: Dependence on the basolateral amygdala. *Proceedings of the National Academy of Sciences.* 2003;100(3):1328.
30. Ouanes S. Glucocorticoid-based therapeutic options for PTSD. *European Psychiatry.* 2016;33.
31. de Quervain DJ, Bentz D, Michael T, Bolt OC, Wiederhold BK, Margraf J, et al. Glucocorticoids enhance extinction-based psychotherapy. *Proc Natl Acad Sci U S A.* 2011;108(16):6621-5.

32. Pufall MA. Glucocorticoids and Cancer. *Adv Exp Med Biol.* 2015;872:315-33.
33. Solano JM, Jacobson L. Glucocorticoids reverse leptin effects on food intake and body fat in mice without increasing NPY mRNA. *American Journal of Physiology-Endocrinology and Metabolism.* 1999;277(4):E708-E16.
34. Joels M. Impact of glucocorticoids on brain function: relevance for mood disorders. *Psychoneuroendocrinology.* 2011;36(3):406-14.
35. Meijer OC, de Kloet ER. A Refill for the Brain Mineralocorticoid Receptor: The Benefit of Cortisol Add-On to Dexamethasone Therapy. *Endocrinology.* 2017;158(3):448-54.
36. Warris LT, van den Heuvel-Eibrink MM, Ariës IM, Pieters R, van den Akker ELT, den Boer ML. Hydrocortisone does not influence glucocorticoid sensitivity of acute lymphoblastic leukemia cells. *Haematologica.* 2015;100(4):e137-e9.
37. Belanoff JK, Rothschild AJ, Cassidy F, DeBattista C, Baulieu E-E, Schold C, et al. An open label trial of C-1073 (mifepristone) for psychotic major depression\*. *Biological Psychiatry.* 2002;52(5):386-92.
38. Gallagher P, Young AH. Mifepristone (RU-486) treatment for depression and psychosis: a review of the therapeutic implications. *Neuropsychiatric disease and treatment.* 2006;2(1):33-42.
39. Vendruscolo LF, Estey D, Goodell V, Macshane LG, Logrip ML, Schlosburg JE, et al. Glucocorticoid receptor antagonism decreases alcohol seeking in alcohol-dependent individuals. *J Clin Invest.* 2015;125(8):3193-7.
40. Green KN, Billings LM, Roozendaal B, McGaugh JL, LaFerla FM. Glucocorticoids increase amyloid-beta and tau pathology in a mouse model of Alzheimer's disease. *J Neurosci.* 2006;26(35):9047-56.
41. Spiga F, Knight DM, Droste SK, Conway-Campbell B, Kershaw Y, MacSweeney CP, et al. Differential effect of glucocorticoid receptor antagonists on glucocorticoid receptor nuclear translocation and DNA binding. *Journal of psychopharmacology (Oxford, England).* 2011;25(2):211-21.
42. Spiga F, Harrison LR, Wood SA, Atkinson HC, MacSweeney CP, Thomson F, et al. Effect of the glucocorticoid receptor antagonist Org 34850 on basal and stress-induced corticosterone secretion. *J Neuroendocrinol.* 2007;19(11):891-900.
43. De Bosscher K, Berghe WV, Beck IME, Van Molle W, Hennuyer N, Hapgood J, et al. A fully dissociated compound of plant origin for inflammatory gene repression. *Proceedings of the National Academy of Sciences of the United States of America.* 2005;102(44):15827.
44. Meijer OC, Koorneef LL, Kroon J. Glucocorticoid receptor modulators. *Ann Endocrinol (Paris).* 2018;79(3):107-11.
45. Femke LG, Henk K, Kloet ERd, Marian J. Rapid non-genomic effects of corticosteroids and their role in the central stress response. *Journal of Endocrinology.* 2011;209(2):153-67.
46. Weikum ER, Knuesel MT, Ortlund EA, Yamamoto KR. Glucocorticoid receptor control of transcription: precision and plasticity via allostery. *Nat Rev Mol Cell Biol.* 2017;18(3):159-74.
47. Surjit M, Ganti KP, Mukherji A, Ye T, Hua G, Metzger D, et al. Widespread negative response elements mediate direct repression by agonist-liganded glucocorticoid receptor. *Cell.* 2011;145(2):224-41.
48. Hudson WH, Youn C, Ortlund EA. The structural basis of direct glucocorticoid-mediated transrepression. *Nature Structural & Molecular Biology.* 2012;20:53.
49. O'Malley BW, Malovannaya A, Qin J. Minireview: nuclear receptor and coregulator proteomics--2012 and beyond. *Mol Endocrinol.* 2012;26(10):1646-50.
50. Martens C, Bilodeau S, Maira M, Gauthier Y, Drouin J. Protein-protein interactions and transcriptional antagonism between the subfamily of NGFI-B/Nur77 orphan nuclear receptors and glucocorticoid receptor. *Mol Endocrinol.* 2005;19(4):885-97.
51. De Bosscher K, Van Craenenbroeck K, Meijer OC, Haegeman G. Selective transrepression versus transactivation mechanisms by glucocorticoid receptor modulators in stress and immune systems. *Eur J Pharmacol.* 2008;583(2-3):290-302.



465 52. Heck S, Kullmann M, Gast A, Ponta H, Rahmsdorf HJ, Herrlich P, et al. A distinct modulating  
466 domain in glucocorticoid receptor monomers in the repression of activity of the transcription factor  
467 AP-1. *The EMBO journal*. 1994;13(17):4087-95.

468 53. Polman JA, de Kloet ER, Datson NA. Two populations of glucocorticoid receptor-binding sites  
469 in the male rat hippocampal genome. *Endocrinology*. 2013;154(5):1832-44.

470 54. Pooley JR, Flynn BP, Grontved L, Baek S, Guertin MJ, Kershaw YM, et al. Genome-Wide  
471 Identification of Basic Helix-Loop-Helix and NF-1 Motifs Underlying GR Binding Sites in Male Rat  
472 Hippocampus. *Endocrinology*. 2017;158(5):1486-501.

473 55. van Weert L, Buurstede JC, Mahfouz A, Braakhuis PSM, Polman JAE, Sips HCM, et al. NeuroD  
474 Factors Discriminate Mineralocorticoid From Glucocorticoid Receptor DNA Binding in the Male Rat  
475 Brain. *Endocrinology*. 2017;158(5):1511-22.

476 56. Reul JM, H. M. Mineralocorticoid and glucocorticoid receptor-mediated control of genomic  
477 responses to stress in the brain AU - Mifsud, Karen R. *Stress*. 2018;21(5):389-402.

478 57. John S, Sabo PJ, Thurman RE, Sung MH, Biddie SC, Johnson TA, et al. Chromatin accessibility  
479 pre-determines glucocorticoid receptor binding patterns. *Nat Genet*. 2011;43(3):264-8.

480 58. Desmet SJ, Dejager L, Clarisse D, Thommis J, Melchers D, Bastiaensen N, et al. Cofactor  
481 profiling of the glucocorticoid receptor from a cellular environment. *Methods Mol Biol*.  
482 2014;1204:83-94.

483 59. Atucha E, Zalachoras I, van den Heuvel JK, van Weert LT, Melchers D, Mol IM, et al. A Mixed  
484 Glucocorticoid/Mineralocorticoid Selective Modulator With Dominant Antagonism in the Male Rat  
485 Brain. *Endocrinology*. 2015;156(11):4105-14.

486 60. Meijer OC, Steenbergen PJ, de Kloet ER. Differential Expression and Regional Distribution of  
487 Steroid Receptor Coactivators SRC-1 and SRC-2 in Brain and Pituitary\*. *Endocrinology*.  
488 2000;141(6):2192-9.

489 61. Winnay JN, Xu J, O'Malley BW, Hammer GD. Steroid receptor coactivator-1-deficient mice  
490 exhibit altered hypothalamic-pituitary-adrenal axis function. *Endocrinology*. 2006;147(3):1322-32.

491 62. Lachize S, Apostolakis EM, van der Laan S, Tijssen AMI, Xu J, de Kloet ER, et al. Steroid  
492 receptor coactivator-1 is necessary for regulation of corticotropin-releasing hormone by chronic  
493 stress and glucocorticoids. *Proceedings of the National Academy of Sciences*. 2009;106(19):8038.

494 63. Zalachoras I, Verhoeve SL, Toonen LJ, van Weert LTCM, van Vlodrop AM, Mol IM, et al.  
495 Isoform switching of steroid receptor co-activator-1 attenuates glucocorticoid-induced anxiogenic  
496 amygdala CRH expression. *Molecular Psychiatry*. 2016;21:1733.

497 64. Stashi E, Wang L, Mani SK, York B, O'Malley BW. Research resource: loss of the steroid  
498 receptor coactivators confers neurobehavioral consequences. *Mol Endocrinol*. 2013;27(10):1776-87.

499 65. Mahfouz A, Lelieveldt BP, Grefhorst A, van Weert LT, Mol IM, Sips HC, et al. Genome-wide  
500 coexpression of steroid receptors in the mouse brain: Identifying signaling pathways and functionally  
501 coordinated regions. *Proc Natl Acad Sci U S A*. 2016;113(10):2738-43.

502 66. Heier CR, Damsker JM, Yu Q, Dillingham BC, Huynh T, Van der Meulen JH, et al. VBP15, a  
503 novel anti-inflammatory and membrane-stabilizer, improves muscular dystrophy without side  
504 effects. *EMBO Mol Med*. 2013;5(10):1569-85.

505 67. Hudson WH, Vera IMSd, Nwachukwu JC, Weikum ER, Herbst AG, Yang Q, et al. Cryptic  
506 glucocorticoid receptor-binding sites pervade genomic NF- $\kappa$ B response elements. *Nature*  
507 *Communications*. 2018;9(1):1337.

508 68. Coghlan MJ, Jacobson PB, Lane B, Nakane M, Lin CW, Elmore SW, et al. A novel  
509 antiinflammatory maintains glucocorticoid efficacy with reduced side effects. *Mol Endocrinol*.  
510 2003;17(5):860-9.

511 69. Koorneef LL, van den Heuvel JK, Kroon J, Boon MR, t Hoen PAC, Hettne KM, et al. Selective  
512 glucocorticoid receptor modulation prevents and reverses non-alcoholic fatty liver disease in male  
513 mice. *Endocrinology*. 2018.

70. Clark RD, Ray NC, Williams K, Blaney P, Ward S, Crackett PH, et al. 1H-Pyrazolo[3,4-g]hexahydro-isoquinolines as selective glucocorticoid receptor antagonists with high functional activity. *Bioorg Med Chem Lett*. 2008;18(4):1312-7.
71. Hunt HJ, Ray NC, Hynd G, Sutton J, Sajad M, O'Connor E, et al. Discovery of a novel non-steroidal GR antagonist with in vivo efficacy in the olanzapine-induced weight gain model in the rat. *Bioorg Med Chem Lett*. 2012;22(24):7376-80.
72. Zalachoras I, Houtman R, Atucha E, Devos R, Tijssen AM, Hu P, et al. Differential targeting of brain stress circuits with a selective glucocorticoid receptor modulator. *Proc Natl Acad Sci U S A*. 2013;110(19):7910-5.
73. Fornari RV, Wichmann R, Atucha E, Desprez T, Eggens-Meijer E, Roozendaal B. Involvement of the insular cortex in regulating glucocorticoid effects on memory consolidation of inhibitory avoidance training. *Front Behav Neurosci*. 2012;6:10.
74. Atucha E, Roozendaal B. The inhibitory avoidance discrimination task to investigate accuracy of memory. *Front Behav Neurosci*. 2015;9:60.
75. Solomon MB, Wulsin AC, Rice T, Wick D, Myers B, McKlveen J, et al. The selective glucocorticoid receptor antagonist CORT 108297 decreases neuroendocrine stress responses and immobility in the forced swim test. *Horm Behav*. 2014;65(4):363-71.
76. Mayer JL, Klumpers L, Maslam S, de Kloet ER, Joels M, Lucassen PJ. Brief treatment with the glucocorticoid receptor antagonist mifepristone normalises the corticosterone-induced reduction of adult hippocampal neurogenesis. *J Neuroendocrinol*. 2006;18(8):629-31.
77. Hoogendijk WJG, Meynen G, Endert E, Hofman MA, Swaab DF. Increased cerebrospinal fluid cortisol level in Alzheimer's disease is not related to depression. *Neurobiology of Aging*. 2006;27(5):780.e1-e2.
78. Csernansky JG, Dong H, Fagan AM, Wang L, Xiong C, Holtzman DM, et al. Plasma cortisol and progression of dementia in subjects with Alzheimer-type dementia. *The American journal of psychiatry*. 2006;163(12):2164-9.
79. Baglietto-Vargas D, Medeiros R, Martinez-Coria H, LaFerla FM, Green KN. Mifepristone alters amyloid precursor protein processing to preclude amyloid beta and also reduces tau pathology. *Biological psychiatry*. 2013;74(5):357-66.
80. Lesuis SL, Weggen S, Baches S, Lucassen PJ, Krugers HJ. Targeting glucocorticoid receptors prevents the effects of early life stress on amyloid pathology and cognitive performance in APP/PS1 mice. *Translational psychiatry*. 2018;8(1):53-.
81. Pineau F, Canet G, Desrumaux C, Hunt H, Chevallier N, Ollivier M, et al. New selective glucocorticoid receptor modulators reverse amyloid-beta peptide-induced hippocampus toxicity. *Neurobiol Aging*. 2016;45:109-22.
82. Meyer M, Gonzalez Deniselle MC, Hunt H, de Kloet ER, De Nicola AF. The selective glucocorticoid receptor modulator CORT108297 restores faulty hippocampal parameters in Wobbler and corticosterone-treated mice. *J Steroid Biochem Mol Biol*. 2014;143:40-8.
83. Meyer M, Lara A, Hunt H, Belanoff J, de Kloet ER, Gonzalez Deniselle MC, et al. The Selective Glucocorticoid Receptor Modulator Cort 113176 Reduces Neurodegeneration and Neuroinflammation in Wobbler Mice Spinal Cord. *Neuroscience*. 2018;384:384-96.
84. Broekema MF, Hollman DAA, Koppen A, van den Ham HJ, Melchers D, Pijnenburg D, et al. Profiling of 3696 Nuclear Receptor-Coregulator Interactions: A Resource for Biological and Clinical Discovery. *Endocrinology*. 2018;159(6):2397-407.
85. Koppen A, Houtman R, Pijnenburg D, Jenning EH, Ruijtenbeek R, Kalkhoven E. Nuclear receptor-coregulator interaction profiling identifies TRIP3 as a novel peroxisome proliferator-activated receptor gamma cofactor. *Mol Cell Proteomics*. 2009;8(10):2212-26.
86. Kalkhoven E, Valentine JE, Heery DM, Parker MG. Isoforms of steroid receptor co-activator 1 differ in their ability to potentiate transcription by the oestrogen receptor. 1998;17(1):232-43.


## Figure Legends


**Fig. 1 Expression of nuclear receptor-associated coregulators in 12 regions of the mouse brain.** The values correspond to the log<sub>2</sub>-transformed ratio of the average expression in each region normalized to the average of expression in the whole brain. Adapted from Mahfouz et al., Proceedings of the National Academy of Sciences (PNAS), 2016.

**Fig. 2. SGRMs CORT108297 and CORT118335 acts as respectively a GR agonist and a GR antagonist in memory consolidation.** A) Acute post-training treatment with CORT108297 (20 mg/kg) or corticosterone (1 mg/kg) led to high retention latencies in an inhibitory avoidance task. Significance: \*\*\* p-value < 0.001. Adapted from Atucha et al., Endocrinology, 2015. B) The administration of CORT118335 (80 mg/kg) an hour before training prevents corticosterone-enhancement of memory consolidation in an inhibitory avoidance task. Significance: \*\*\* p-value < 0.001. Adapted from Zalachoras et al., Proceedings of the National Academy of Sciences (PNAS), 2013.

**Fig. 3 CORT108297 and CORT118335 induce GR binding to coregulators in an intermediate fashion compared to Dexamethasone and RU486.** Each column represents a unique coregulator-derived peptide from a range of 50 nuclear receptor coregulators. In the MARCoNI analysis, the peptides were immobilized on a solid support and incubated with cell lysates containing tagged-GR, a ligand (Dexamethasone, CORT108297, CORT118335 or RU486), and a tag-specific antibody coupled with a fluorophore. The relative interaction between each peptide and the GR was assessed by detecting the fluorescent label. The modulation index represents the log<sub>10</sub>-transformed ratio of the normalized fluorescence value. The values were normalized to the values obtained in control conditions (with Dimethyl Sulfoxide treatment) (unpublished data).

**Fig. 4 SGRMs lead to differential recruitment of GR regulatory elements in the brain, as schematically depicted for CORT108297 and CORT118335.** A) GR agonistic-like interactions promoted in CORT108297 context may underlie beneficial effects in memory consolidation. B) GR antagonistic-like interactions promoted in CORT118335 context may be responsible for the antagonism of GR-mediated memory consolidation.





Dexamethasone

CORT108297

CORT118335

RU486

Accepted manuscript

0

0.5

1.0


1.5

2.0

Modulation Index

A

CORT108297

**BEHAVIOR**

Enhances memory consolidation

**HIPPOCAMPAL GENE EXPRESSION**

Combined agonism and antagonism


**PATHOLOGY**

Similar to Mifepristone. Beneficial effects in:

- Alzheimer's disease (AD) model
- Amyotrophic lateral sclerosis (ALS) model
- Chronic stress

B

CORT118335

**BEHAVIOR**

Blocks memory consolidation

**HIPPOCAMPAL GENE EXPRESSION**

Predominantly antagonism

**PATHOLOGY**

To be investigated