

HW 6

Problem 1. Suppose A is an $n \times n$ matrix and $D = \text{Diag}(d)$ is a diagonal matrix made from the $n \times 1$ vector d . What does left-multiplication by D do to A ? (Think elementary operations.) What does right-multiplication do?

Consider the matrix

$$\begin{bmatrix} 3 & 4 & 0 \\ 1 & -1 & 1 \\ 0 & 9 & 2 \end{bmatrix}$$

and let $D = \text{Diag}(a, b, c)$. Compute $\tilde{A} := D^{-1}AD$ and then solve for a, b, c so that \tilde{A} becomes a symmetric matrix.

Problem 2. Consider the tridiagonal s.p.d:

$$A = \begin{bmatrix} a_1 & b_1 & & & & \\ b_1 & a_2 & b_2 & & & \\ & b_2 & a_3 & b_3 & & \\ & & \ddots & \ddots & \ddots & \\ & & & b_{n-2} & a_{n-1} & b_{n-1} \\ & & & & b_{n-1} & a_n \end{bmatrix}$$

Then A has the Cholesky factorization $A = LL^T$ for some matrix

$$L = \begin{bmatrix} x_1 & & & & & \\ y_1 & x_2 & & & & \\ & y_2 & x_3 & & & \\ & & \ddots & \ddots & & \\ & & & y_{n-1} & x_n \end{bmatrix}$$

Find an algorithm for computing L . First, formally compute the first few rows and columns of LL^T in terms of the unknowns x_i and y_j . Then, match them to the entries of A and solve repeatedly. You should find that the algorithm requires n square-root operations and $3n - 3$ other arithmetic operations.

Problem 3. Draw an $n \times n$ square grid in the plane, meaning that each side has n nodes, and so that $(0, 0)$ is the bottom left corner and all the edges have length 1.

Start by assuming $n = 3$.

- (a) Enumerate the nodes using the lexicographic order given by the coordinates of the nodes and write down the Laplacian matrix L .
- (b) Assume that the following Dirichlet boundary values are given:

$$\begin{aligned} g_{\text{left}} &= 0 \\ g_{\text{bottom}} &= 0 \\ g_{\text{right}} &= 2 \\ g_{\text{top}} &= 3 \end{aligned}$$

Solve the Dirichlet problem

$$\begin{cases} Lu = 0 & \text{on the interior nodes} \\ u = g & \text{on the boundary nodes} \end{cases}$$

How many equations and how many unknowns are there? Does it matter that the definition of the boundary data at the corners is ambiguous?

Now repeat (a) and (b) with $n = 4$.