



Theoretical Study of a Cylindrical Graphene-based Localized Surface Plasmon Spaser

Leila Prélat¹ (leilaprelat@gmail.com), Mauro Cuevas²,
Nicolás Passarelli³, Raúl Bustos Marún⁴, Ricardo Depine¹

[1] Departamento de Física, UBA and IFIBA, CONICET

[2] CONICET and Facultad de Ingeniería, UA

[3] Instituto de Ciencias de Materiales de Barcelona, CSIC

[4] IFEG, CONICET and Facultad de Ciencias Químicas, UNC

September 17, 2020

We studied localized surface plasmons (LSP) of a cylinder coated with a graphene monolayer. The H_z field is (p polarization):

$$H_z^{(1)}(\rho, \varphi) = \sum_{n=-\infty}^{\infty} c_n J_n(k_1 \rho) e^{in\varphi} \quad \rho < R$$

$$H_z^{(2)}(\rho, \varphi) = \sum_{n=-\infty}^{\infty} [A_o i^n J_n(k_2 \rho) + a_n H_n^{(1)}(k_2 \rho)] e^{in\varphi} \quad \rho > R$$

Applying boundary conditions to the field H_z ^[1] \rightarrow formulas a_n, c_n


Requiring the common denominator of the coefficients a_n and c_n to be zero, the following fully-retarded dispersion relation is obtained:

$$\mu_2 h_n - \mu_1 j_n + \frac{4\pi i \omega}{c} R \mu_1 \mu_2 j_n h_n = 0$$


We considered two different interior media:

- ▶ Non-dispersive (dye): the ε of the medium is $\varepsilon_1 = \text{Re}(\varepsilon_1) + i\text{Im}(\varepsilon_1)$
- ▶ Dispersive (mix of nanocrystal and dye): for the ε of the nanocrystal we used the DL model and the ε of the dye is $\varepsilon_{\text{cr}} + i\varepsilon_{\text{ci}}$

We solved the dispersion relation and we found the optical gain vs μ_c of the graphene (tunable parameter):




(a) Non-dispersive
Dielectric medium




(b) Dispersive
Metallic-like medium

Extinction cross-section near the optical gains:



(a) Non-dispersive
Dielectric medium



(b) Dispersive
Metallic-like medium

For the critical values we observed: the extinction sign changes, the absorption minimizes and the scattering maximizes