

Landolt C and snellen e acuity: differences in strabismus amblyopia?

Answer (EN)

The paper does not give a definitive answer on whether the Snellen E versus Landolt C acuity differs specifically in strabismus amblyopia. It reports that recognition of different optotypes can differ, and that conventional evaluation may miss differences in the low-acuity range—which is exactly the range where amblyopia patients fall. The stated study design covers “39 with amblyopia due to strabismus,” but the abstract stops short of reporting an explicit result for that subgroup; it only states that such differences “cannot be excluded” (arXiv:SciQ:ctx).

中文速览

论文未明确给出「斜视性弱视中 Landolt C 与 Snellen E 视力是否有差异」的定论。它指出：不同视标在同一视角下的识别率本就不同，且因视标多在正常视力者身上评价，「低视力区段的差异无法排除」——而弱视患者正处于该区段。研究纳入了 39 例斜视性弱视患者，但摘要未直接给出该亚组的明确结论 (arXiv:SciQ:ctx)。

Source evidence

“Since optotypes are evaluated on individuals with good visual acuity and without eye disorders, differences in the lower visual acuity range cannot be excluded.”

“100 patients (age 8 - 90 years, median 60.5 years) with various eye disorders, among them 39 with amblyopia due to strabismus.”

Final decision: maybe