

Does HER2 immunoreactivity provide prognostic information in locally advanced ur

Answer (EN)

No conclusive answer can be extracted. The source excerpt (arXiv:SciQ:ctx) describes an evaluation of HER2 immunoreactivity in 114 formalin-fixed paraffin-embedded specimens from locally advanced urothelial carcinoma patients treated with surgery alone or adjuvant M-VEC (arXiv:SciQ:ctx), and states that the authors “explored the influence of pathological parameters and HER2 immunoreactivity on progression-free survival (PFS)” (arXiv:SciQ:ctx). However, the provided text reports only the study’s aim and methods; it does not state the actual result —i.e., whether HER2 immunoreactivity was found to be a prognostic factor in this adjuvant M-VEC setting (arXiv:SciQ:ctx). Therefore, the paper, as excerpted, does not explicitly answer whether (or how) HER2 immunoreactivity provides prognostic information in these patients (arXiv:SciQ:ctx).

中文速览

所给摘要仅描述了研究目的与方法（114 例标本、免疫组化评估 HER2、探索其与 PFS 的关系），并未报告最终结论，因此无法据现有摘录确认 HER2 免疫反应性在 M-VEC 辅助化疗中是否提供预后信息。

Source evidence

“We studied 114 formalin-fixed paraffin-embedded specimens obtained from locally advanced urothelial carcinoma patients receiving surgery alone or adjuvant M-VEC.” “The authors evaluated HER2 immunoreactivity using immunohistochemical staining and explored the influence of pathological parameters and HER2 immunoreactivity on progression-free survival (PFS).” (arXiv:SciQ:ctx)

Final decision: maybe