

Syncope during bathing in infants, a pediatric form of water-induced urticaria?

Answer (EN)

Yes/maybe question: does the paper establish that bathing syncope in infants is a pediatric form of water-induced urticaria? The source paper does not definitively confirm this. It reports eight infants (2–15 months) who became pale, hypotonic, still and unreactive on bath immersion, with recovery in seconds after withdrawal. The initially considered diagnoses (seizure, gastroesophageal reflux) were doubted, and instead “the hypothesis of an equivalent of aquagenic urticaria” was considered as an alternative explanation (arXiv:SciQ:ctx). The paper presents this only as a hypothesis under consideration, not as a confirmed diagnosis or stated fact. Therefore the source supports “maybe” —the authors hypothesized a urticaria-equivalent mechanism for bathing syncope, but the claim is not asserted as established in the paper.

中文速览

结论为「可能」。论文报告了 8 名在洗澡时出现苍白、肌张力低下、静止无反应的婴儿，离开浴水并刺激后数秒恢复；曾被怀疑的发作或胃食管反流诊断不成立，于是「考虑了一种相当于水源性荨麻疹的假说」。但这仅是提出假说，论文并未确证为水源性荨麻疹的小儿类型。

Source evidence

“The hypothesis of an equivalent of aquagenic urticaria was then considered.” (arXiv:SciQ:ctx)

Final decision: maybe