

Are the long-term results of the transanal pull-through equal to those of the tr

Answer (EN)

The provided source does not state whether long-term results of the transanal pull-through (TERPT) equal those of the transabdominal (ABD) pull-through. The excerpt contains only the study design and aim, not any comparative results: it reports that records of 41 patients older than 3 years who underwent a pull-through (TERPT, $n = 20$; ABD, $n = 21$) were reviewed, and that overstretching of the anal sphincters in TERPT “remains a critical issue that may impact the continence.” No outcome data, statistical comparison, or conclusion of equivalence appears in the provided text (arXiv:SciQ:ctx).

中文速览

所给来源未给出经肛/经腹拖出术长期结果是否相当的结论。摘要仅报告研究设计与目的（共 41 例 >3 岁患儿：TERPT $n=20$ ，ABD $n=21$ ），并提示经肛术式可能对括约肌过度牵拉而影响控便，但未提供结果数据或两者对比结论（arXiv:SciQ:ctx）。

Source evidence

“This study examined the long-term outcome of TERPT versus conventional transabdominal (ABD) pull-through for HD. Records of 41 patients more than 3 years old who underwent a pull-through for HD (TERPT, $n = 20$; ABD, $n = 21$) were reviewed.”

The excerpt ends before any results are reported; the outcome comparison is not stated.