

30-Day and 1-year mortality in emergency general surgery laparotomies: an area o

Answer (EN)

Based strictly on the provided excerpt, the paper' s own 30-day and 1-year mortality results are **not stated** in this source. The text only provides:

- **Background/context:** "recent studies suggesting the 30-day mortality to be 14-15%"for emergency surgery (arXiv:SciQ:ctx).
- **Study aim:** to analyse "the 30-day mortality, age-related 30-day mortality and 1-year mortality following emergency laparotomy" (arXiv:SciQ:ctx).
- **Design:** a "retrospective study of patients who underwent emergency laparotomy from June 2010 to May 2012" (arXiv:SciQ:ctx).

No outcome figures (neither 30-day nor 1-year mortality) are reported in the provided material, so they cannot be extracted without inventing or inferring data. The question' s findings (if included in the full paper) are not present in this excerpt.

中文速览

来源摘要未给出本研究实际结果。仅含背景（近期研究提示急诊手术 30 天死亡率为 14-15%）、研究目的（分析急诊剖腹术后 30 天、年龄相关 30 天及 1 年死亡率）和研究设计（2010 年 6 月至 2012 年 5 月的回顾性队列）。原文未提供本研究的死亡率数据，无法从所给文本中提取。

Source evidence

"The aim of this study was to analyse the 30-day mortality, age-related 30-day mortality and 1-year mortality following emergency laparotomy." (arXiv:SciQ:ctx)

"recent studies suggesting the 30-day mortality to be 14-15%" (arXiv:SciQ:ctx)