

Suitably impressive thesis title

Lennart Golks

Department of Physics

University of Otago

*A thesis submitted for the degree of
Doctor of Philosophy*

November 2025

Abstract


Your abstract text goes here. Check your departmental regulations, but generally this should be less than 300 words. See the beginning of Chapter ?? for more.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Pellentesque sit amet nibh volutpat, scelerisque nibh a, vehicula neque. Integer placerat nulla massa, et vestibulum velit dignissim id. Ut eget nisi elementum, consectetur nibh in, condimentum velit. Quisque sodales dui ut tempus mattis. Duis malesuada arcu at ligula egestas egestas. Phasellus interdum odio at sapien fringilla scelerisque. Mauris sagittis eleifend sapien, sit amet laoreet felis mollis quis. Pellentesque dui ante, finibus eget blandit sit amet, tincidunt eu neque. Vivamus rutrum dapibus ligula, ut imperdiet lectus tincidunt ac. Pellentesque ac lorem sed diam egestas lobortis.

Suspendisse leo purus, efficitur mattis urna a, maximus molestie nisl. Aenean porta semper tortor a vestibulum. Suspendisse viverra facilisis lorem, non pretium erat lacinia a. Vestibulum tempus, quam vitae placerat porta, magna risus euismod purus, in viverra lorem dui at metus. Sed ac sollicitudin nunc. In maximus ipsum nunc, placerat maximus tortor gravida varius. Suspendisse pretium, lorem at porttitor rhoncus, nulla urna condimentum tortor, sed suscipit nisi metus ac risus.

Change again

Suitably impressive thesis title



Lennart Golks
Department of Physics
University of Otago

A thesis submitted for the degree of
Doctor of Philosophy

November 2025

Acknowledgements

Personal

I would like to thank Alex Elliott for his wonderful help and support. None of this would be possible otherwise.

This is where you thank your advisor, colleagues, and family and friends.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Vestibulum feugiat et est at accumsan. Praesent sed elit mattis, congue mi sed, porta ipsum. In non ullamcorper lacus. Quisque volutpat tempus ligula ac ultricies. Nam sed erat feugiat, elementum dolor sed, elementum neque. Aliquam eu iaculis est, a sollicitudin augue. Cras id lorem vel purus posuere tempor. Proin tincidunt, sapien non dictum aliquam, ex odio ornare mauris, ultrices viverra nisi magna in lacus. Fusce aliquet molestie massa, ut fringilla purus rutrum consectetur. Nam non nunc tincidunt, rutrum dui sit amet, ornare nunc. Donec cursus tortor vel odio molestie dignissim. Vivamus id mi erat. Duis porttitor diam tempor rutrum porttitor. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Sed condimentum venenatis consectetur. Lorem ipsum dolor sit amet, consectetur adipiscing elit.

Aenean sit amet lectus nec tellus viverra ultrices vitae commodo nunc. Mauris at maximus arcu. Aliquam varius congue orci et ultrices. In non ipsum vel est scelerisque efficitur in at augue. Nullam rhoncus orci velit. Duis ultricies accumsan feugiat. Etiam consectetur ornare velit et eleifend.

Suspendisse sed enim lacinia, pharetra neque ac, ultricies urna. Phasellus sit amet cursus purus. Quisque non odio libero. Etiam iaculis odio a ex volutpat, eget pulvinar augue mollis. Mauris nibh lorem, mollis quis semper quis, consequat nec metus. Etiam dolor mi, cursus a ipsum aliquam, eleifend venenatis ipsum. Maecenas tempus, nibh eget scelerisque feugiat, leo nibh lobortis diam, id laoreet purus dolor eu mauris. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Nulla eget tortor eu arcu sagittis euismod fermentum id neque. In sit amet justo ligula. Donec rutrum ex a aliquet egestas.

Institutional

If you want to separate out your thanks for funding and institutional support, I don't think there's any rule against it. Of course, you could also just remove the subsections and do one big traditional acknowledgement section.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut luctus tempor ex at pretium. Sed varius, mauris at dapibus lobortis, elit purus tempor neque, facilisis sollicitudin felis nunc a urna. Morbi mattis ante non augue blandit pulvinar. Quisque nec euismod mauris. Nulla et tellus eu nibh auctor malesuada quis imperdiet quam. Sed eget tincidunt velit. Cras molestie sem ipsum, at faucibus quam mattis vel. Quisque vel placerat orci, id tempor urna. Vivamus mollis, neque in aliquam consequat, dui sem volutpat lorem, sit amet tempor ipsum felis eget ante. Integer lacinia nulla vitae felis vulputate, at tincidunt ligula maximus. Aenean venenatis dolor ante, euismod ultrices nibh mollis ac. Ut malesuada aliquam urna, ac interdum magna malesuada posuere.

Abstract

Your abstract text goes here. Check your departmental regulations, but generally this should be less than 300 words. See the beginning of Chapter ?? for more.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Pellentesque sit amet nibh volutpat, scelerisque nibh a, vehicula neque. Integer placerat nulla massa, et vestibulum velit dignissim id. Ut eget nisi elementum, consectetur nibh in, condimentum velit. Quisque sodales dui ut tempus mattis. Duis malesuada arcu at ligula egestas egestas. Phasellus interdum odio at sapien fringilla scelerisque. Mauris sagittis eleifend sapien, sit amet laoreet felis mollis quis. Pellentesque dui ante, finibus eget blandit sit amet, tincidunt eu neque. Vivamus rutrum dapibus ligula, ut imperdiet lectus tincidunt ac. Pellentesque ac lorem sed diam egestas lobortis.

Suspendisse leo purus, efficitur mattis urna a, maximus molestie nisl. Aenean porta semper tortor a vestibulum. Suspendisse viverra facilisis lorem, non pretium erat lacinia a. Vestibulum tempus, quam vitae placerat porta, magna risus euismod purus, in viverra lorem dui at metus. Sed ac sollicitudin nunc. In maximus ipsum nunc, placerat maximus tortor gravida varius. Suspendisse pretium, lorem at porttitor rhoncus, nulla urna condimentum tortor, sed suscipit nisi metus ac risus.

Change again

Contents

List of Figures	ix
List of Abbreviations	xi
1 Introduction	1
1.1 Radiative transform equation	1
1.2 Bayes Theorem	1
1.3 GMRF	1
1.4 Inverse Rosenblatt transform	1
2 Linear Forward model	3
2.1 Sampling	3
2.1.1 two hyperparameter DAG	3
2.1.2 four hyperparameter DAG	3
2.2 Extending Model to Temperature and Pressure	3
2.2.1 Squared Inverse Rosenblatt transform	4
3 Nonlinear Forward model	5
3.1 Sampling	5
3.1.1 two or four hyperparameter DAG	5
4 Nonlinear to Linear Forward model	7
4.1 Strategie	7
4.2 local linear Map	7
4.2.1 Machine learning vs Gaussian elimination	7
Appendices	
A Posterior of Bayesian Hierachical model	11
B Convergence of the Metropolis-Hastings	13
C Randomize then Optimize - RTO	15
D Inverting Matrices - QR factorization	17

E	Taylor expansion of $g(\lambda)$	19
F	Radiation transfer and absorption line shape	21
G	whispering gallery resonator	23
	References	25

List of Figures

G.1 whispering gallery resonator	24
--	----

List of Abbreviations

i.i.d.	independent and identically distributed
MRF	Markov Random Field
GMRF	Gaussian Markov Random Field
MTC	Marginal Then Conditional sampler
GOMOS	Global Ozone Monitoring by Occultation of Stars
MCMC	Markov Chain Monte-Carlo
MH	Metropolis-Hastings

1

Introduction

- 1.1 Radiative transform equation
- 1.2 Bayes Theorem
- 1.3 GMRF
- 1.4 Inverse Rosenblatt transform

2

Linear Forward model

- algorithmic box ? or in section?
- define posterior

2.1 Sampling

2.1.1 two hyperparameter DAG

Marginal – Gibbs-MH

Conditional – RTO

2.1.2 four hyperparameter DAG

Marginal – t-walk

Conditional – RTO

2.2 Extending Model to Temperature and Pressure

- algorithmic box ? updating scheme
- write down DAG
- define posterior

2.2.1 Squared Inverse Rosenblatt transform

3

Nonlinear Forward model

- algorithmic box? updating scheme

3.1 Sampling

3.1.1 two or four hyperparameter DAG

MTC or t-walk

4

Nonlinear to Linear Forward model

4.1 Strategie

- scheme image

4.2 local linear Map

4.2.1 Machine learning vs Gaussian elimination

Appendices



Posterior of Bayesian Hierarchical model

Here we show how to obtain the posterior covariance and mean of our hierarchical Bayesian model in ?? - ??. We do not consider the hyper-parameters and start with the joint probability distribution of $(\mathbf{x}^T, \mathbf{y}^T)^T$, where $\mathbf{x} \in \mathcal{X}$ and $\mathbf{y} \in \mathcal{Y}$ do not intersect. For more details we refer to Chapter 2 in [1] and to the book of Rue and Held [2].

The exponent of the normal Gaussian can be rewritten into:

$$-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \mathbf{Q}(\mathbf{x} - \boldsymbol{\mu}) = -\frac{1}{2}\mathbf{x}^T \mathbf{Q} \mathbf{x} + \mathbf{x}^T \mathbf{Q} \boldsymbol{\mu} + \text{const.} \quad (\text{A.1})$$

We like to bring the joint distribution into a similar form so that we can compare the linear and second order terms and find the precision matrix and mean of the joint distribution.

In general the joint distribution to find the expression for the posterior distribution

We can express this posterior through the likelihood and prior probability by Bayesian theorem, with a constant and positive normalization constant:

$$\pi(\mathbf{x}|\mathbf{y}) \propto \pi(\mathbf{y}|\mathbf{x})\pi(\mathbf{x}) \quad (\text{A.2})$$

Taking the logarithmic function of this formulation we can find an expression for

the the posterior covariance, with the $\text{Var}(\mathbf{x}) = \mathbf{Q}_x^{-1}$ and $\text{Var}(\mathbf{y}) = \mathbf{Q}_y^{-1}$.

$$\ln \pi(\mathbf{x}|\mathbf{y}) \propto \ln \pi(\mathbf{y}|\mathbf{x}) + \ln \pi(\mathbf{x}) \quad (\text{A.3})$$

$$= -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \mathbf{Q}_x (\mathbf{x} - \boldsymbol{\mu}) - \frac{1}{2}(\mathbf{y} - \mathbf{A}\mathbf{x})^T \mathbf{Q}_y (\mathbf{y} - \mathbf{A}\mathbf{x}) \quad (\text{A.4})$$

$$= -\frac{1}{2} \left[\mathbf{x}^T [\mathbf{Q}_x + \mathbf{A}^T \mathbf{Q}_y \mathbf{A}] \mathbf{x} + \mathbf{x}^T [-\mathbf{A}^T \mathbf{Q}_y] \mathbf{y} \right. \quad (\text{A.5})$$

$$\left. + \mathbf{y}^T [-\mathbf{Q}_y \mathbf{A}] \mathbf{x} + \mathbf{y}^T [\mathbf{Q}_y] \mathbf{y} - 2\mathbf{x}^T \mathbf{Q}_x \boldsymbol{\mu} \right] + \text{const.} \quad (\text{A.6})$$

Hence we deal with a Gaussian distribution, we consider second order terms only and rearrange to the precision matrix.

$$-\frac{1}{2} \begin{bmatrix} \mathbf{x}^T [\mathbf{Q}_x + \mathbf{F}^T \mathbf{Q}_y \mathbf{F}] + \mathbf{y}^T [-\mathbf{Q}_y \mathbf{F}] & \mathbf{y}^T [\mathbf{Q}_y] + \mathbf{x}^T [-\mathbf{F}^T \mathbf{Q}_y] \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \quad (\text{A.7})$$

$$= \begin{bmatrix} \mathbf{x}^T & \mathbf{y}^T \end{bmatrix} \underbrace{\begin{bmatrix} \mathbf{Q}_x + \mathbf{F}^T \mathbf{Q}_y \mathbf{F} & -\mathbf{F}^T \mathbf{Q}_y \\ -\mathbf{Q}_y \mathbf{F} & \mathbf{Q}_y \end{bmatrix}}_{\text{precision matrix}} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \quad (\text{A.8})$$

We denote the precision matrix of the joint field as:

$$\mathbf{Q}_{xy} = \begin{bmatrix} \mathbf{Q}_{aa} & \mathbf{Q}_{ab} \\ \mathbf{Q}_{ba} & \mathbf{Q}_{bb} \end{bmatrix} = \begin{bmatrix} \mathbf{Q}_x + \mathbf{F}^T \mathbf{Q}_y \mathbf{F} & -\mathbf{F}^T \mathbf{Q}_y \\ -\mathbf{Q}_y \mathbf{F} & \mathbf{Q}_y \end{bmatrix} \quad (\text{A.9})$$

The mean is defined through the linear term.

$$\frac{-2\mathbf{x}^T \mathbf{Q}_x \boldsymbol{\mu}}{-2} = \begin{bmatrix} \mathbf{x}^T & 0 \end{bmatrix} \begin{bmatrix} \mathbf{Q}_x \boldsymbol{\mu} \\ 0 \end{bmatrix} \quad (\text{A.10})$$

Comparing to the linear term of Equation A.1 we can formulate an expression for the joint mean:

$$\Rightarrow \boldsymbol{\mu}_{xy} = \mathbf{Q}_{xy}^{-1} \begin{bmatrix} \mathbf{Q}_x \boldsymbol{\mu} \\ 0 \end{bmatrix} \quad (\text{A.11})$$

The mean of the conditional distribution $\mathbf{x}|\mathbf{y}$ is given by:

$$\boldsymbol{\mu}_{x|\mathbf{y}} = \boldsymbol{\mu}_x + \mathbf{Q}_{ba}^{-1} \mathbf{Q}_{ab} (\mathbf{x} - \boldsymbol{\mu}_y) \quad (\text{A.12})$$

$$\boldsymbol{\mu}_{x|\mathbf{y}} = \boldsymbol{\mu} + (\mathbf{Q}_x + \mathbf{F}^T \mathbf{Q}_y \mathbf{F})^{-1} \mathbf{F}^T \mathbf{Q}_y (\mathbf{x} - \mathbf{F}\boldsymbol{\mu}), \quad (\text{A.13})$$

and the covariance of $\mathbf{x}|\mathbf{y}$ is given by:

$$\mathbf{Q}_{x|\mathbf{y}} = \mathbf{Q}_{aa} = \mathbf{Q}_x + \mathbf{F}^T \mathbf{Q}_y \mathbf{F}, \quad (\text{A.14})$$

as illustrated through Theorem 2.5 in [2].

B

Convergence of the Metropolis-Hastings

If we show that the detailed balance condition holds and that the state space is irreducible and aperiodic under the transition matrix \mathbf{P} , we generate a Markov chain with a unique stationary distribution proportional to $\pi(\mathbf{x}, \boldsymbol{\theta}|\mathbf{y})$. Since the posterior is strictly positive $\pi(\mathbf{x}, \boldsymbol{\theta}|\mathbf{y}) \geq 0$ on the finite state space $\Omega(\mathcal{X}, \theta)$ the generated chain is irreducible. Further, it is possible to reject any proposed state and stay in the current state, which leads to aperiodicity. The detailed balance holds for the case that $\mathbf{j} = \mathbf{i}$, but if $\mathbf{j} \neq \mathbf{i}$ it is not trivial. In case we accept $\{\mathbf{x}, \boldsymbol{\theta}\}^{(n+1)} = \mathbf{j}$ as the new state we have $\pi(\mathbf{j}|\mathbf{y})g(\mathbf{i}|\mathbf{j}) > \pi(\mathbf{i}|\mathbf{y})g(\mathbf{j}|\mathbf{i})$. This gives us $\alpha(\mathbf{j}|\mathbf{i}) = 1$ and $\alpha(\mathbf{i}|\mathbf{j}) = \frac{\pi_{\mathbf{i}}g(\mathbf{j}|\mathbf{i})}{\pi_{\mathbf{j}}g(\mathbf{i}|\mathbf{j})}$ and satisfies the detailed balance:

$$\cancel{\pi_{\mathbf{j}}} \frac{\pi_{\mathbf{i}}}{\cancel{\pi_{\mathbf{j}}}} g(\mathbf{j}|\mathbf{i}) = \pi_{\mathbf{i}} g(\mathbf{j}|\mathbf{i}) \quad .$$

If $\pi(\mathbf{j}|\mathbf{y})g(\mathbf{i}|\mathbf{j}) < \pi(\mathbf{i}|\mathbf{y})g(\mathbf{j}|\mathbf{i})$ then $\alpha(\mathbf{i}|\mathbf{j}) = 1$ and $\alpha(\mathbf{j}|\mathbf{i}) = \frac{\pi_{\mathbf{j}}g(\mathbf{i}|\mathbf{j})}{\pi_{\mathbf{i}}g(\mathbf{j}|\mathbf{i})}$, this satisfies the detailed balance as well.

In conclusion the Metropolis-Hastings algorithm samples from a unique distribution proportional to the posterior distribution.



Randomize then Optimize - RTO

$$\pi(\mathbf{x}|\mathbf{y}, \boldsymbol{\theta}) \propto \pi(\mathbf{y}|\mathbf{x}, \boldsymbol{\theta})\pi(\mathbf{x}|\boldsymbol{\theta}) \quad (\text{C.1})$$

$$\propto \exp \left[(\mathbf{F}\mathbf{x} - \mathbf{y})^T \boldsymbol{\Sigma}^{-1} (\mathbf{F}\mathbf{x} - \mathbf{y}) + (\mathbf{x} - \boldsymbol{\mu})^T \mathbf{Q} (\mathbf{x} - \boldsymbol{\mu}) \right] \quad (\text{C.2})$$

$$= \exp \|\hat{\mathbf{F}}\mathbf{x} - \hat{\mathbf{y}}\|^2 \quad (\text{C.3})$$

where

$$\hat{\mathbf{F}} = \begin{bmatrix} \boldsymbol{\Sigma}^{-1/2} \mathbf{F} \\ \mathbf{Q}^{1/2} \end{bmatrix}, \quad \hat{\mathbf{y}} = \begin{bmatrix} \boldsymbol{\Sigma}^{-1/2} \mathbf{y} \\ \mathbf{Q}^{1/2} \boldsymbol{\mu} \end{bmatrix} \quad (\text{C.4})$$

One sample from the posterior can be computed by minimizing the following with respect to \mathbf{x}

$$\mathbf{x} = \arg \min_{\mathbf{x}} \|\hat{\mathbf{F}}\mathbf{x} - (\hat{\mathbf{y}} + \boldsymbol{\eta})\|^2, \quad \boldsymbol{\eta} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}) \quad (\text{C.5})$$

We can solve this and rewrite to

$$\frac{\partial}{\partial \mathbf{x}} \left[(\hat{\mathbf{F}}\mathbf{x} - (\hat{\mathbf{y}} + \boldsymbol{\eta}))^T (\hat{\mathbf{F}}\mathbf{x} - (\hat{\mathbf{y}} + \boldsymbol{\eta})) \right] = 0 \quad (\text{C.6})$$

$$\Leftrightarrow \mathbf{x}^T \hat{\mathbf{F}}^T \hat{\mathbf{F}} + \hat{\mathbf{F}}^T \hat{\mathbf{F}} \mathbf{x} - \hat{\mathbf{F}}^T (\hat{\mathbf{y}} + \boldsymbol{\eta}) - (\hat{\mathbf{y}} + \boldsymbol{\eta})^T \hat{\mathbf{F}} \mathbf{x} = 0 \quad (\text{C.7})$$

We can argue through the symmetry of the inner product that and the symmetry of the precision matrix

$$\hat{\mathbf{F}}^T \hat{\mathbf{F}} \mathbf{x} = \hat{\mathbf{F}}^T (\hat{\mathbf{y}} - \boldsymbol{\eta}) \quad (\text{C.8})$$

$$\Leftrightarrow (\mathbf{F}^T \mathbf{Q}_y \mathbf{F} + \mathbf{Q}) \mathbf{x} = \mathbf{F}^T \mathbf{Q}_y \mathbf{y} + \mathbf{Q} \boldsymbol{\mu} - \hat{\mathbf{F}}^T \boldsymbol{\eta} \quad (\text{C.9})$$

If we substitute $-\hat{\mathbf{F}}^T \boldsymbol{\eta} = \mathbf{v}_1 + \mathbf{v}_2$ we end up with

$$(\mathbf{F}^T \boldsymbol{\Sigma}^{-1} \mathbf{F} + \mathbf{Q}) \mathbf{x} = \mathbf{F}^T \boldsymbol{\Sigma}^{-1} \mathbf{y} + \mathbf{Q} \boldsymbol{\mu} + \mathbf{v}_1 + \mathbf{v}_2 \quad (\text{C.10})$$

where $\mathbf{v}_1 \sim \mathcal{N}(\mathbf{0}, \mathbf{F}^T \boldsymbol{\Sigma}^{-1} \mathbf{F})$ and $\mathbf{v}_2 \sim \mathcal{N}(\mathbf{0}, \mathbf{Q})$ are independent random variables.
 mayeb introduce... x^2 time nomral variubale

D

Inverting Matrices - QR factorization

E

Taylor expansion of $g(\lambda)$

We Taylor expand the function $g(\lambda)$ around $\lambda = \lambda' - \Delta\lambda$

$$g(\lambda) = \ln \det \underbrace{(\mathbf{F}^T \mathbf{F} + \lambda \mathbf{L})}_{\mathbf{B}} \quad (\text{E.1})$$

$$g(\lambda') - g(\lambda) = \ln \det(\mathbf{F}^T \mathbf{F} + \lambda' \mathbf{L}) - \ln \det(\mathbf{F}^T \mathbf{F} + \lambda \mathbf{L}) \quad (\text{E.2})$$

$$= \ln \det \left[\frac{(\mathbf{F}^T \mathbf{F} + (\lambda + \Delta\lambda) \mathbf{L})}{(\mathbf{F}^T \mathbf{F} + \lambda \mathbf{L})} \right] \quad (\text{E.3})$$

$$= \ln \det \left[1 + \frac{\Delta\lambda \mathbf{L}}{\mathbf{B}} \right] \quad (\text{E.4})$$

$$= \sum_{r=1}^{\infty} \frac{(-1)^{r+1}}{r!} \text{tr}((\mathbf{B}^{-1} \mathbf{L})^r) (\Delta\lambda)^r \quad (\text{E.5})$$

, where we use the identity from [3] at page 29. So the derivatives of $g(\lambda)$ are:

$$g^{(r)}(\lambda) = (-1)^{r+1} \text{tr}((\mathbf{B}^{-1} \mathbf{L})^r) \quad (\text{E.6})$$

$$\approx (-1)^{r+1} \sum_{k=1}^p \mathbf{z}_k^T (\mathbf{B}^{-1} \mathbf{L})^r \mathbf{z}_k \quad (\text{E.7})$$

Here we use a Monte Carlo estimate and draw p vectors $\mathbf{z}_k \in \mathbb{R}^n$, where each vector element $z_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{U}(\{-1, 1\})$ and $i = 1, \dots, n$.

F

Radiation transfer and absorption line
shape

G

whispering gallery resonator




Figure G.1: whispering gallery resonator

References

- [1] Christopher M Bishop and Nasser M Nasrabadi. *Pattern recognition and machine learning*. Vol. 4. 4. Springer, 2006.
- [2] Havard Rue and Leonhard Held. *Gaussian Markov random fields: theory and applications*. Chapman and Hall/CRC, 2005.
- [3] Israel Gohberg, Seymour Goldberg, and Nahum Krupnik. *Traces and determinants of linear operators*. Vol. 116. Birkhäuser, 2012.