



CS215 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: wangqi@sustech.edu.cn

Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|c$, then $a|c$.



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} transitive?



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} transitive?

No. $(1, 2), (2, 1) \in R_{\neq}$ but $(1, 1) \notin R_{\neq}$.



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R transitive?



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R transitive?

Yes.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Combining Relations: Since **relations are sets**, we can *combine* relations via **set operations**.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Combining Relations: Since **relations are sets**, we can *combine* relations via **set operations**.

Set operations: **union, intersection, difference, etc.**



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, $R_2 - R_1$?



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, $R_2 - R_1$?

We may also combine relations by **matrix operations**.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$

$$S \circ R = \{(1, b), (3, a), (3, b)\}$$



Implementation of Composite

- **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$
 $R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B
 $S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C



Implementation of Composite

- **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$
 $R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B
 $S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C
 $S \circ R = \{(1, b), (1, a), (2, a)\}$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$$S \circ R = \{(1, b), (1, a), (2, a)\}$$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix}$$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$

$$M_R \odot M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^k = ? \text{ for } k > 3$$



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.

If $(a, b) \in R$ and $(b, c) \in R$, then by the definition of composition, we have $(a, c) \in R^2 \subseteq R$.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.

If $(a, b) \in R$ and $(b, c) \in R$, then by the definition of composition, we have $(a, c) \in R^2 \subseteq R$.

“only if” part: by induction.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.

How many of these pairs are there?



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.

How many of these pairs are there?

How many subsets on $n(n-1)$ elements are there?



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.



Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.



Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.



Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.

Transitive Relation: A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.



Representing Relations


- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$

Representing Relations

- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*


Representing Relations

- Some ways to represent *n*-ary relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*




Representing Relations


- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*




reflexive



irreflexive



symmetric



antisymmetric

Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ are not in R .



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ are not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ are not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?

How to make R reflexive by minimum number of additions?



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ are not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?

How to make R reflexive by minimum number of additions?

Add $(2, 2)$ and $(3, 3)$

Then $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 2), (3, 3)\} \supseteq R$



Closures of Relations

- Let $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$ on $A = \{1, 2, 3\}$.

Is this relation R reflexive?

No. Why? $(2, 2)$ and $(3, 3)$ are not in R .

The question is what is the minimal relation $S \supseteq R$ that is reflexive?

How to make R reflexive by minimum number of additions?

Add $(2, 2)$ and $(3, 3)$

Then $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 2), (3, 3)\} \supseteq R$

The minimal set $S \supseteq R$ is called the reflexive closure of R .



Reflexive Closure

- The set S is called *the reflexive closure of R* if it:



Reflexive Closure

- The set S is called *the reflexive closure of R* if it:
 - ◇ contains R
 - ◇ is reflexive
 - ◇ is minimal (is contained in every reflexive relation Q that contains R ($R \subseteq Q$), i.e., $S \subseteq Q$)



Closures on Relations

- Relations can have different properties:
 - reflexive
 - symmetric
 - transitive



Closures on Relations

- Relations can have different properties:
 - reflexive
 - symmetric
 - transitive

We define:

- reflexive closures
- symmetric closures
- transitive closures



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (symmetric closure)

$$R = \{(1, 2), (1, 3), (2, 2)\} \text{ on } A = \{1, 2, 3\}$$

What is the symmetric closure S of R ?



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (symmetric closure)

$$R = \{(1, 2), (1, 3), (2, 2)\} \text{ on } A = \{1, 2, 3\}$$

What is the symmetric closure S of R ?

$$R = \{(1, 2), (1, 3), (2, 2)\} \cup \{(2, 1), (3, 1)\}$$



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is R transitive?



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is R transitive? No

How to make it transitive?



Closures

- **Definition** Let R be a relation on a set A . A relation S on A with property P is called *the closure of R with respect to P* if S is subset of every relation Q ($S \subseteq Q$) with property P that contains R ($R \subseteq Q$).

S is the **minimal set** containing R satisfying the property P .

Example: (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is R transitive? No

How to make it transitive?

$$S = \{(1, 2), (2, 2), (2, 3)\} \cup \{(1, 3)\}$$



Transitive Closure

- We can represent the relation on the graph.



Transitive Closure

- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path



Transitive Closure

- We can represent the relation on the graph.

Finding a **transitive closure** corresponds to **finding all pairs of elements that are connected with a directed path**

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Transitive Closure


- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Transitive Closure


- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Transitive Closure



- We can represent the relation on the graph.

Finding a **transitive closure** corresponds to **finding all pairs of elements that are connected with a directed path**

Example:

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure: $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



Paths in Directed Graphs

- **Definition** A *path* from a to b in the directed graph G is a *sequence of edges* $(x_0, x_1), (x_1, x_2), \dots, (x_{n-1}, x_n)$ in G , where n is nonnegative and $x_0 = a$ and $x_n = b$. A path of length $n \geq 1$ that begins and ends at the same vertex is called a *circuit* or *cycle*.



Paths in Directed Graphs

- **Definition** A *path* from a to b in the directed graph G is a *sequence of edges* $(x_0, x_1), (x_1, x_2), \dots, (x_{n-1}, x_n)$ in G , where n is nonnegative and $x_0 = a$ and $x_n = b$. A path of length $n \geq 1$ that begins and ends at the same vertex is called a *circuit* or *cycle*.

Theorem Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.



Path Length

- **Theorem** Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.

Proof (by induction)



Path Length

- **Theorem** Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.

Proof (by induction)



Path of length 1



Path Length

- **Theorem** Let R be relation on a set A . There is a path of length n from a to b if and only if $(a, b) \in R^n$.

Proof (by induction)



Path of length 1



Path of length 1

Path of length n

Path of length $n+1$

Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$



Connectivity Relation


- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .


$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$


Example:

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:


$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$

$$R^4 = \emptyset$$



Connectivity Relation

- **Definition** Let R be a relation on a set A . The *connectivity relation* R^* consists of **all pairs** (a, b) s.t. there is a path (**of any length**) between a and b in R .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

Example:

$$A = \{1, 2, 3, 4\}$$


$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$

$$R^4 = \emptyset$$

$$R^* = \{(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)\}$$



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)

- ◇ There are at most n different elements we can visit on a path if the path does not have loops



Connectivity


- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)

- ◇ There are at most n different elements we can visit on a path if the path does not have loops



- ◇ Loops may increase the length but the same node is visited more than once



Connectivity


- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

Proof (by intuition)

- ◇ There are at most n different elements we can visit on a path if the path does not have loops



- ◇ Loops may increase the length but the same node is visited more than once



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$R^* = \bigcup_{k=1}^n R^k$$



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$R^* = \bigcup_{k=1}^n R^k$$

Theorem: The transitive closure of a relation R equals the connectivity relation R^* .



Connectivity

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$R^* = \bigcup_{k=1}^n R^k$$

Theorem: The transitive closure of a relation R equals the connectivity relation R^* .

Recall Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof



- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
2. $R^* \subseteq S$ whenever S is a transitive relation containing R



- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
2. $R^* \subseteq S$ whenever S is a transitive relation containing R

1. If $(a, b) \in R^*$ and $(b, c) \in R^*$, then there are paths from a to b and from b to c in R . Thus, there is a path from a to c in R . This means that $(a, c) \in R^*$.



- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
 2. $R^* \subseteq S$ whenever S is a transitive relation containing R
2. Suppose that S is a transitive relation containing R .



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
 2. $R^* \subseteq S$ whenever S is a transitive relation containing R
2. Suppose that S is a transitive relation containing R .
Then S^n is also transitive and $S^n \subseteq S$. Why?



Connectivity

- **Theorem:** The transitive closure of a relation R equals the connectivity relation R^* .

Proof

1. R^* is transitive
2. $R^* \subseteq S$ whenever S is a transitive relation containing R

2. Suppose that S is a transitive relation containing R .

Then S^n is also transitive and $S^n \subseteq S$. Why?

We have $S^* \subseteq S$. Thus, $R^* \subseteq S^* \subseteq S$



Find Transitive Closure

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.



Find Transitive Closure

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$\mathbf{M}_{R^*} = \mathbf{M}_R \vee \mathbf{M}_R^{[2]} \vee \mathbf{M}_R^{[3]} \vee \dots \vee \mathbf{M}_R^{[n]}$$



Find Transitive Closure

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

$$\mathbf{M}_{R^*} = \mathbf{M}_R \vee \mathbf{M}_R^{[2]} \vee \mathbf{M}_R^{[3]} \vee \dots \vee \mathbf{M}_R^{[n]}$$

Example

$$\mathbf{M}_R = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

$$\mathbf{M}_{R^*} = ?$$



Simple Transitive Closure Algorithm

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

procedure transClosure (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$A := B := \mathbf{M}_R$;

for $i := 2$ to n

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

return B

// B is the zero-one matrix for R^*



Simple Transitive Closure Algorithm

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

procedure transClosure (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$A := B := \mathbf{M}_R$;

for $i := 2$ to n

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

return B

// B is the zero-one matrix for R^*

This algorithm takes $\Theta(n^4)$ time.



Simple Transitive Closure Algorithm

- **Lemma:** Let A be a set with n elements, and R a relation on A . If there is a path from a to b with $a \neq b$, then there exists a path of length $\leq n - 1$.

procedure transClosure (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$A := B := \mathbf{M}_R$;

for $i := 2$ to n

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

return B

// B is the zero-one matrix for R^*

This algorithm takes $\Theta(n^4)$ time. Why?



Roy-Warshall Algorithm

```
procedure Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)  
  // computes  $R^*$  with zero-one matrices  
   $W := \mathbf{M}_R$ ;  
  for  $k := 1$  to  $n$   
    for  $i := 1$  to  $n$   
      for  $j := 1$  to  $n$   
         $w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$   
  return  $W$   
  //  $W$  is the zero-one matrix for  $R^*$ 
```



Roy-Warshall Algorithm

procedure Warshall (\mathbf{M}_R : zero-one $n \times n$ matrix)

// computes R^* with zero-one matrices

$W := \mathbf{M}_R$;

for $k := 1$ to n

for $i := 1$ to n

for $j := 1$ to n

$w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$

return W

// W is the zero-one matrix for R^*

$w_{ij} = 1$ means there is a path from i to j going only through nodes $\leq k$.

$$W_{ij}^{[k]} = W_{ij}^{[k-1]} \vee \left(W_{ik}^{[k-1]} \wedge W_{kj}^{[k-1]} \right)$$



Roy-Warshall Algorithm

```
procedure Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)  
  // computes  $R^*$  with zero-one matrices  
   $W := \mathbf{M}_R$ ;  
  for  $k := 1$  to  $n$   
    for  $i := 1$  to  $n$   
      for  $j := 1$  to  $n$   
         $w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$   
  return  $W$   
  //  $W$  is the zero-one matrix for  $R^*$ 
```

$w_{ij} = 1$ means there is a path from i to j going only through nodes $\leq k$.


$$W_{ij}^{[k]} = W_{ij}^{[k-1]} \vee \left(W_{ik}^{[k-1]} \wedge W_{kj}^{[k-1]} \right)$$

This algorithm takes $\Theta(n^3)$ time.



Example


Find the matrices W_0 , W_1 , W_2 , W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .



Let $v_1 = a$, $v_2 = b$, $v_3 = c$, $v_4 = d$.

Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .




Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .




Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .



Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ \textcolor{red}{1} & 0 & 1 & \textcolor{red}{1} \end{bmatrix}$$



Example

Find the matrices W_0, W_1, W_2, W_3 , and W_4 . The matrix W_4 is the **transitive closure** of R .



Let $v_1 = a, v_2 = b, v_3 = c, v_4 = d$.

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \color{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \color{red}{1} \\ 1 & 0 & 0 & 1 \\ \color{red}{1} & 0 & 1 & \color{red}{1} \end{bmatrix}$$

$$W_4 = \begin{bmatrix} \color{red}{1} & 0 & \color{red}{1} & 1 \\ 1 & 0 & 1 & \color{red}{1} \\ 1 & 0 & \color{red}{1} & 1 \\ \color{red}{1} & 0 & 1 & \color{red}{1} \end{bmatrix}$$

n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.
 - The sets A_i 's are called the *domains* of R .



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.
 - The sets A_i 's are called the *domains* of R .
 - The *degree* of R is n .



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.
 - The sets A_i 's are called the *domains* of R .
 - The *degree* of R is n .
 - R is *functional* in domain A_i if it contains **at most one** n -tuple (\dots, a_i, \dots) for any value a_i within domain A_i .



Relational Databases

- A *relational database* is essentially an *n*-ary relation *R*.



Relational Databases

- A *relational database* is essentially an *n*-ary relation R .
- A domain A_i is a *primary key* for the database if the relation R is *functional* in A_i .




Relational Databases

- A *relational database* is essentially an n -ary relation R .
- A domain A_i is a *primary key* for the database if the relation R is *functional* in A_i .
- A *composite key* for the database is a set of domains $\{A_i, A_j, \dots\}$ such that R contains **at most 1 n -tuple** $(\dots, a_i, \dots, a_j, \dots)$ for each composite value $(a_i, a_j, \dots) \in A_i \times A_j \times \dots$.



Relational Databases

E-R Diagram



Selection Operators

- Let A be any *n -ary domain* $A = A_1 \times \cdots \times A_n$, and let $C : A \rightarrow \{T, F\}$ be any *condition* (predicate) on elements (n -tuples) of A .



Selection Operators

- Let A be any *n -ary domain* $A = A_1 \times \cdots \times A_n$, and let $C : A \rightarrow \{T, F\}$ be any *condition* (predicate) on elements (n -tuples) of A .
- The *selection operator* s_C is the operator that maps any (n -ary) relation R on A to the n -ary relation of all n -tuples from R that *satisfy* C .



Selection Operators

- Let A be any n -ary domain $A = A_1 \times \cdots \times A_n$, and let $C : A \rightarrow \{T, F\}$ be any *condition* (predicate) on elements (n -tuples) of A .
- The *selection operator* s_C is the operator that maps any (n -ary) relation R on A to the n -ary relation of all n -tuples from R that *satisfy* C .

$$- \forall R \subseteq A,$$

$$\begin{aligned} s_C(R) &= R \cap \{a \in A \mid s_C(a) = T\} \\ &= \{a \in R \mid s_C(a) = T\}. \end{aligned}$$



Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$



Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$

- Suppose that we have a domain

$$\begin{aligned} &\textit{UpperLevel}(\textit{name}, \textit{standing}, \textit{ssn}) \\ &::= [(\textit{standing} = \textit{junior}) \vee (\textit{standing} = \textit{senior})] \end{aligned}$$



Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$

- Suppose that we have a domain

$$\begin{aligned} &\textit{UpperLevel}(\textit{name}, \textit{standing}, \textit{ssn}) \\ &:\equiv [(\textit{standing} = \textit{junior}) \vee (\textit{standing} = \textit{senior})] \end{aligned}$$

- Then, $\textit{SUpperLevel}$ is the selection operator that takes any relation R on A (database of students) and produces a relation consisting of just the upper-level classes (juniors and seniors).



Projection Operators

- Let $A = A_1 \times \cdots \times A_n$ be any n -ary domain, and let $\{i_k\} = (i_1, \dots, i_m)$ be a sequence of indices all falling in the range 1 to n .
i.e., where $1 \leq i_k \leq n$ for all $1 \leq k \leq m$.



Projection Operators

- Let $A = A_1 \times \cdots \times A_n$ be any *n-ary domain*, and let $\{i_k\} = (i_1, \dots, i_m)$ be a sequence of indices all falling in the range 1 to n .

i.e., where $1 \leq i_k \leq n$ for all $1 \leq k \leq m$.

- Then the *projection operator* on n -tuples

$$P_{\{i_k\}} : A \rightarrow A_{i_1} \times \cdots \times A_{i_m}$$

is defined by

$$P_{\{i_k\}}(a_1, \dots, a_n) = (a_{i_1}, \dots, a_{i_m})$$



Projection Example

- Suppose that we have a tenary domain

$$\textit{Cars} = \textit{Model} \times \textit{Year} \times \textit{Color} \ (n = 3)$$



Projection Example

- Suppose that we have a ternary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence $\{i_k\} = \{1, 3\} \quad (m = 2)$



Projection Example

- Suppose that we have a ternary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence $\{i_k\} = \{1, 3\}$ ($m = 2$)

- Then the projection $P_{\{i_k\}}$ simply maps each tuple $(a_1, a_2, a_3) = (model, year, color)$ to its image:

$$(a_{i_1}, a_{i_2}) = (a_1, a_3) = (model, color)$$



Projection Example

- Suppose that we have a tenary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence $\{i_k\} = \{1, 3\}$ ($m = 2$)

- Then the projection $P_{\{i_k\}}$ simply maps each tuple $(a_1, a_2, a_3) = (model, year, color)$ to its image:

$$(a_{i_1}, a_{i_2}) = (a_1, a_3) = (model, color)$$

- This operator can be usefully applied to a whole relation $R \subseteq Cars$ (database of cars) to obtain a list of *model/color* combinations available.



Join Operator

- Puts two relations together to form a sort of *combined relation*.



Join Operator

- Puts two relations together to form a sort of *combined relation*.
- If the tuple (A, B) appears in R_1 , and the tuple (B, C) appears in R_2 , then the tuple (A, B, C) appears in the *join* $J(R_1, R_2)$.



Join Operator

- Puts two relations together to form a sort of *combined relation*.
- If the tuple (A, B) appears in R_1 , and the tuple (B, C) appears in R_2 , then the tuple (A, B, C) appears in the *join* $J(R_1, R_2)$.
- A, B, C can also be sequences of elements rather than single elements.



Join Example

- Suppose that R_1 is a teaching assignment table, relating *Professors* to *Courses*.



Join Example

- Suppose that R_1 is a teaching assignment table, relating *Professors* to *Courses*.
- Suppose that R_2 is a room assignment table relating *Courses* to *Rooms* and *Times*.



Join Example

- Suppose that R_1 is a teaching assignment table, relating *Professors* to *Courses*.
- Suppose that R_2 is a room assignment table relating *Courses* to *Rooms* and *Times*.
- Then $J(R_1, R_2)$ is like your **class schedule**, listing *(professor, course, room, time)*.



Next Lecture

- relation II ...

