
Binary Star System

Zheyu Lu Rui Liu Shuqi Xu
University of California, Berkeley
{zheyulu, rui_liu, sqxu}@berkeley.edu

Abstract

We create a fictional binary star solar system which has two suns, one Earth and one moon and simulate the orbital motion of these celestial bodies. Based on the observation data from a local position on Earth, we use machine learning techniques and successfully learn the time evolution curve of the moon phase and the altitude and azimuth of the other three celestial bodies including the two suns and the moon. Furthermore, we demonstrate the underlying physics of these time evolution curves which can help us better understand the physical principles dominating the astronomical phenomena. Below is the link to our project directory:

<https://github.com/Leo-godel/Project-S-final>

1 Introduction

We create a fictional binary star solar system which has two suns, one Earth and one moon. In our early project, we generate our data by using SpaceEngine and OCR. This time, however, we come up with a new method to generate the altitude and azimuth data of the celestial bodies by analytically solving our model which generalized and simplified our previous method. Our current method can be used to quickly generate observation data from an arbitrary position on Earth. Combining the two methods can provide us with abundant data for learning the orbital motion and positions of the celestial bodies and the moon phase.

Our results can be used to further study and predict both solar and lunar eclipse phenomena since we can accurately predict the time evolution curve of the sun's and the moon's position on the sky. That will be an interesting work for us in the future.

The paper is organized as follows: In Section 2, we firstly introduce data generation in Section 2.1 which includes the two different methods we used to generate our data and we also verify the consistency between using the two methods. For the second method, we also introduce the theory and show the analytical derivation of the coordinate transformation. In Section 2.2, we discuss the physical intuition when we analyzed the data and chose the relevant features. This can be viewed as preprocessing the data with reasonable physical consideration and certain domain knowledge. The main learning results of the position of the celestial bodies and the moon phase are shown in Section 3. In Section 4, we conclude the results we learned and also discuss the underlying physics which is related to the observed astronomical phenomena.

2 Method

2.1 Data generation

2.1.1 First method: SpaceEngine screenshots & OCR

One way to generate data is to simulate the binary star system in SpaceEngine. But SpaceEngine doesn't provide users with API functions to directly export data, so we can only take screenshots and

then use optical character recognition (OCR) to extract data from the raw pictures. We managed to generate 27,000 data points with 3-year span in 3 different locations: Berkeley, Shanghai and Sydney with very high recognition accuracy (at least 99%). The data includes time, spherical coordinates, *i.e.*, altitude and azimuth of two suns and one moon in observer's topocentric coordinate system, the size of the three celestial bodies, and the phase of the moon.

2.1.2 Second method: analytical derivation of the altitude and azimuth of celestial body

In addition to the data generated by using OCR on screenshots from SpaceEngine, we also simulated all celestial bodies' orbital motion by using Mathematica in the two-sun center of mass coordinate system, where the four celestial bodies, *i.e.*, the two suns, the Earth, and the moon, are arranged to be along the x -axis at $t = 0$ as shown in Figure 1.




Figure 1: The initial condition of the binary star system

To begin with, we generate the orbital motion, *i.e.*, x, y, z coordinates for all celestial bodies in the coordinate system S shown in Figure 1. Our goal is to transform these coordinates into observations (altitude and azimuth) in observer's topocentric coordinate system S_3 on Earth (defined below). In order to derive such transformation, we need to introduce three coordinate systems:

1. The center of mass reference system of Earth without rotation. We call it S_1 which is shown in Figure 2.
2. The fixed center of mass reference system of Earth. We call it S_2 which is shown in Figure 3. The x -axis is pointing to the intersection of equator and prime meridian and the z -axis is pointing toward north pole. ξ_0 here is the obliquity of Earth. S_2 and S_1 are related by the Euler angles $(\frac{\pi}{2}, \xi_0, \gamma_0)$.
3. The local observer's reference system which is fixed relative to the Earth. We call it S_3 which is shown in Figure 4. The x -axis is pointing toward south and the y -axis is pointing toward east. S_2 and S_3 are related by the Euler angles $(\phi, \theta, 0)$ and a translation with length R_e , *i.e.*, the radius of the Earth. Note that the latitude and longitude are given by $90 - \theta$ and ϕ respectively.

Now, for any celestial body, we want to find the transformation between (x, y, z) and (x_3, y_3, z_3) .

First, we transform (x, y, z) into (x_1, y_1, z_1) which is simply given by

$$\begin{aligned} x_1 &= y_e - y \\ y_1 &= -x_e + x \\ z_1 &= 0 \end{aligned} \tag{1}$$

where (x_e, y_e, z_e) is the Earth's coordinates in S .

Finally, we transform (x_2, y_2, z_2) into (x_3, y_3, z_3) . As we mentioned, S_3 and S_2 are related by the Euler angles $(\phi, \theta, 0)$ so that we have

$$\begin{pmatrix} x_3 \\ y_3 \\ z_3 \end{pmatrix} = R^{-1}(\phi, \theta, 0) \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} \quad (5)$$

so that the transformation is given by

$$\begin{aligned} x_3 &= \cos \phi \cos \theta \left[\cos \xi_0 \cos (\gamma_0 + \Omega t) x_1 + \sin (\gamma_0 + \Omega t) y_1 \right] \\ &\quad + \sin \phi \cos \theta \left[-\cos \beta_0 \sin (\gamma_0 + \Omega t) x_1 + \cos (\gamma_0 + \Omega t) y_1 \right] \\ &\quad - \sin \theta \sin \xi_0 x_1 \\ y_3 &= -\sin \phi \left[\cos \xi_0 \cos (\gamma_0 + \Omega t) x_1 + \sin (\gamma_0 + \Omega t) y_1 \right] \\ &\quad + \cos \phi \left[-\cos \beta_0 \sin (\gamma_0 + \Omega t) x_1 + \cos (\gamma_0 + \Omega t) y_1 \right] \\ z_3 &= \cos \phi \sin \theta \left[\cos \xi_0 \cos (\gamma_0 + \Omega t) x_1 + \sin (\gamma_0 + \Omega t) y_1 \right] \\ &\quad + \sin \phi \sin \theta \left[-\cos \beta_0 \sin (\gamma_0 + \Omega t) x_1 + \cos (\gamma_0 + \Omega t) y_1 \right] \\ &\quad + \cos \theta \sin \beta_0 x_1 - R_e \end{aligned} \quad (6)$$

and the altitude and azimuth are simply given by

$$\begin{aligned} \text{altitude} &= \arctan\left(\frac{z}{\sqrt{x^2 + y^2}}\right) \\ \text{azimuth} &= \arctan\left(\frac{y}{x}\right) \end{aligned} \quad (7)$$

Note that in theory, one can generate the azimuth and the altitude data in any arbitrary location on Earth which paves the way for further predicting when and where on Earth can we observe both solar and lunar eclipses.

2.1.3 Data verification

The above mentioned two completely different methods for generating data yield almost identical results. Figure 5 and Figure 6 show the comparison between the azimuth and the altitude data of a sun measured in Berkeley generated by the two different methods. Notice that the number of orange points is much smaller than that of the blue points since the consumption of generating data by using SpaceEngine is much higher than that by using analytical model.




Figure 5: Azimuth data comparison




Figure 6: Altitude data comparison

The definition of the altitude in SpaceEngine is different from the traditional topocentric coordinate system. The altitude in SpaceEngine ranges from -180° to 180° and 0° is defined to point to the South pole, while the traditional topocentric coordinate system ranges from 0° to 360° and it starts from the North pole. We decided to follow the traditional topocentric coordinate system definition in the following training part.

2.2 Physical reasoning of the characteristic frequency

From the time evolution plot of the altitude, we can notice that this quantity is periodic. In order to get a sense of what features are relevant in learning such 1-d curve, we firstly use matlab to do a rough fitting of the time evolution curve using a physically reasonable function given by:

$$f(t) = A_1 \sin(\omega_1 t + \phi_1) + A_2 \sin(\omega_2 t + \phi_2) \quad (8)$$

The fitting result is shown in Figure 7 for the result of sun and in Figure 8. It is obvious that there are two main characteristic frequencies which are physically important in our system, that is the two frequencies corresponding to the two angular velocities given by

$$\omega_1 = 2\pi f_1$$

$$\omega_2 = 2\pi f_2$$

Therefore, it is reasonable to use the combination of the Fourier features $\sin \omega_1 t$, $\cos \omega_1 t$, $\sin \omega_2 t$, and $\cos \omega_2 t$ to learn such 1-d function.

Actually, these two frequencies are corresponding to the self-rotation of the Earth and the revolution of the Earth in our binary star solar system which is physically reasonable.




Figure 7: Matlab fitting result of the altitude time evolution curve of the sun.




Figure 8: Matlab fitting result of the altitude time evolution curve of the moon.

2.3 Polynomial Fourier features for learning time evolution curve

In our learning process, we choose "polynomial-type" combination of the Fourier features mentioned above as our features in order to capture any additional "nonlinear correlation effect" in the sense

of Fourier features. That is, for example, we use the following kind of features for learning in the degree = 2 case:

$$\begin{aligned} &1, \sin(\omega_1 t), \cos(\omega_1 t), \sin(\omega_2 t), \cos(\omega_2 t), \sin^2(\omega_1 t), \cos^2(\omega_1 t), \sin^2(\omega_2 t), \cos^2(\omega_2 t) \\ &\sin(\omega_1 t) \cos(\omega_1 t), \sin(\omega_1 t) \sin(\omega_2 t), \sin(\omega_1 t) \cos(\omega_2 t), \cos(\omega_1 t) \sin(\omega_2 t) \\ &\sin(\omega_1 t) \cos(\omega_2 t), \sin(\omega_2 t) \cos(\omega_2 t) \end{aligned}$$

It is clearly that the degree parameter D plays a role as a hyperparameter in our training process which can take values larger than 1. Considering the number of polynomial features scales roughly as D^3 which increases very fast, we focus on using D in range 2 to 5. Through our training, we pick up the optimal hyperparameter $D = 3$ considering both performance and the time consumption.

3 Results

Our learning results for the sun and the moon are shown in Figure 9 and Figure 10 respectively. The mean squared error(MSE) of the test data for the sun's altitude is 9.4 which means that our prediction only deviates from the true value by around $\sqrt{9.4} \approx 3$ degrees on average which is very accurate. The MSE of the test data for the moon's altitude is 27.6 which means that our prediction deviates from the true value by around $\sqrt{27.6} \approx 5$ degrees on average which is also very accurate.

Our learning result for the moon phase is shown in Figure 11. The MSE of the test data for the moon phase is 0.002 which is clearly very accurate.




Figure 9: Learning result of the altitude time evolution curve of the sun. The curve shows performance on test dataset. Orange dots are the prediction on test data and blue dots are the original test data.




Figure 10: Learning result of the altitude time evolution curve of the moon. The curve shows performance on test dataset. Orange dots are the prediction on test data and blue dots are the original test data.

4 Conclusion

1. There are three main physically important characteristic frequencies in our system which are corresponding to the self-rotation of the Earth, the revolution of the Earth, and the revolution of the moon.




Figure 11: Learning result of the phase time evolution curve of the moon. The curve shows performance on test dataset. Orange curve is the prediction on test data and blue curve is the original test data.

2. The time evolution of the altitude and azimuth of the two suns and the moon are periodic and the periodicity is captured by the three characteristic frequencies mentioned above.
3. The azimuth curve of the sun basically has the period of one day since it will always change from 0 to 360 within one day.
4. Similarly, the azimuth curve of the moon also has the period of one day since it will also change from 0 to 360 within one day.
5. The altitude curve of the sun is basically a combination of two sinusoidal curves with two frequencies corresponding to the self-rotation of the Earth and the revolution of the Earth. This is physically reasonable.
6. The altitude curve of the moon is basically a combination of two sinusoidal curves with two frequencies corresponding to the self-rotation of the Earth and the revolution of the moon. This is physically reasonable.
7. The phase of the moon has the periodicity of around one month which is around 29.5 days.

References

[1] <https://www.wikipedia.org/>