
a Final AL performance variability by dataset
P. aeruginosa AMPs
SBI STAAM
E. coli AMPs
S. aureus AMPs
HemoPI2
ENVZ ECOLI
FKBP3 HUMAN
DN7A SACS2
SOX30 HUMAN
MAFG MOUSE sub
SDA BACSU
YNZC BACSU sub
POLG PESV sub
A0A247D711 LISMN
Substitution
AMP
HemoPI2
Std of final AL performance (Top-10% acquired %) across 10 seeds
b Per-seed selection trajectories
MAFG MOUSE sub
E. coli AMPs
Top-10% acquired (%)
Sequences screened
c Seed spread across screening rounds
Substitution
AMP
HemoPI2
Std across seeds of Top-10% acquired (%)
Sequences screened
d Final AL performance vs initial-seed label spread
Pearson r = 0.11
Final AL performance (x random)
E. coli AMPs
P. aeruginosa AMPs
HemoPI2
S. aureus AMPs
Initial-seed label spread, relative to this dataset
(0 = average seed, +1 = one std above)
e Final AL performance vs best initial-seed label
Pearson r = 0.18
Final AL performance (x random)
E. coli AMPs
P. aeruginosa AMPs
HemoPI2
S. aureus AMPs
Best initial-seed label, min-max within dataset
(0 = pool worst, 1 = pool best)
f Final AL performance vs initial-seed length range
Pearson r = 0.30
Final AL performance (x random)
E. coli AMPs
P. aeruginosa AMPs
HemoPI2
S. aureus AMPs
Initial-seed sequence-length range, min-max within dataset
(0 = all equal length, 1 = full dataset span)