
BIG DATA COMPUTING

Homework 1

Concentration of measures and statistical significance

Leonardo Di Nino : 1919479

October 21, 2023

Assignment 1

Question (a)

Given such a random graph model $G(V, E)$ where $|V| = n$ and $\mathbb{P}((u, v) \in E) = p$ for each $u, v \in V$, the total number of possible cliques of size k is given by $C = \mathcal{C}_{n,k} = \binom{n}{k}$, i.e. the number of subset of K nodes. Each of these cliques is associated to a different subset of nodes $\{V'_i\}_{i=1}^C$ such that the clique exists only if the induced subgraph $G(V'_i)$ has a number of edges equal to $\binom{k}{2}$. Stating that each edge is a random variable independent from all the others, we can then associate to each of those cliques a binary random variable $Z_i \sim \text{Ber}(q)$: $Z_i = 1$ if the cliques exists, with a probability equal to $q = p^{\binom{k}{2}} = \frac{k(k-1)}{2}$ since the clique needs such a number of edges in order to exists. So we have the succession of random variables $\{Z_i\}_{i=1}^C$ and $Z = \sum_{i=1}^C Z_i$ is the total number of cliques of size k . Finally we have the following result just leveraging the linearity of expected value:

$$\mathbb{E}[Z] = \mathbb{E}\left[\sum_{i=1}^C Z_i\right] = \sum_{i=1}^C \mathbb{E}[Z_i] = Cq = \binom{n}{k} p^{\frac{k(k-1)}{2}} \quad (1)$$

This is a closed form solution for $\mathbb{E}[Z]$, but we can easily bound it since we have the following inequality for binomial coefficients:

$$\left(\frac{n}{k}\right)^k \leq \binom{n}{k} \leq \left(\frac{en}{k}\right)^k \quad (2)$$

So just applying eq.(2) to eq.(1) we get:

$$\left(\frac{n}{k}\right)^k p^{\frac{k(k-1)}{2}} \leq \mathbb{E}[Z] \leq \left(\frac{en}{k}\right)^k p^{\frac{k(k-1)}{2}} \quad (3)$$

Question (b)

We now want to provide an upper bound for the probability for the existence of a clique of size at least equal to $\frac{epn}{1-\epsilon}$ for $0 < \epsilon < 1$.

Before moving forward let's consider the feasibility of the question also to figure out in which dynamic k ranges. In order to have consistency we have to impose $k \leq n$, i.e. $\frac{epn}{1-\epsilon} \leq n$, so that $0 \leq \epsilon \leq 1 - pe$. Still we want $1 - pe \geq 0$ so that also $p \geq \frac{1}{e}$. In this setting we are considering a high value for k : we are likely to consider $\epsilon \approx 0$ and $p \approx \frac{1}{e}$ to retrieve non trivial bounds, as we can see.

Now moving on with probabilistic modeling we can introduce the event I_u as the binary random variable whose realization is 1 if the clique of size u does exist. It is interesting to show that $I_{u+1} \subset I_u$ since $I_{u+1} = I_u \cap N_{u+1}$ being N_u the event "There is a $(u+1)$ -th node that is connected to all the u nodes forming a u sized clique". We have that the following holds:

$$\begin{aligned} I_k &\supset I_{k+1} \supset I_{k+2} \supset \dots \supset I_n \\ \mathbb{P}(I_k) &> \mathbb{P}(I_{k+1}) > \mathbb{P}(I_{k+2}) > \dots > \mathbb{P}(I_n) \end{aligned} \quad (4)$$

This few considerations in modeling are actually very important, since the event of interest is that the graph has at least a clique of at least size k :

$$\mathbb{P}\left(\bigcup_{u=k}^n (I_u = 1)\right) = \mathbb{P}(I_k = 1) \quad (5)$$

since we have $I_k \supset I_{k+1} \supset I_{k+2} \supset \dots \supset I_n$ we have that $\bigcup_{u=k}^n (I_u = 1) = I_k$.

Finally we can just consider to bound $\mathbb{P}(I_k = 1)$. Firstly stating that Z still is the total number of cliques of size k and we are considering that a clique must exist we can leverage the

fact that all Z_i are binary random variables as defined in question (a), so that:

$$\mathbb{P}(I_k = 1) = \mathbb{P}(Z \geq 1) = \mathbb{P}\left(\sum_{i=1}^C Z_i \geq 1\right) = \mathbb{P}\left(\bigcup_{i=1}^C (Z_i = 1)\right) \quad (6)$$

Now we can apply Boole's inequality to eq.(6):

$$\mathbb{P}\left(\bigcup_{i=1}^C (Z_i = 1)\right) \leq \sum_{i=1}^C \mathbb{P}(Z_i = 1) \quad (7)$$

from this last naturally follows the following, since $\sum_{i=1}^C \mathbb{P}(Z_i = 1) = \mathbb{E}[Z]$:

$$\mathbb{P}\left(\bigcup_{i=1}^C (Z_i = 1)\right) \leq \mathbb{E}[Z] \quad (8)$$

and now we can use the upper bound we derived in eq.(3) making some further assumptions leveraging $p \in [0, 1]$, that implies that exponentiation is decreasing when using p as a base:

$$\mathbb{P}\left(\bigcup_{i=1}^C (Z_i = 1)\right) \leq \mathbb{E}[Z] \leq \left(\frac{en}{k}\right)^k p^{\frac{k(k-1)}{2}} \leq \left(\frac{en}{k}\right)^k p^{k^2} \leq \left(\frac{en}{k}\right)^k p^k \Big|_{k=\frac{epn}{1-\epsilon}} \quad (9)$$

Chaining everything we have the following:

$$\mathbb{P}(Z \geq 1) \leq \left(\frac{epn}{k}\right)^k \quad (10)$$

We can now evaluate the term bounding on the right in $k = \frac{epn}{1-\epsilon}$:

$$\left(\frac{epn}{k}\right)^k \Big|_{k=\frac{epn}{1-\epsilon}} = \left(\frac{epn}{\frac{epn}{1-\epsilon}}\right)^{\frac{epn}{1-\epsilon}} = (1-\epsilon)^{\frac{epn}{1-\epsilon}} \quad (11)$$

Finally we have our upper bound:

$$\mathbb{P}(Z \geq 1) \leq (1-\epsilon)^{\frac{epn}{1-\epsilon}} \quad (12)$$

Assignment 2

Question (a)

Given the assumptions we can configure a proper probabilistic model. In particular being A a measurable set, we can define a uniform probability distribution over it such that for each $a \in A$ we have the following probability measure: $f(a) = \frac{1}{\mu(A)} \mathbf{1}_A(a)$, being $\mu(A)$ the Lebesgue integral over A , i.e. its very area: this models the fact that the pixels are uniformly distributed in A .

We then can consider the following probability, given that $X \subseteq A$ and basic measure theory results:

$$\mathbb{P}(a \in X | X \subseteq A) = \int_X \frac{d\mu(X)}{\mu(a)} = \frac{\mu(X)}{\mu(A)} \quad (13)$$

From now on to lighten notation we'll consider $\mu(A) = A$, $p = \mathbb{P}(a \in X | X \subseteq A)$ and $\mu(X) = X$, so that $\mathbb{P}(a \in X | X \subseteq A) = p = \frac{X}{A}$, and naturally $X = Ap$.

We want to provide an unbiased estimator of X using repeated IID sampling leveraging these uniformity assumptions. So we can define for each pixel sampled $Z \sim \text{Ber}(p)$ as the binary random variable assuming value 1 if $Z \in X$ so that this distribution is parametrized exactly by $p = \mathbb{P}(a \in X | X \subseteq A)$: this is realized by the routine `sample()`.

Given this sampling scheme and an actual sample $\{Z_i\}_{i=1}^m$, we know that the maximum likelihood estimator for p is the sample mean \bar{Z}_m and it's trivial to show that it is also an unbiased estimator for p :

$$\mathbb{E}[\bar{Z}_m] = \mathbb{E}\left[\frac{1}{m} \sum_{i=1}^m Z_i\right] = \frac{1}{m} \sum_{i=1}^m \mathbb{E}[Z_i] = \frac{1}{m} mp = p \quad (14)$$

Now we can leverage that $X = Ap$ is a function $\phi(p)$ and the fact that maximum likelihood estimation is invariant under continuous mapping: $\hat{\phi}_{MLE}(p) = \phi(\hat{p}_{MLE})$. This finally gives us the estimator for X that is $\hat{X} = A\bar{Z}_m$. Given the primitive `sample()` we have the following procedure:

Algorithm 1 Unbiased estimator of X

Require: $A : \text{float}, m : \text{int}$

$S \leftarrow \mathbf{0}_n$

▷ Pre-allocating a vector for the samples

for $i = 1, \dots, m$ **do**

$S[i] \leftarrow \text{sample}()$

end for

return $\frac{A}{n} \sum_{i=1}^m S[i]$

Question(b)

Given a precision ϵ and a degree of confidence δ we want to bound m such that

$$\mathbb{P}(|\hat{X} - X| \leq \epsilon X) \geq 1 - \delta$$

We can play a bit with this expression:

$$\begin{aligned} \mathbb{P}(|\hat{X} - X| \leq \epsilon X) &\geq 1 - \delta \\ \mathbb{P}(|A\hat{p} - Ap| \leq \epsilon X) &\geq 1 - \delta \\ \mathbb{P}(A|\hat{p} - p| \leq \epsilon X) &\geq 1 - \delta \\ \mathbb{P}(|\bar{Z}_m - \mathbb{E}[Z]| \leq \frac{\epsilon X}{A}) &\geq 1 - \delta \end{aligned} \quad (15)$$

Hoeffding's inequality is a good choice in our case, since it is an exponentially decreasing

bound. Hoeffding's inequality is usually in the form

$$\mathbb{P}(|\bar{X}_m - \mathbb{E}[X]| \leq \kappa) \geq 1 - 2e^{\frac{-2m\kappa^2}{(b-a)^2}}$$

under the hypothesis that X is a bounded random variable, which is true for a Bernoulli being $b = 1$ and $a = 0$. So finally we have the following:

$$\begin{aligned} \mathbb{P}(|\bar{Z}_m - \mathbb{E}[Z]| \leq \frac{\epsilon X}{A}) &\geq 1 - \delta \\ \mathbb{P}(|\bar{Z}_m - \mathbb{E}[Z]| \leq \frac{\epsilon X}{A}) &\geq 1 - 2e^{-2m(\frac{\epsilon X}{A})^2} \end{aligned} \tag{16}$$

Comparing directly δ and the term in Hoeffding's inequality we get:

$$\begin{aligned} \delta &= 2e^{-2m(\frac{\epsilon X}{A})^2} \\ \log\left(\frac{\delta}{2}\right) &= -2m\left(\frac{\epsilon X}{A}\right)^2 \\ m\left(\frac{\epsilon X}{A}\right)^2 &= \frac{1}{2}\log\left(\frac{2}{\delta}\right) \\ m &= \frac{A^2\log(\frac{2}{\delta})}{2\epsilon^2 X^2} \end{aligned} \tag{17}$$

The last expression is the lower bound on m respecting the initial request.

Assignment 3

Question (a)

We can model the situation as a hypothesis testing problem with the following hypothesis set:

$$\begin{cases} \mathcal{H}_0 : \text{There is at least one node whose degree is 600 as effect of randomness;} \\ \mathcal{H}_1 : \text{At least one node whose degree is 600 is a prove of existence of strong social structures} \end{cases}$$

We are going to find a formal acceptance or rejection of \mathcal{H}_0 through a bound on the p-value, i.e. the probability for the observed data under the null hypothesis.

Question (b)

We can be a little bit more formal in defining what we are dealing with. First of all we can retrieve the average degree from $|V| = 5000, |E| = 10^6$: $\hat{d} = \frac{2|E|}{|V|} = 400$.

Now we'll focus on a single node of interest u and we'll define D_u as the random variable that associate to u its degree. From the assumptions $\mathbb{E}[D_u] = \hat{d} = 400$, and given this we can also retrieve the following estimate for the parameter p characterizing the underlying Erdős–Rényi model in the null hypothesis: $\hat{p} \approx \frac{\hat{d}}{|V|} = 0.08$.

We then can model the degree of a fixed node through a random variable D that under \mathcal{H}_0 is equal to the sum of IID bernoulli random variables; in particular $D_u = \sum_{i=1}^{|V|-1} e_i$, being in this case $e_i \sim \text{Ber}(\hat{p})$ the bernoulli for the existence of the edge between the fixed node in question and all the other nodes. We have from the assumptions that $\mathbb{E}[D] = 400$: all these can give us the following upper tail Chernoff bound over the probability that $D \geq 600$:

$$\mathbb{P}(D \geq (1 + \delta)\mathbb{E}[D]) \leq \exp\left\{-\frac{\delta^2}{2 + \delta}\mathbb{E}[D]\right\} \quad (18)$$

being $(1 + \delta)\mathbb{E}[D] = 400(1 + \delta) = 600$. So we have $\delta = \frac{1}{2}$ and the following bound holds:

$$\mathbb{P}(D \geq 600) \leq \exp\left\{-\frac{\delta^2}{2 + \delta}\mathbb{E}[D]\right\} \Big|_{\delta=\frac{1}{2}} \quad (19)$$

$$\mathbb{P}(D \geq 600) \leq \exp\left\{-\frac{\frac{1}{4}}{\frac{5}{2}} * 400\right\}$$

$$\mathbb{P}(D \geq 600) \leq \exp -40$$

$$\mathbb{P}(D \geq 600) \leq 4.24 * 10^{-18}$$

What we are interested in is the probability that at least one vertex has such a degree: so we consider now the random variables $\{D_u\}_{u=1}^{|V|}$ and the following holds through Boole's inequality:

$$\mathbb{P}\left(\bigcup_{u=1}^{|V|} (D_u \geq 600)\right) \leq \sum_{u=1}^{|V|} \mathbb{P}(D_u \geq 600) \quad (20)$$

Since for each term in the right side of this last inequality eq.(19) holds, we can bound each term in the sum and get the following:

$$\mathbb{P}(\bigcup_{u=1}^{|V|}(D_u \geq 600)) \leq \sum_{u=1}^{|V|} \mathbb{P}(D_u \geq 600) \leq \sum_{u=1}^{|V|} 4.24 * 10^{-18} \quad (21)$$

Finally we have

$$\mathbb{P}(\bigcup_{u=1}^{|V|}(D_u \geq 600)) \leq 2.12 * 10^{-14} \quad (22)$$

Now consider a standard level for hypotesis testing $\alpha = 0.05$; due the Bonferroni correction we have that the actual level for our test is $\alpha' = \frac{\alpha}{|V|} = 10^{-5}$. The strong upper bound we have on the p-value implies the rejection of the null hypotesis for every level of confidence higher than $2.12 * 10^{-14}$, as α' effectively is. In the end, in support against the thesis of prof. Knowitbetter, we could not but reject \mathcal{H}_0 .