



Hilbert functions of chopped ideals

Nonlinear Algebra Seminar

MAX PLANCK INSTITUTE
FOR MATHEMATICS
IN THE SCIENCES

Leonie Kayser (joint with Fulvio Gesmundo & Simon Telen)

kayser@mis.mpg.de

June 21, 2023



Motivation: Eigenvalue methods for polynomial system solving

Task: Given 0-dim'l system $J \subseteq S = \mathbb{C}[x_0, \dots, x_n]$, compute $Z = \{z_1, \dots, z_r\} = V(J) \subseteq \mathbb{P}^n$

- ▶ For t large enough, $h_{S/J}(t) := \dim_{\mathbb{C}}(S/J)_t = r$ and $J_t = I(Z)_t$
- ▶ **Multiplication map:** $g \in S_e$, $M_g: (S/J)_d \xrightarrow{g} (S/J)_{d+e}$
- ▶ Under “suitable conditions” $M_h^{-1}M_g: (S/J)_d \rightarrow (S/J)_d$ has left eigenpairs

$$\{ (\text{ev}_{z_i}, \frac{g}{h}(z_i)) \mid i = 1, \dots, r \}, \quad \text{ev}_{z_i}(f) = f(z_i)/h(z_i)$$

↪ Translate problem into large eigenvalue problem, solve numerically

- ▶ For this need $h_{S/J}(d+e) = h_{S/J}(d) = r$, want $d, d+e$ **as small as possible**

Example: J saturated

If $J = I(Z)$ and Z is a general set of points, then $h_{S/I(Z)} = \min\{h_S(t), r\}$.

Hence $d = \min\{t \mid h_S(t) \geq r\}$ and $e = 1$ work.

Motivation: Symmetric tensor decomposition

Task: Given $F \in T = \mathbb{C}[X_0, \dots, X_n]$ of degree D , calculate decomposition

$$F = L_1^D + \dots + L_r^D, \quad L_i \in T_1, \quad r = \text{rk}(F) \text{ minimal}$$

▷ If $r < h_S(\lfloor \frac{D}{2} \rfloor) - n$, then **generically unique** summands

$$[F] \dashrightarrow Z = \{[L_1], \dots, [L_r]\} \subseteq \mathbb{P}(T_1)$$

▷ Equations of Z are contained in the kernel of the *Catalecticant map*

$$C_F(d, D-d): S_d \rightarrow T_{D-d}, \quad g \mapsto g(\partial_0, \dots, \partial_n)F$$

▷ $I(Z)_d \subseteq \text{Ker } C_F(d, D-d)$ with equality for $d \leq \lfloor \frac{D}{2} \rfloor$ and F general

↪ Obtain all equations on Z in a single *low degree* d

Key example: $F \in \mathbb{C}[X_0, X_1, X_2]_{10}$ of rank 18, obtain equations of degree ≤ 5

The funny word in the title

Definition (Chopped ideal)

The *chopped ideal* of a homogeneous ideal $I \subseteq S$ in degree d is $I_{\langle d \rangle} := \langle I_d \rangle_S$.

From now on $Z \subseteq \mathbb{P}^n$ is a general set of r points,
 $I = I(Z)$, $d = \min \{ t \mid h_S(t) \geq r \}$.

- ▷ Can we recover Z from $I(Z)_{\langle d \rangle}$?
- ▷ When does $(I(Z)_{\langle d \rangle})_{d+e} = I(Z)_{d+e}$?
- ▷ What is the Hilbert function $h_{I(Z)_{\langle d \rangle}}(t)$?



Example: $Z = 18$ points in the plane

t	...	3	4	5	6	7
$h_S(t)$...	10	15	21	28	36
$h_I(t)$...	0	0	3	10	18
$h_{I_{\langle 5 \rangle}}(t)$...	0	0	3	9	18





Figure 1: Three quintics $\langle q_1, q_2, q_3 \rangle_{\mathbb{C}} = I_5$ passing through 18 general points (left) and the missing split sextic $cc' \in I_6$ (right).

t	0	1	2	3	4	5	6	7
$h_S(t)$	1	3	6	10	15	21	28	36
$h_{S/I}(t)$	1	3	6	10	15	18	18	18
$h_{S/I_{\langle 5 \rangle}}(t)$	1	3	6	10	15	18	19	18



Recovering the points from their chopped ideal

▷ Generally $I_{\langle d \rangle} \subsetneq I$, but maybe

$$I \stackrel{?}{=} (I_{\langle d \rangle})^{\text{sat}} := \bigcup_{k \geq 0} (I_{\langle d \rangle} : \mathfrak{m}^k) \quad \Longleftrightarrow \quad V(I) \stackrel{?}{=} \underset{\text{schemes}}{V(I_{\langle d \rangle})} \subseteq \mathbb{P}^n$$

Theorem

Let $Z \subseteq \mathbb{P}^n$ be a general set of r points and $d \in \mathbb{N}$.

1. If $r > \binom{n+d}{n} - n$, then $V(I_{\langle d \rangle})$ is a positive-dimensional complete intersection.
2. If $r = \binom{n+d}{n} - n$, then $V(I_{\langle d \rangle})$ is a complete intersection of d^n points.
3. If $r < \binom{n+d}{n} - n$, then $I_{\langle d \rangle}$ cuts out Z scheme-theoretically.

In particular, $I = (I_{\langle d \rangle})^{\text{sat}}$ if and only if $r < \binom{n+d}{n} - n$ or $r = 1$ or $(n, r) = (2, 4)$.

Towards the expected Hilbert function

- ▶ Graded components of $I_{\langle d \rangle}$ are images of multiplication map

$$\mu_e: S_e \otimes_{\mathbb{C}} I_d \rightarrow I_{d+e}, \quad g \otimes f \mapsto g \cdot f$$

- ▶ One may expect μ_e to have *maximal rank*, i.e. to be injective or surjective:

$$h_{I_{\langle d \rangle}}(t) \stackrel{?}{=} \min\{h_I(t), h_S(t-d) \cdot h_I(d)\}$$

↪ $e = 1$: **Ideal generation conjecture (IGC)** predicting number of minimal generators of I

- ▶ This turns out to be too optimistic; μ_e has elements in its kernel, for example

$$f_1 \otimes f_2 - f_2 \otimes f_1 \in \text{Ker } \mu_d, \quad f_1, f_2 \in I_d$$

- ▶ This *does* happen, e.g. $r = 52$ points in \mathbb{P}^3 , then μ_5 does not have maximal rank

Thank you! Questions?

Better luck next time ;(

Towards the expected Hilbert function – for real

- ▶ The kernel of μ_e contains the Koszul syzygies Ksz_e generated by

$$gf_i \otimes f_j - gf_j \otimes f_i, \quad g \in S_{e-d}, \quad f_i, f_j \in I_d$$

- ▶ Expecting $\text{Ker } \mu_e = \text{Ksz}_e$, a first estimate of $\dim_{\mathbb{C}} \text{Ker } \mu_e$ is $h_S(e-d) \cdot \binom{h_I(d)}{2}$
- ▶ Expect the syzygies to also only have Koszul syzygies, correct by $h_S(e-2d) \cdot \binom{h_I(d)}{3}$
- ▶ And these also only have Koszul syzygies and ...
- ▶ This leads to the following estimate for $h_{S/I_{\langle d \rangle}}(t)$:

$$h_S(t) - \underbrace{h_S(t-d)h_I(d)}_{\text{gen's of } I_d} + \underbrace{h_S(t-2d)\binom{h_I(d)}{2}}_{\text{Koszul syzygies}} - \underbrace{h_S(t-3d)\binom{h_I(d)}{3}}_{\text{Koszul syzygy syzygies}} \pm \dots$$

- ▶ On the other hand, as soon as $h_{I_{\langle d \rangle}}(t_0) \geq h_I(t_0)$, then $I_t = (I_{\langle d \rangle})_t$ for $t \geq t_0$

The main conjecture

Expected syzygy conjecture (ESC)

$$h_{S/I_{\langle d \rangle}}(t) = \begin{cases} \sum_{k \geq 0} (-1)^k \cdot h_S(t - kd) \cdot \binom{h_I(d)}{k} & t < t_0, \\ r & t \geq t_0, \end{cases}$$

where t_0 is the least integer $> d$ such that the sum is at most r .

- ▷ This is always a lower bound due to Fröberg
- ▷ Alternative expression for the ideal:

$$h_{I_{\langle d \rangle}}(t) = \begin{cases} \sum_{k \geq 1} (-1)^{k-1} \cdot h_S(t - kd) \cdot \binom{h_I(d)}{k} & t < t_0, \\ h_I(t) & t \geq t_0, \end{cases}$$

Is the complicated alternating sum really needed?

- ▷ For \mathbb{P}^2 the (ESC) “actually” says $h_{I_{\langle d \rangle}}(t) = \min\{h_I(d) \cdot h_S(t - d), h_I(t)\}$
- ▷ This is no longer true in higher dimension – in general n summands are required
- ▷ **Smallest example:** 52 points in \mathbb{P}^3

$$h_{S/I_{\langle 5 \rangle}}(t) = \begin{cases} h_S(t) - 4h_S(t-5) + 6h_S(t-10) & t < 11, \\ 52 & t \geq 11 \end{cases}$$




Figure 2: The Hilbert function of the chopped ideal of 52 general points in \mathbb{P}^3 .

Theorem

Conjecture (ESC) is true in the following cases:

- ▷ $r_{\max} := h_S(d) - (n + 1)$ for all d in all dimensions n .
- ▷ In the plane for $r_{\min} = \frac{1}{2}(d + 1)^2$ when d is odd.
- ▷ $r \leq \frac{1}{n}((n + 1)h_S(d) - h_S(d + 1))$ and $[n \leq 4$ or generally whenever (IGC) holds].
- ▷ In a large number of individual cases in low dimension (next slide).

The length of the saturation gap is bounded above by

$$\min \{ e > 0 \mid (I_{\langle d \rangle})_{d+e} = I_{d+e} \} \leq (n - 1)d - (n + 1).$$

Whenever $I_{\langle d \rangle}$ is non-saturated, one has $\operatorname{reg}_{\text{CM}} S/I_{\langle d \rangle} = \operatorname{reg}_{\text{H}} S/I_{\langle d \rangle} - 1 = d + e - 1$.

Verification using computer algebra

- ▷ Testing the conjecture for particular values of (n, r) :
 - Sample r random points from $\mathbb{P}^n(\mathbb{Q})$
 - Calculate $h_{S/I(Z)_{\langle d \rangle}}(t)$ using a computer algebra system
 - If the sample satisfies (ESC), then the conjecture is true for general such Z

Theorem

The map $Z \mapsto h_{S/I(Z)_{\langle d \rangle}}(t)$ is upper semicontinuous on the set $U \subseteq (\mathbb{P}^n)^r$ of points with generic Hilbert function.

- ▷ To speed up computation, perform calculations over a finite field \mathbb{F}_p
- ▷ Using Macaulay2 we verified the conjecture in the following cases

n	2	3	4	5	6	7	8	9	10
r	≤ 1825	≤ 1534	≤ 991	≤ 600	≤ 447	≤ 316	≤ 333	≤ 204	≤ 259
d	≤ 58	≤ 18	≤ 9	≤ 6	≤ 4	≤ 3	≤ 3	≤ 2	≤ 2

Visualization of the saturation gaps in \mathbb{P}^2

- ▷ ESC predicts exactly how large the difference between I and $I_{\langle d \rangle}$ is




Figure 3: The saturation gaps for all values of $r \leq 102$ in \mathbb{P}^2 .

Visualization of the saturation gaps in \mathbb{P}^3







Figure 4: The saturation gaps for all values of $r \leq 116$ in \mathbb{P}^3 .

- ▷ Characteristic $p > 0$? \rightsquigarrow Should carry over.
- ▷ Proving the conjecture in \mathbb{P}^2 ?
- ▷ Improve code to verify more cases
- ▷ Generalizations multi-graded setting, e.g. points in $\mathbb{P}^n \times \mathbb{P}^m$
- ▷ State a conjecture for the minimal free resolution of $I(Z)_{\langle d \rangle}$

Thank you! Questions?

Preprint soon™

-  A. Bernardi, E. Carlini, M. V. Catalisano, A. Gimigliano, and A. Oneto.
The hitchhiker guide to: Secant varieties and tensor decomposition.
Mathematics, 6(12):314, 2018.
-  Ralf Fröberg.
An inequality for hilbert series of graded algebras.
Mathematica Scandinavica, 56(2), 1985.
-  Fulvio Gesmundo, Leonie Kayser, and Simon Telen.
Hilbert functions of chopped ideals.
In preparation, 2023.



Anna Lorenzini.

The minimal resolution conjecture.

Journal of Algebra, 156(1), 1993.



Simon Telen.

Solving Systems of Polynomial Equations.

PhD thesis, KU Leuven, Leuven, Belgium, 2020.



Simon Telen and Nick Vannieuwenhoven.

A normal form algorithm for tensor rank decomposition.

ACM Trans. on Math. Soft., 48(4):1–35, 2022.