Joint natural boundaries of the full square-lattice Ising susceptibility

Independent complex couplings and exact physical-slice descendants

21 October 2026

Abstract

We study the complete connected zero-field response of the infinite nearest-neighbor square-lattice Ising model with two independent complex couplings. We give explicit high- and low-temperature domains in $\mathbb{C}^2$, identify them as components of an independently specified spectral complement, construct the physical full-sum continuations there, and prove local nonextendability at every boundary point. At the selected regular boundary points the obstruction is stable along arbitrary nonramified holomorphic transverse curves, with genuine local uniformity in complex dispersion parameters. Its negative odd-particle edge mechanism uses exact pole periods and a single dominant index in selected high Taylor coefficients of the entire sum. The collinear isotropic case is treated by a separate, explicitly proved even–odd tail estimate. Exact pullbacks recover both prescribed fixed-positive-ratio temperature domains for every ratio, including all exceptional boundary points. The map $s = \sinh z$ then gives the full isotropic interior and exterior descendants with their distinct physical germs. The onsite term, all lattice sites, all particle indices, the plus-phase subtraction and the physical relation $\chi = \beta\mathcal{X}$ are retained. No conclusion about singularity throughout the interior of the spectral set is asserted.

1 Statements and the scope of the parent theorem

The variables K_x, K_y are independent coupling coordinates. Put

$$z = 2K_x = x + \mathrm{i}y, \qquad w = 2K_y = u + \mathrm{i}v,$$
$$S = \sinh z, \quad T = \sinh w, \quad A = \cosh z \cosh w, \quad p = ST.$$

For real positive couplings, $\mathcal{X}_H$ is the sum of all two-spin correlations in the unique high-temperature Gibbs state; $\mathcal{X}_L$ is the sum of all connected two-spin correlations in the infinite-volume plus phase. Their precise definitions and the physical interface are in Section 3.

Define independently of any susceptibility

$$\Sigma = \{(K_x, K_y) : A = Sa + Tb \text{ for some } a, b \in [-1, 1]\}.$$

The two domains used in this paper are explicitly

$$\mathcal{D}_H = \big\{|y| < \pi/2,\ |v| < \pi/2,\ \cosh x \tanh|u| < \cos y, \quad \cosh u \tanh|x| < \cos v\big\}, \tag{1.1}$$

$$\mathcal{D}_L = \big\{x > 0,\ u > 0, \quad \cosh x \tanh u > |\cos y|, \quad \cosh u \tanh x > |\cos v|\big\}. \tag{1.2}$$

These are open sets in the original coupling space, using the displayed invertible linear change of coordinates. The HT domain is unbounded along decoupled directions; boundedness is asserted only for its positive-ratio temperature slices.

Theorem 1.1 (Joint physical natural boundaries)**.** *The following assertions hold for both phases of the full bulk reduced susceptibility.*

1. *The sets $\mathcal{D}_H$ and $\mathcal{D}_L$ are precisely the components of $\mathbb{C}^2 \setminus \Sigma$ containing, respectively, the real positive regions $0 < p < 1$ and $p > 1$. They are contractible and pseudoconvex. On them $|p| < 1$ and $|p| > 1$, respectively.*

2. *The physical germs extend to single-valued jointly holomorphic functions $\mathcal{X}_H \in \mathcal{O}(\mathcal{D}_H)$ and $\mathcal{X}_L \in \mathcal{O}(\mathcal{D}_L)$. Their complete odd and connected-even form-factor sums converge normally on compact subsets, with every fixed mixed complex derivative.*

3. *At every point of the respective boundary, there is no holomorphic function on a neighborhood agreeing with that physical continuation on an accumulating connected component of its domain side. These are thus natural boundaries also for continuation in unramified Riemann domains over $\mathbb{C}^2$, reached from the given physical germ.*

4. *The proof gives more than an ambient obstruction. At a dense set on every regular boundary face, the complete physical sum is nonextendable along every holomorphic transverse curve on which the local edge parameter is nonramified. The estimates are locally uniform for a fixed selected phase and compact admissible subfamilies of bounded holomorphic complex-parameter families on common polydisks in a regular-edge chart. No uniformity over the primes or through $\Im B = 0$ is required. The degenerate equal-coupling restriction has the additional dense local obstructions of Proposition 6.8.*

Here regular faces and their edge parameters are specified by Definition 2.8 and Proposition 2.10. The final assertion is part of the parent construction needed for the exact restrictions below.

For every finite $\lambda > 0$, put $\iota_\lambda(\zeta) = (\zeta/2, \lambda\zeta/2)$. Let

$$P_\lambda(a) = \cosh a \tanh(\lambda a), \qquad Q_\lambda(a) = \cosh(\lambda a) \tanh a, \qquad Y_\lambda = \frac{\pi}{2\max(1,\lambda)}.$$

Both functions on $[0,\infty)$ have increasing inverses. Define

$$\begin{aligned} H_\lambda(b) &= \min\{P_\lambda^{-1}(\cos b), Q_\lambda^{-1}(\cos(\lambda b))\}, && |b| < Y_\lambda, \\ L_\lambda(b) &= \max\{P_\lambda^{-1}(|\cos b|), Q_\lambda^{-1}(|\cos(\lambda b)|)\}, && b \in \mathbb{R}, \\ \Omega_{H,\lambda} &= \{a + \mathrm{i}b : |b| < Y_\lambda,\ |a| < H_\lambda(b)\}, \\ \Omega_{L,\lambda} &= \{a + \mathrm{i}b : a > L_\lambda(b)\}. \end{aligned}$$

Theorem 1.2 (Exact recovery of every positive-ratio slice)**.** *For every finite positive λ,*

$$\iota_\lambda^{-1}(\mathcal{D}_H) = \Omega_{H,\lambda}, \qquad \iota_\lambda^{-1}(\mathcal{D}_L) = \Omega_{L,\lambda}.$$

The restrictions of the two joint physical functions are the respective physical reduced susceptibilities, holomorphic on these exact domains, and nonextendable through every boundary point in the componentwise sense of Theorem 1.1. They retain the complete lattice sum and every odd or nonzero even particle order. On a physical line with fixed $J_x > 0$, $J_y = \lambda J_x$, the same conclusions hold for the energy-field response

$$\chi(\zeta) = \frac{\zeta}{2J_x}\,\mathcal{X}(\iota_\lambda(\zeta)).$$

This is the complete statement of the supplied anisotropic Theorem 1.1, on precisely its domains. Its proof below uses the curve-level part of Theorem 1.1 and its analytic construction, rather than inferring a restriction obstruction solely from ambient nonextension.

Corollary 1.3 (Exact isotropic descendants)**.** *Set $\lambda = 1$ and $s = \sinh\zeta$. The reduced HT physical germ is single valued and holomorphic on $|s| < 1$, with value one at zero; the reduced connected-plus-phase LT physical germ is single valued and holomorphic on $|s| > 1$. Every point of $|s| = 1$ is a natural-boundary point for each germ. Their complete normalized representations are*

$$\mathcal{X}_H(s) = \frac{(1-s^4)^{1/4}}{s} \sum_{N\geq 1,\ N \text{ odd}} C_N(1/s), \qquad \mathcal{X}_L(s) = (1-s^{-4})^{1/4} \sum_{N\geq 2,\ N \text{ even}} C_N(s),$$

where the coefficients are the full-site integrals in Section 6. The LT onsite term is $1-(1-s^{-4})^{1/4}$. The map sinh *is a biholomorphism from $\Omega_{H,1}$ to the disk and is the universal covering from $\Omega_{L,1}$ to the exterior, with deck translations $\zeta \mapsto \zeta + 2\pi\mathrm{i}k$. The energy-field results follow with $\beta = \zeta/(2J)$ on that physical continuation. No single-valued inverse temperature on the whole exterior is required.*

1.1 Sources, genuine joint content, and proof dependencies

The starting geometry is related to the unsquared anisotropic Nickel equation studied by Boukraa, Hassani and Maillard [4, Section 6]. That paper distinguishes multi-variable finite-integral singularities from the full-susceptibility question. The present argument must supply, and below supplies, separate whole-sum obstructions. In particular the set Σ has interior; neither its interior nor every finite-integral candidate is declared a singularity of the physical sum.

The provided LT, HT and ANISO manuscripts [7, 8, 9] are mathematical source material. Their geometric identities, contour methods and candidate estimates have been reconstructed at the needed scope. None of their natural-boundary conclusions is an imported hypothesis. The real-temperature exact-solution interface used here is identified precisely in Section 3.1; all continuation and infinite-tail claims needed for the three theorems are proved in this manuscript.

The domain construction is genuinely joint. For example $z = \operatorname{arsinh}(1/2),\ w = \mathrm{i}\pi/6$ is an interior point of $\mathcal{D}_H$ and lies on no positive-real-ratio temperature line. The independent complex coordinates $(C, B) = (A/S, T/S)$ have nonzero Jacobian at every regular edge. Their common polydisks, constant-B transversals, and variable-B curve estimates are used explicitly. Conversely the exact identity $\mathcal{X}_H(K_x, 0) = e^{2K_x}$ shows why restriction nonextension needs its own proof.

The main dependency order is as follows. Independent spectral geometry gives the domains, regular faces, exact pullbacks and local collars. The physical interface and double Pfaffians give the joint full-sum holomorphic functions. Positive edges use a first singular derivative and a bounded whole tail. Negative odd edges use exact high Taylor coefficients and uniform index separation. The isotropic local lemmas supply the collinear case. Section 7 assembles these results into Theorems 1.1–1.2 and then derives Corollary 1.3. The quantitative dependency table and verification scope are recorded at the end.

Contents

1 Statements and the scope of the parent theorem **1**
1.1 Sources, genuine joint content, and proof dependencies 3

2 Independent complex spectral geometry **5**
2.1 The physical components and their branches 7
2.2 Exact fixed-ratio pullbacks 9
2.3 Regular boundary faces in two complex variables 10
2.4 Fixed-ratio regularity is a separate assertion 12
2.5 Exceptional boundary points and componentwise continuation 14
2.6 Pseudoconvexity and the Hartogs question 16

3 The physical observable and joint holomorphy **16**
3.1 The precise imported physical interface 17
3.2 Double contours and the full lattice 18
3.3 Energy-field normalization 20

4 Regular spectral edges: a genuinely complex-parameter theorem **20**

5 The full odd sum at a regular negative spectral edge **24**
5.1 Statement, branch, and the physical side 24
5.2 A uniform Gaussian numerator bound and all boundary jets 25
5.3 An exact reduction to simple-pole periods 27
5.4 Bounds for every winding and every particle index 28
5.5 An exact symmetric pole coordinate 29
5.6 The complete residue density and its integration cycle 33
5.7 A uniform beta evaluation of the first two windings 34
5.8 Exact finite differences and a single dominant index 35
5.9 Exhausting the remaining indices and windings 37
5.10 Nonextension and uniformity 39

6 The isotropic restriction: a unified even–odd proof **39**
6.1 Compact convergence and a dense arithmetic family 41
6.2 The first complete singular coefficient 42
6.3 A uniform boundary-tail theorem 44
6.4 Real-domain differentiation in the intermediate window 48
6.5 Dense local obstructions and the exact physical identities 53

7 Closure of the joint theorem and its exact descendants **54**
7.1 The complete joint obstruction 54
7.2 All physical ratios, including the collinear one 56
7.3 Derivation of the complete isotropic theorems from the slice theorem 57

8 Dependency map and verification scope **57**

2 Independent complex spectral geometry

This section concerns the explicitly defined spectral set, not the singularities of a susceptibility. In particular, no assertion about a complete particle sum is used in its proofs. Write

$$\begin{aligned} z = 2K_x = x + \mathrm{i}y, \qquad w = 2K_y = u + \mathrm{i}v, \\ S = \sinh z, \quad T = \sinh w, \quad A = \cosh z \cosh w, \quad p = ST, \end{aligned}$$

and define the closed set

$$\Sigma = \{(K_x, K_y) : A = Sa + Tb \text{ for some } (a, b) \in [-1, 1]^2\}. \tag{2.1}$$

The two real functions that describe this set are

$$F_1 = \cosh^2 u \cos^2 y - \cosh^2 x \sinh^2 u, \qquad F_2 = \cosh^2 x \cos^2 v - \cosh^2 u \sinh^2 x. \tag{2.2}$$

The use of z, w instead of K_x, K_y is just an invertible linear coordinate change; all domains below are domains in the original $\mathbb{C}^2$ coupling space.

Theorem 2.1 (Exact spectral set)**.** *For two independent complex couplings, including zero, opposite-sign, and collinear real-part configurations,*

$$\Sigma = \{F_1 F_2 \le 0\}. \tag{2.3}$$

Proof. The independent reflections $z \mapsto -z$ and $w \mapsto -w$ preserve Σ, by changing the sign of the corresponding representing coefficient. They also preserve F_1, F_2. We may therefore first suppose $x, u > 0$. Set

$$\begin{aligned} h = \sinh x, \quad c = \cosh x, \quad H = \sinh u, \quad C_0 = \cosh u, \\ t = \cos y, \quad q = \sin y, \quad V = \cos v, \quad Q = \sin v. \end{aligned}$$

For

$$D = \Im(\overline{S}T), \qquad N_a = \Im(\overline{A}T), \qquad N_b = \Im(\overline{S}A),$$

direct multiplication gives

$$\begin{aligned} D &= hC_0tQ - cHqV, \\ N_a &= ctVQ - hHC_0q, \qquad N_b = hcHQ - C_0tqV. \end{aligned} \tag{2.4}$$

When $D \neq 0$, the unique real solution of $A = Sa + Tb$ is $a = N_a/D,\ b = N_b/D$.

For the sign analysis put

$$\begin{aligned} a_0 = h^2, \quad b_0 = H^2, \quad A_0 = 1 + a_0, \quad B_0 = 1 + b_0, \\ X = t^2, \quad Y = V^2, \quad X_0 = \frac{A_0 b_0}{B_0}, \quad Y_0 = \frac{B_0 a_0}{A_0}. \end{aligned}$$

Expanding (2.4) and cancelling the common cross terms gives the identities

$$\begin{aligned} F_1 &= B_0(X - X_0), & F_2 &= A_0(Y - Y_0), \\ N_a^2 - D^2 &= F_2 K, & K &= (B_0 - Y)X - b_0, \\ N_b^2 - D^2 &= -F_1 L, & L &= a_0 - (A_0 - X)Y. \end{aligned} \tag{2.5}$$

Both K and L vanish at (X_0, Y_0), increase with X, and decrease with Y. If F_1, F_2 have strictly opposite signs, both differences of squares in (2.5) are strictly negative. Consequently $D \neq 0$ and $|a|, |b| < 1$; the point belongs to Σ.

Suppose instead that F_1, F_2 have the same strict sign and $D \neq 0$. For $0 < Y < 1$, membership would impose

$$X \leq \frac{b_0}{B_0 - Y}, \quad X \geq A_0 - \frac{a_0}{Y} \quad \text{if } F_1, F_2 > 0,$$

and the reverse two inequalities if $F_1, F_2 < 0$. But

$$\frac{b_0}{B_0 - Y} - \left(A_0 - \frac{a_0}{Y}\right) = \frac{A_0(Y-1)(Y-Y_0)}{Y(B_0 - Y)}. \tag{2.6}$$

For the positive-sign case $Y > Y_0$, this expression is negative; for the negative-sign case $Y < Y_0$, it is positive. Each case contradicts the required inequalities. At $Y = 0$, $F_2 < 0$ and $L = a_0 > 0$, so the negative-sign case is impossible directly from (2.5). At $Y = 1$, $K = b_0(X - 1)$ and $L = X - 1$; either same-sign case requires $X = 1$, which gives $D = 0$.

It remains to exclude a same-strict-sign point with $D = 0$ from Σ. Membership would give $N_a = N_b = 0$. Since $F_1 F_2 \neq 0$, (2.5) implies $K = L = 0$. Here $Y > 0$, and (2.6) then implies $Y = Y_0$ or $Y = 1$. The first contradicts $F_2 \neq 0$, whereas the second gives $X = 1$. At $X = Y = 1$ the segment condition is

$$cC_0 \leq h + H.$$

As

$$(cC_0)^2 - (h + H)^2 = (hH - 1)^2,$$

this requires $hH = 1$, and hence $F_1 = F_2 = 0$, again a contradiction.

The equality cases have a particularly useful exact proof. At a point with $F_1 = 0$, $S \neq 0$, and $\tanh(z/2)$ is finite and nonzero. The two endpoint roots

$$r_+ = e^w \tanh(z/2), \qquad r_- = e^{-w} \tanh(z/2) \tag{2.7}$$

satisfy

$$\frac{r_+ + r_+^{-1}}{2} = \frac{A - T}{S}, \qquad \frac{r_- + r_-^{-1}}{2} = \frac{A + T}{S}. \tag{2.8}$$

Moreover,

$$|\tanh(z/2)|^2 = \frac{\cosh x - \cos y}{\cosh x + \cos y}. \tag{2.9}$$

Thus $F_1 = 0$ means that at least one of r_+, r_- has modulus one: the respective equations are $\cos y = \cosh x \tanh u$ and $\cos y = -\cosh x \tanh u$. The corresponding right side of (2.8) belongs to $[-1, 1]$. This supplies a representing pair with $b = 1$ or $b = -1$. Interchanging z, w proves membership when $F_2 = 0$.

For completeness, the zero-real-part cases can be checked without any determinant division. Let $x = 0, u > 0$. If $\cos v \neq 0$, the real part of the equation $A = Sa + Tb$ forces

$$b = \frac{\cosh u}{\sinh u} \cos y.$$

Its necessary condition is $|\cos y| \leq \tanh u$. Under that condition $\sin y \neq 0$, and

$$a = -\frac{\cos y \sin v}{\sinh u \sin y}$$

has modulus at most one, so the condition is also sufficient. Indeed $\cos^2 y \le \sinh^2 u/\cosh^2 u \le \sinh^2 u/(\sinh^2 u + \sin^2 v)$. Here $F_2 = \cos^2 v > 0$, and the condition is precisely $F_1 \le 0$. If $\cos v = 0$, all terms are imaginary, and the choice

$$a = 0, \qquad b = \tanh u \cos y$$

always represents A; here $F_2 = 0$. The case $u = 0, x > 0$ follows by exchange. Finally, if $x = u = 0$, the coefficients S, T are imaginary and $A = \cos y \cos v$ is real. Membership is equivalent to $A = 0$, represented by $a = b = 0$, and is exactly $F_1 F_2 = 0$. The initial reflections now cover all real-part signs. □

2.1 The physical components and their branches

Theorem 2.2 (All components; explicit physical domains)**.** *The components of $\mathbb{C}^2 \setminus \Sigma$ with $F_1, F_2 > 0$ are indexed by $m, n \in \mathbb{Z}$ and are*

$$F_1 > 0, \quad F_2 > 0, \quad |y - m\pi| < \pi/2, \quad |v - n\pi| < \pi/2.$$

Each is star-shaped about the point $(z, w) = (\mathrm{i}m\pi, \mathrm{i}n\pi)$. The components with $F_1, F_2 < 0$ are the four sets specified by the independent signs of x, u; each is contractible.

The component containing the origin is exactly

$$\mathcal{D}_H = \left\{ \begin{array}{l} |y| < \pi/2, \quad |v| < \pi/2, \\ \cosh x \tanh |u| < \cos y, \quad \cosh u \tanh |x| < \cos v \end{array} \right\}. \tag{2.10}$$

The component containing the positive real low-temperature region is

$$\mathcal{D}_L = \left\{ \begin{array}{l} x > 0, \quad u > 0, \\ \cosh x \tanh u > |\cos y|, \quad \cosh u \tanh x > |\cos v| \end{array} \right\}. \tag{2.11}$$

They satisfy the strict inequalities

$$|p| < 1 \text{ on } \mathcal{D}_H, \qquad |p| > 1 \text{ on } \mathcal{D}_L. \tag{2.12}$$

Proof. The sign description follows from Theorem 2.1. For $F_1, F_2 > 0$, both cosines are nonzero, so a connected component remains in a single indicated cosine strip in each variable. Within a fixed pair of strips, set $\eta = y - m\pi,\ \theta = v - n\pi$. The inequalities are

$$\cosh x \tanh |u| < \cos \eta, \qquad \cosh u \tanh |x| < \cos \theta.$$

Replacing (x, u, η, θ) by $t(x, u, \eta, \theta)$, $0 \le t \le 1$, decreases the left sides and increases the right sides. This proves star-shapedness and excludes further component subdivision.

For $F_1, F_2 < 0$, neither x nor u can vanish. Writing $a = |x|,\ b = |u|$, the conditions are

$$\cosh a \tanh b > |\cos y|, \qquad \cosh b \tanh a > |\cos v|.$$

Each left side is nondecreasing in both a, b, and strictly increasing in the variable inside its hyperbolic tangent. Choose once and for all $R > 0$ with $\sinh R > 1$. First increase (a, b) continuously to $(\max(a, R), \max(b, R))$. Then move these two coordinates linearly to (R, R), throughout staying in $[R, \infty)^2$. During the second move both left sides are at least $\cosh R \tanh R = \sinh R > 1$. Finally move y, v linearly to zero. These three moves give a continuous contraction within the fixed sign

component. The signs cannot change without leaving the set. This proves all component claims and the physical identifications (2.10)–(2.11).

The modulus identity, with the notation in the proof of Theorem 2.1, is

$$|p|^2 = (A_0 - X)(B_0 - Y). \tag{2.13}$$

For $F_1, F_2 > 0$,

$$A_0 - X < \frac{A_0}{B_0}, \qquad B_0 - Y < \frac{B_0}{A_0},$$

so $|p| < 1$. For $F_1, F_2 < 0$, both inequalities reverse, so $|p| > 1$. These arguments allow zero real parts as well. On the positive real quadrant the common sign of F_1, F_2 is the sign of $1 - (\sinh x \sinh u)^2$, which proves the claimed physical identification. □

Remark 2.3. In particular $\{|p| = 1\} \subset \Sigma$. The more specific corner identity is

$$A^2 - (\epsilon S + \delta T)^2 = (p - \epsilon\delta)^2, \qquad \epsilon, \delta \in \{1, -1\},$$

and therefore

$$\bigcup_{\epsilon,\delta\in\{1,-1\}} \{A = \epsilon S + \delta T\} = \{p^2 = 1\} \subset \Sigma.$$

The branches $(1 - p^2)^{1/4}$ on $\mathcal{D}_H$ and $(1 - p^{-2})^{1/4}$ on $\mathcal{D}_L$ are defined directly by the binomial series and their positive real physical values. No additional covering is needed for these factors. On $\mathcal{D}_L$, S, T are nonzero. On $\mathcal{D}_H$ the zeros of S and T are respectively $z = 0$ and $w = 0$; these are genuine complex subvarieties of the joint domain. Any particle representation used there must account for its removable normalization at these axes.

Lemma 2.4 (A holomorphic modulus description)**.** *Within the two central strips in* (2.10)*, the four inequalities defining $\mathcal{D}_H$ can equivalently be written*

$$|e^w \tanh(z/2)| < 1, \quad |e^{-w} \tanh(z/2)| < 1, \quad |e^z \tanh(w/2)| < 1, \quad |e^{-z} \tanh(w/2)| < 1.$$

The domain $\mathcal{D}_L$ is equivalently described by

$$\begin{aligned} |e^{-w} \tanh(z/2)| < 1, \quad |e^{-w} \coth(z/2)| < 1, \\ |e^{-z} \tanh(w/2)| < 1, \quad |e^{-z} \coth(w/2)| < 1. \end{aligned}$$

The functions in the second display are evaluated where they are finite; its inequalities themselves force $x, u > 0$.

Proof. For the first pair in the HT assertion, (2.9) gives

$$e^{2|u|} \frac{\cosh x - \cos y}{\cosh x + \cos y} < 1,$$

equivalent to $\cos y > \cosh x \tanh |u|$. Exchange variables for the other pair. In the first LT pair, multiplication forces $e^{-2u} < 1$, hence $u > 0$; the second pair similarly forces $x > 0$. If $r = |\tanh(z/2)|^2$, the first pair is $e^{-2u} < r < e^{2u}$, which is equivalent to $|\cos y| < \cosh x \tanh u$. Exchange variables again. □

Lemma 2.5 (Canonical decaying root)**.** *For $K = (K_x, K_y) \notin \Sigma$ and real q, the equation*

$$S\rho^2 - 2(A - T\cos q)\rho + S = 0 \tag{2.14}$$

has exactly one root of modulus less than one, interpreting that root as zero when $S = 0$. This root is holomorphic in the independent coupling variables and locally jointly holomorphic with the complex angle q. On a compact subset of $\mathbb{C}^2 \setminus \Sigma$, its modulus has a uniform gap below one for all real q. On both positive real physical components it is the positive decaying root.

Proof. When $S \neq 0$, put $W = (A - T\cos q)/S$. By the definition of Σ, $W \notin [-1, 1]$. The two roots are reciprocal, neither has modulus one, and exactly one lies in the unit disk. Equivalently, it is $\rho = W - \sqrt{W^2 - 1}$, with the square-root branch asymptotic to W at infinity. Its uniqueness makes these local holomorphic roots agree on overlaps. At $S = 0$, $A - T\cos q \neq 0$; the implicit function theorem at $\rho = 0$ supplies the same holomorphic root. Compactness gives both the modulus gap and a common thin complex angular neighborhood on any compact parameter set. For positive real S, T,

$$(A - S - T)(A + S + T) = (p - 1)^2 > 0$$

on either physical component, and hence $W > 1$. The interior root is then real and belongs to $(0, 1)$. □

2.2 Exact fixed-ratio pullbacks

Theorem 2.6 (Exact slice domains)**.** *For every finite $\lambda > 0$, let*

$$\iota_\lambda(\zeta) = (\zeta/2, \lambda\zeta/2), \qquad Y_\lambda = \frac{\pi}{2\max(1, \lambda)},$$

and, for $a \geq 0$, let

$$P_\lambda(a) = \cosh a \, \tanh(\lambda a), \qquad Q_\lambda(a) = \cosh(\lambda a) \, \tanh a.$$

Both functions are strictly increasing bijections of $[0, \infty)$ onto itself. Define

$$\begin{aligned} H_\lambda(b) &= \min\{P_\lambda^{-1}(\cos b),\ Q_\lambda^{-1}(\cos(\lambda b))\}, && |b| < Y_\lambda, \\ L_\lambda(b) &= \max\{P_\lambda^{-1}(|\cos b|),\ Q_\lambda^{-1}(|\cos(\lambda b)|)\}, && b \in \mathbb{R}. \end{aligned}$$

Then the exact equalities of open sets are

$$\iota_\lambda^{-1}(\mathcal{D}_H) = \Omega_{H,\lambda} = \{a + \mathrm{i}b : |b| < Y_\lambda,\ |a| < H_\lambda(b)\}, \tag{2.15}$$

$$\iota_\lambda^{-1}(\mathcal{D}_L) = \Omega_{L,\lambda} = \{a + \mathrm{i}b : a > L_\lambda(b)\}. \tag{2.16}$$

Both domains are connected and simply connected. There is a unique $z_c > 0$ with

$$\sinh z_c \sinh(\lambda z_c) = 1, \qquad P_\lambda(z_c) = Q_\lambda(z_c) = 1.$$

The function H_λ is positive inside its interval and extends continuously by zero at both endpoints, and $0 \leq L_\lambda \leq z_c$.

Proof. The derivatives

$$P'_\lambda(a) = \sinh a \tanh(\lambda a) + \lambda \cosh a \, \operatorname{sech}^2(\lambda a), \quad Q'_\lambda(a) = \lambda \sinh(\lambda a) \tanh a + \cosh(\lambda a) \operatorname{sech}^2 a$$

are positive on $[0, \infty)$, and both functions tend to infinity. On the slice, $(x, y, u, v) = (a, b, \lambda a, \lambda b)$. The two strip conditions in (2.10) are exactly $|b| < Y_\lambda$; its remaining two inequalities are $P_\lambda(|a|) < \cos b$ and $Q_\lambda(|a|) < \cos(\lambda b)$. This proves (2.15). In (2.11) the positive-real-part conditions are exactly $a > 0$, and the other inequalities invert to (2.16).

The positive function $\sinh a \sinh(\lambda a)$ increases strictly from zero to infinity. The identities

$$P_\lambda(a)^2 - 1 = \frac{\sinh^2 a \sinh^2(\lambda a) - 1}{\cosh^2(\lambda a)}, \quad Q_\lambda(a)^2 - 1 = \frac{\sinh^2 a \sinh^2(\lambda a) - 1}{\cosh^2 a}$$

prove the assertions about z_c and L_λ. At either endpoint of the HT interval at least one of its two cosine arguments is zero, which proves the endpoint limit for H_λ. The HT domain is star-shaped by Theorem 2.2, or directly by simultaneous scaling of a, b. The LT domain is homeomorphic to $(0, \infty) \times \mathbb{R}$ via $(a, b) \mapsto (a - L_\lambda(b), b)$. □

Lemma 2.7 (Isotropic covering geometry)**.** *At $\lambda = 1$,*

$$\iota_1^{-1}(\Sigma) = \{\zeta : |\sinh \zeta| = 1\},$$

$$\Omega_{H,1} = \{|b| < \pi/2, \ |a| < \operatorname{arsinh}(\cos b)\}, \qquad \Omega_{L,1} = \{a > \operatorname{arsinh}(|\cos b|)\}.$$

The map $s = \sinh \zeta$ maps $\Omega_{H,1}$ biholomorphically onto $\{|s| < 1\}$, and maps $\Omega_{L,1}$ as the universal covering of $\{|s| > 1\}$, with deck translations $\zeta \mapsto \zeta + 2\pi \mathrm{i} k$, $k \in \mathbb{Z}$.

Proof. Here $S = T = s$, $A = 1 + s^2$. For $s \neq 0$, the spectral condition is $s + s^{-1} \in [-2, 2]$, whose preimage is exactly the unit circle; $s = 0$ is not spectral. Also $P_1 = Q_1 = \sinh$, giving the displayed domains. On the central strip $|\Im \zeta| < \pi/2$, sinh is injective: the alternatives for equal hyperbolic sines are $\zeta_1 = \zeta_2 + 2\pi \mathrm{i} k$ or $\zeta_1 = -\zeta_2 + (2k + 1)\pi \mathrm{i}$, and only the first with $k = 0$ is possible there. The domain $\Omega_{H,1}$ is bounded, its boundary maps into $|s| = 1$, and its derivative has no zero inside. Its image is consequently both open and relatively closed in the disk, and contains zero; it is the whole disk.

For the exterior, put $\omega = e^\zeta$. The equation $\omega - \omega^{-1} = 2s$ has, for $|s| > 1$, exactly one root with $|\omega| > 1$. A root of modulus one would give $s \in \mathrm{i}[-1, 1]$. The outside root varies holomorphically, for example by using $\sqrt{s^2 + 1} = s\sqrt{1 + s^{-2}}$. Its domain is the continuous radial epigraph

$$\{ re^{\mathrm{i}b} : r > \exp(\operatorname{arsinh} |\cos b|) \}.$$

Exponentiation from the simply connected epigraph $\Omega_{L,1}$ is its universal covering, and the outside-root map is a biholomorphism to $|s| > 1$. The stated deck group follows. □

2.3 Regular boundary faces in two complex variables

Definition 2.8. A point of $\partial \mathcal{D}_H$ or $\partial \mathcal{D}_L$ will be called a regular joint edge point if $S, T \neq 0$, $D = \Im(\overline{S}T) \neq 0$, exactly one of its unique representing coefficients a, b is 1 or -1, the other lies in $(-1, 1)$, and $p^2 \neq 1$.

Proposition 2.9 (Density including exceptional strata)**.** *Regular joint edge points are dense in each of $\partial \mathcal{D}_H$ and $\partial \mathcal{D}_L$. More precisely, the points on either boundary for which $x, u \neq 0$ and exactly one of F_1, F_2 vanishes are regular and are dense. Thus the excluded zero-real-part, collinear, and intersection strata contain no relatively open boundary patch.*

Proof. First suppose $x, u \neq 0$, $F_1 = 0$, $F_2 \neq 0$. The algebra (2.5), valid also for negative real parts because it only uses squares, reduces to

$$N_a^2 - D^2 = -\frac{\sinh^2 u}{\cosh^2 u} F_2^2 < 0, \qquad N_b^2 - D^2 = 0. \tag{2.17}$$

Thus $D \neq 0$, $|a| < 1$, and $|b| = 1$. Furthermore (2.13) gives

$$|p|^2 = 1 - \frac{F_2}{\cosh^2 u} \quad (F_1 = 0), \qquad |p|^2 = 1 - \frac{F_1}{\cosh^2 x} \quad (F_2 = 0). \tag{2.18}$$

At an HT face the other F_i is positive, and at an LT face it is negative. Consequently $p^2 \neq 1$. The case $F_2 = 0, F_1 \neq 0$ follows by exchange.

Here is an explicit approximation argument covering all remaining boundary configurations. For $a, b \geq 0$, set

$$r_1(a, b) = \cosh a \tanh b, \qquad r_2(a, b) = \cosh b \tanh a.$$

Both $r_i < 1$, both equal one, or both exceed one according as $\sinh a \sinh b < 1$, $= 1$, or > 1. The closure of $\mathcal{D}_H$ is

$$|y|, |v| \leq \pi/2, \qquad r_1(|x|, |u|) \leq \cos y, \qquad r_2(|x|, |u|) \leq \cos v. \tag{2.19}$$

Indeed, contracting all four coordinates toward zero strictly opens every active inequality, proving the reverse inclusion in this closure description. If $\sinh |x| \sinh |u| < 1$, the angular fiber of (2.19) is the closed rectangle

$$|y| \leq \arccos r_1, \qquad |v| \leq \arccos r_2.$$

If the product is one, both intervals collapse to zero. Every boundary point has at least one active rectangular face. Approximate its real-part magnitudes by $a_j, b_j > 0$ with $\sinh a_j \sinh b_j < 1$. For an active first face choose $|y_j| = \arccos r_1(a_j, b_j)$, with the sign giving $y_j \to y$, and choose v_j strictly inside the second interval with $v_j \to v$. If the first interval has collapsed at the limit, either sign is available. Preserve the original signs of nonzero real parts; choose either sign at a zero real part. This gives $x_j, u_j \neq 0$, $F_1 = 0, F_2 > 0$, tending to the given point. Use the other face if it is the active one. This also covers corners, strip endpoints, and the collapsed fiber at real product one.

For $\mathcal{D}_L$, the closure is exactly

$$x, u \geq 0, \qquad r_1(x, u) \geq |\cos y|, \qquad r_2(x, u) \geq |\cos v|. \tag{2.20}$$

Increasing both real parts by any positive amount opens all its inequalities and proves the closure assertion. A boundary point must have $\sinh x \sinh u \leq 1$: if the product exceeds one, both $r_i > 1$ and the point is interior. It must also have at least one active cosine inequality. For example, $x = 0$ forces the second inequality to be active, while $u = 0$ forces the first. Again choose $x_j, u_j > 0$ with product less than one, tending to x, u. At an active first face choose

$$|\cos y_j| = r_1(x_j, u_j), \quad y_j \to y, \qquad |\cos v_j| < r_2(x_j, u_j), \quad v_j \to v.$$

These choices are always possible in the appropriate local cosine intervals. When $r_i \to 0$, these intervals shrink to a cosine zero; when $r_i \to 1$, their endpoints approach a multiple of π. The strict choice in the second interval remains possible since $r_2(x_j, u_j) > 0$. This gives $F_1 = 0, F_2 < 0$, tending to the prescribed point. The other active face is treated by exchange. □

Proposition 2.10 (Holomorphic coordinates on a regular face)**.** *Where $S \neq 0$, set*

$$C = A/S, \qquad B = T/S.$$

Then

$$\det \frac{\partial(C,B)}{\partial(z,w)} = \frac{p^2-1}{S^3}. \tag{2.21}$$

At a regular joint edge, (C,B) are therefore local holomorphic coordinates and $\Im B \neq 0$. If the endpoint is $b = \epsilon$, use $h = C - \epsilon B$: locally the boundary is $\Im h = 0$, with $\Re h \in (-1,1)$, and the physical domain occupies exactly one of its two sides. If the endpoint is $a = \epsilon$, exchange the transfer axes for the same conclusion.

On each regular boundary patch, points at which its free coefficient is

$$\cos(2\pi k/q), \qquad q \text{ an odd prime}, \quad 1 \le k \le (q-1)/2,$$

are dense. This assertion specifies a dense geometric family only; it does not assert singularity of a sum there.

Proof. The four derivatives are

$$C_z = -\frac{\cosh w}{S^2}, \quad C_w = \frac{\cosh z\, T}{S}, \quad B_z = -\frac{T\cosh z}{S^2}, \quad B_w = \frac{\cosh w}{S},$$

which give (2.21). Also $D = |S|^2 \Im B$. In these coordinates the spectral condition is simply

$$C \in [-1,1] + B[-1,1].$$

For nonreal B, this is a closed parallelogram in the C-plane. Near an interior point of its edge with $b = \epsilon$, the condition for the corresponding representing coefficient is

$$b - \epsilon = \frac{\Im(C - \epsilon B)}{\Im B};$$

the other coefficient remains strictly between its endpoints. Hence one side is spectral and the other is the unique local complementary side. Its intersection with the physical component is that whole side after shrinking the neighborhood. Holding B fixed permits h to range over an open real interval on the edge. The stated prime cosine values are dense in $(-1,1)$, since for arbitrarily large odd primes the angles $2\pi k/q$ approximate every angle in $(0,\pi)$. The coordinate description then proves the density assertion. □

Remark 2.11. On an HT regular face $F_1 = 0$, the endpoint is $b = \operatorname{sign} u$; on an HT regular face $F_2 = 0$, it is $a = \operatorname{sign} x$. This follows from (2.8), since both cosines are positive in the physical strips. On an LT regular face $F_1 = 0$, the endpoint is $b = \operatorname{sign}(\cos y)$; on $F_2 = 0$, it is $a = \operatorname{sign}(\cos v)$. In particular both signs of endpoint must be treated in any proof for the full physical boundaries.

2.4 Fixed-ratio regularity is a separate assertion

Joint boundary density alone need not restrict to a specified complex line. The following result supplies the separate one-variable statement for every $\lambda \neq 1$.

Proposition 2.12 (Every anisotropic slice has dense regular arcs)**.** *Fix $\lambda > 0, \lambda \neq 1$, and restrict S, T, A, p, D to $(z, w) = (\zeta, \lambda\zeta)$. On each boundary $\partial\Omega_{H,\lambda}$ and $\partial\Omega_{L,\lambda}$, all but a locally finite set of points have $D \neq 0$, exactly one endpoint coefficient, and a nonzero derivative of the corresponding meromorphic edge map*

$$h_\epsilon(\zeta) = \frac{A - \epsilon S}{T} \quad (a = \epsilon), \qquad g_\epsilon(\zeta) = \frac{A - \epsilon T}{S} \quad (b = \epsilon). \tag{2.22}$$

Such points lie on real analytic boundary arcs, and are dense on both full boundaries. On each such arc, the prime cosine values in Proposition 2.10 are dense in its free coefficient.

Proof. We first prove the required local finiteness of the collinear candidates. Off the imaginary axis, both real parts $x, \lambda x$ are nonzero. The strict opposite-sign case in (2.5) gives $D \neq 0$; the strict face identity (2.17) excludes $D = 0$ on a single zero face as well. Consequently a collinear candidate off the imaginary axis has $F_1 = F_2 = 0$, hence $|p| = 1$. Conversely $|p| = 1$ implies membership in Σ, by (2.12) applied to all complementary components. Thus

$$\{\, \zeta \in \iota_\lambda^{-1}(\Sigma) : \Re\zeta \neq 0,\ D = 0 \,\} = \{\, \zeta : \Re\zeta \neq 0,\ D = 0,\ |p| = 1 \,\}. \tag{2.23}$$

For $x > 0$, the modulus equation is

$$(\sinh^2 x + \sin^2 y)(\sinh^2(\lambda x) + \sin^2(\lambda y)) = 1. \tag{2.24}$$

If $|\sin y \sin(\lambda y)| < 1$, its left side starts below one and strictly increases to infinity with x. Its positive x derivative at the root gives a unique positive real analytic solution $x = X(y)$. At a simultaneous zero of $\cos y, \cos(\lambda y)$, the root tends continuously to zero.

If there are no simultaneous cosine zeros, X is analytic on all of $\mathbb{R}$. Since X is even near zero, $X'(0) = 0$, and $X(0) = z_c$. Direct differentiation therefore gives

$$\frac{\mathrm{d}}{\mathrm{d}y} D(X(y) + \mathrm{i}y)\Big|_{y=0} = \lambda \coth(\lambda z_c) - \coth z_c \neq 0. \tag{2.25}$$

For the last inequality, $t \coth t$ is strictly increasing on $t > 0$, because its derivative has the positive numerator $\sinh t \cosh t - t$. The analytic function $D(X(y) + \mathrm{i}y)$ is therefore nonzero identically and has locally isolated zeros.

Simultaneous cosine zeros occur exactly when $\lambda = r/s$ in lowest positive terms with r, s both odd; their locations are $y_j = (2j+1)\pi s/2$. Both X and $D(X(y) + \mathrm{i}y)$ have period πs. Each interval between successive y_j's is a translate of the interval containing zero, so (2.25) proves nonidentity on every such interval. It remains to exclude accumulation at its endpoints. Put $\zeta = \mathrm{i}y_j + \eta$, $\eta = x + \mathrm{i}\delta$. For signs $\sigma, \tau \in \{1, -1\}$,

$$S = \mathrm{i}\sigma \cosh \eta, \qquad T = \mathrm{i}\tau \cosh(\lambda\eta),$$

and direct Taylor expansion gives

$$D = \sigma\tau(\lambda^2 - 1)x\delta + O(|\eta|^4). \tag{2.26}$$

Equation (2.24), written analytically in x^2, δ^2, has linear part $(1 + \lambda^2)(x^2 - \delta^2)$. The implicit function theorem gives

$$X(y_j + \delta) = |\delta| + O(|\delta|^3).$$

Inserting this into (2.26) shows $D \neq 0$ for all sufficiently small nonzero δ, since $\lambda \neq 1$. There is no endpoint accumulation.

On the imaginary axis the candidate condition is $\cos y \cos(\lambda y) = 0$, a locally finite set. At a point where the cosine zeros are not simultaneous, $|p| = |\sin y \sin(\lambda y)| < 1$; continuity and (2.23) exclude nearby off-axis collinear candidates. The simultaneous case was just handled. Reflection $x \mapsto -x$, leaving y fixed, supplies the same conclusion on the other side. The complete set of collinear candidates is consequently locally finite in the ζ-plane.

The remaining exceptions are also locally finite. The zeros of S, T, and of $p^2 - 1$, are discrete. Each of the four functions in (2.22) is nonconstant meromorphic: at zero its leading term is $1/(\lambda\zeta)$ or $1/\zeta$. Its poles and critical points are locally finite. Let E_λ be the union of these sets and the collinear candidates.

At a boundary point outside E_λ, the unique representing coefficients cannot both be strict: otherwise continuity would place a whole neighborhood inside the spectral set, contrary to being approached by the complementary domain. They cannot both be endpoints, by the corner identity in Remark 2.3. Exactly one is an endpoint. Its edge map has nonzero derivative. For example, at $a = \epsilon$,

$$\Im h_\epsilon = \Im(S/T)(a - \epsilon)$$

for the unique real representing coefficients near the point; $\Im(S/T) \neq 0$. Hence the spectral set is locally one closed side of the nonsingular real analytic arc $\Im h_\epsilon = 0$, and the physical domain occupies the other side.

Neither physical boundary has an isolated point: the HT boundary is the two continuous graphs $a = \pm H_\lambda(b)$, completed at their endpoints, and the LT boundary is the continuous graph $a = L_\lambda(b)$. Removing the locally finite set E_λ therefore leaves a dense subset of both boundaries. The edge map is a local biholomorphism and maps each small regular arc to a real interval. Prime cosine density now proves the final assertion. □

Remark 2.13. The regular right boundary of the HT lens has positive endpoint, and its regular left boundary has negative endpoint, since x and $u = \lambda x$ have the same sign. On the LT boundary the endpoint signs are those in Remark 2.11, with $v = \lambda y$. Thus the negative odd endpoint problem occurs on the actual HT physical boundary for every anisotropic positive ratio; it is not removed by selecting the physical component. The ratio $\lambda = 1$ is deliberately excluded from Proposition 2.12; its collinear degeneracy is the entire isotropic boundary, described instead by Lemma 2.7.

2.5 Exceptional boundary points and componentwise continuation

Lemma 2.14 (Uniform local collar)**.** *Let $D = \mathcal{D}_H$ or $\mathcal{D}_L$, and let $P \in \partial D$. For every open neighborhood U of P, there is an open neighborhood $V \subset U$ of P such that $D \cap V$ is contained in a single connected component of $D \cap U$. Consequently every component E of $D \cap U$ with $P \in \overline{E}$ contains $D \cap V$. The same assertion holds for every $\Omega_{H,\lambda}$ and $\Omega_{L,\lambda}$.*

Proof. It suffices to work in a small ball contained in U. Use real coordinates (x, u, y, v), and first consider $\mathcal{D}_H$. Its two nontrivial defining functions in the central strips are

$$g_1 = \cos y - \cosh x \tanh|u|, \qquad g_2 = \cos v - \cosh u \tanh|x|.$$

At $P = (x_0, u_0, y_0, v_0)$, use the fixed direction $\nu = -P$. For an active inequality $g_1(P) = 0$, its derivative in that direction is

$$y_0 \sin y_0 + x_0 \sinh x_0 \tanh|u_0| + |u_0| \cosh x_0 \operatorname{sech}^2 u_0 > 0. \tag{2.27}$$

All terms are nonnegative in the closed central strips. If the last term vanishes then $u_0 = 0$, and activity forces $|y_0| = \pi/2$, making the first term positive. If $u_0 \neq 0$, the last term is positive.

Although $|u|$ is not differentiable at zero, when $u_0 = 0$ the chosen direction has zero u component. The directional derivative along ν is then well-defined and continuous in a neighborhood of P. The exchanged calculation applies to an active g_2. Thus there is a fixed small $t > 0$ and a neighborhood of P on which every active inequality increases uniformly along $Q + s\nu$, $0 \le s \le t$. Inactive inequalities remain positive after shrinking this neighborhood and t. The central strip conditions are preserved by the same translation for nearby points and sufficiently small t; at a strip endpoint it moves strictly inward. It follows that $P + t\nu \in D$ and, for all sufficiently near $Q \in D$, the entire segment $Q + [0, t]\nu$ lies in $D \cap U$.

For $\mathcal{D}_L$, use instead the direction $\nu = (1, 1, 0, 0)$ and the defining functions

$$g_1 = \cosh x \tanh u - |\cos y|, \qquad g_2 = \cosh u \tanh x - |\cos v|.$$

On $x, u \ge 0$, their derivatives in this direction are

$$\sinh x \tanh u + \cosh x \operatorname{sech}^2 u > 0, \qquad \sinh u \tanh x + \cosh u \operatorname{sech}^2 x > 0.$$

The same translation also opens $x, u > 0$. The absolute cosine terms are fixed during the translation, so they create no differentiability issue. This proves the same segment property.

In either case choose a small ball about $P + t\nu$ contained in $D \cap U$. Shrink V about P until every translated endpoint $Q + t\nu$, $Q \in D \cap V$, lies in that ball. Join any two such endpoints by a straight segment in the ball and use their translation segments to join the original points. Thus $D \cap V$ lies in one component of $D \cap U$. A component accumulating at P meets $D \cap V$ and hence is that component.

On a fixed-ratio slice the HT direction $-P$ preserves the slice; the same derivative proof applies. For LT use the real direction $(1, \lambda, 0, 0)$, or equivalently the horizontal translation $\zeta \mapsto \zeta + s$ of the epigraph $a > L_\lambda(b)$. The preceding ball-and-segment argument is then entirely within the complex line. □

Definition 2.15. For $f \in \mathcal{O}(D)$ and $P \in \partial D$, a local holomorphic continuation through P means a holomorphic function F on an open neighborhood U of P which agrees with f on some connected component E of $D \cap U$ satisfying $P \in \overline{E}$. This definition allows agreement on only the component reached by a continuation path; it does not require agreement on every component of $D \cap U$.

Corollary 2.16 (Propagation from dense genuine obstructions)**.** *Let D be any of the domains in Lemma 2.14, and let $f \in \mathcal{O}(D)$. Suppose a dense subset $G \subset \partial D$ consists of points at which no holomorphic function on a neighborhood can agree with f on the full intersection with D. Then f has no local holomorphic continuation through any point of ∂D, even in the componentwise sense of Definition 2.15.*

Proof. If F, U, E gave such a continuation at P, Lemma 2.14 would give $V \subset U$ with $D \cap V \subset E$, and therefore $F = f$ on $D \cap V$. Choose $Q \in G \cap V$ and a neighborhood of Q contained in V. Restricting F gives the forbidden continuation at Q. □

Remark 2.17. Proposition 2.9, Proposition 2.12, and Corollary 2.16 are a geometric bridge. To apply them to a susceptibility, the dense obstructions must be proved for the entire physical sum, with its normalization and tail included. For a fixed-ratio corollary, its local singularity calculation must remain valid along the corresponding analytic curve $B = B(t)$ in the coordinates of Proposition 2.10; a bare joint nonextension assertion does not imply the same for its restriction. Nothing here claims that all of Σ, which has interior, is a singularity set.

2.6 Pseudoconvexity and the Hartogs question

Proposition 2.18 (An explicit plurisubharmonic exhaustion)**.** *Every component of* $\mathbb{C}^2 \setminus \Sigma$ *has a continuous plurisubharmonic exhaustion. In particular the explicit physical components above are pseudoconvex.*

Proof. Set

$$m(K) = \min_{(a,b)\in[-1,1]^2} |A - Sa - Tb|.$$

Compactness of the square makes m continuous, and its zero set is exactly Σ. On a complementary component, each $-\log|A - Sa - Tb|$ is pluriharmonic, and

$$-\log m(K) = \sup_{(a,b)\in[-1,1]^2} -\log|A - Sa - Tb|$$

is continuous and locally bounded above. The submean inequality on every complex line, or the usual locally bounded supremum criterion, shows that it is plurisubharmonic. Consequently

$$\Phi(K) = |K_x|^2 + |K_y|^2 + \max\{0, -\log m(K)\}$$

is continuous and plurisubharmonic. It tends to infinity at every finite boundary point, because $m \to 0$, and at infinity by its quadratic term. More explicitly, any sequence in a bounded sublevel set has a convergent subsequence with m bounded away from zero; its limit lies in the same complementary component. Thus its sublevel sets are relatively compact there. □

Remark 2.19. The exhaustion resolves the immediate geometric Hartogs test: these components are not domains with a compulsory Hartogs filling caused by failure of pseudoconvexity. It does not prove that a particular physical function fails to extend. That assertion still requires genuine whole-sum obstructions and the local continuation argument above.

3 The physical observable and joint holomorphy

Throughout this paper $z = 2K_x$ and $w = 2K_y$ are independent complex coordinates. Write

$$S = \sinh z, \qquad T = \sinh w, \qquad A = \cosh z \cosh w, \qquad p = ST.$$

No inverse temperature is introduced as an additional independent variable in $\mathbb{C}^2$. For real positive K_x, K_y, the dimensionless uniform field is denoted by η. The zero-field responses are

$$\mathcal{X}_H(K_x, K_y) = \sum_{(m,l)\in\mathbb{Z}^2} \langle\sigma_{0,0}\sigma_{m,l}\rangle, \qquad 0 < p < 1, \tag{3.1}$$

$$\mathcal{X}_L(K_x, K_y) = \sum_{(m,l)\in\mathbb{Z}^2} \big(\langle\sigma_{0,0}\sigma_{m,l}\rangle_+ - M^2\big), \qquad p > 1, \tag{3.2}$$

$$M^2 = (1 - p^{-2})^{1/4}.$$

The high-temperature state is the unique Gibbs state. The subscript + means the infinite-volume plus state; the fourth root in the last line is positive. These definitions include the origin, both axes and all quadrants. In particular the low-temperature onsite contribution is $1 - M^2$, not 1.

3.1 The precise imported physical interface

Only the real, noncritical spin form factors are imported from the exact solution. The relevant finite-cylinder formulas at arbitrary separation are [1, (35)–(39)]; their finite-lattice matrix-element factorization is proved in [2, Theorem 6.1], with the paramagnetic vacuum change in its Remark 6.2. In our convention put

$$B = T/S > 0, \quad C = A/S, \quad \cosh\gamma(q) = C - B\cos q, \quad \gamma(q) > 0.$$

Then the infinite-plane coefficient is

$$f_N(m,l) = \frac{B^{N^2/2}}{N!} \int_{\mathbb{T}^N} \prod_{i=1}^{N} \frac{e^{-|m|\gamma(q_i)+ilq_i}}{\sinh\gamma(q_i)} \prod_{i<j} \left\{ \frac{\sin((q_i-q_j)/2)}{\sinh((\gamma(q_i)+\gamma(q_j))/2)} \right\}^2 \prod_{i=1}^{N} \frac{\mathrm{d}q_i}{2\pi}. \tag{3.3}$$

With positive real square roots the two correlations are

$$\langle \sigma_0 \sigma_{m,l} \rangle = \xi_H \sum_{\substack{N\geq 1 \\ N \text{ odd}}} f_N(m,l), \qquad \xi_H = (1-p^2)^{1/4} p^{-1/2}, \tag{3.4}$$

$$\langle \sigma_0 \sigma_{m,l} \rangle_+ = \xi_L \left(1 + \sum_{\substack{N\geq 2 \\ N \text{ even}}} f_N(m,l) \right), \qquad \xi_L = (1-p^{-2})^{1/4} = M^2. \tag{3.5}$$

This is the normalization of [3, (1.8)–(1.10)]. In particular the anisotropy exponent in (3.3) is $N^2/2$.

Here are the limit and state identifications needed in using that interface. On a real noncritical compact set the dispersion has a uniform positive gap and is analytic in an angular strip. The finite-circumference corrections $\eta, \Lambda^{-1}, \log\xi_T$ in [1] tend uniformly to zero, by the exponentially convergent trapezoidal rule applied to their sum-minus-integral expressions. Here is a quantitative version that also covers the highest particle indices. If L is the circumference, the unnormalized Ramond and Neveu–Schwarz sums of a strip-analytic function differ by $O(Le^{-hL})$: their Fourier expansions select the multiples of L, with respective factors 1 and $(-1)^k$. Apply this to γ and to $\log(1-e^{-\gamma(q)-\gamma(q')})$. The mixed double-grid difference of $\log\sinh((\gamma(q)+\gamma(q'))/2)$ in [1, (25)] gives

$$|\Lambda_L^{-1}| + \sup_q |\eta_L(q)| \leq CLe^{-hL}, \qquad |\log\xi_{T,L}| \leq CL^2 e^{-2hL}.$$

Thus at each fixed separation the entire correction is bounded by $C_{m,l}C^N$, uniformly in L, N. Terms with $N > L$ vanish on the discrete grid. In particular no estimate at a fixed particle number is being used to pass an unbounded particle sum through the thermodynamic limit. For the real roots $0 < \rho(q) = e^{-\gamma(q)} \leq r < 1$,

$$\left| \frac{\rho(q)-\rho(q')}{1-\rho(q)\rho(q')} \right| \leq r < 1.$$

The matching argument in Lemma 4.1 is an algebraic identity on discrete angular grids as well as on continuous circles. It gives a common bound $\exp\{-cN^2 + O(N\log(N+1))\}$ at fixed separation, including the finite-size one-body corrections and factorial normalization. Consequently the particle sum passes through the cylinder limit; each fixed-index Riemann sum gives (3.3).

Every local cylinder limit is a translation-invariant and spin-flip invariant Gibbs state. The Gibbs-state theorem [5, Theorem 1] applies with the vertex-transitive group chosen to be translations: the two unequal positive nearest-neighbor couplings satisfy its nonnegativity, invariance, irreducibility and summability hypotheses. That state is the equal mixture of the plus and minus states. Its two-spin correlations equal those of the plus state. This proves the state identification in (3.5); subtracting $M^2 = \xi_L$ removes exactly its zero-particle term. No response of a zero-mean mixed state is substituted for (3.2).

3.2 Double contours and the full lattice

Let ζ, η now denote independent contour variables, and set

$$\mathcal{D}(\zeta,\eta;z,w) = A - \frac{S}{2}(\zeta+\zeta^{-1}) - \frac{T}{2}(\eta+\eta^{-1}), \qquad a(v,v') = \frac{v-v'}{1-vv'}.$$

If $X = \prod_i \zeta_i$ and $Y = \prod_i \eta_i$, the same coefficient has the symmetric representation

$$f_N(m,l) = \frac{p^{N/2}}{N!} \oint \cdots \oint X^{|m|-1} Y^{|l|-1} \prod_{i<j} a(\zeta_i,\zeta_j) a(\eta_i,\eta_j) \prod_{i=1}^{N} \frac{\mathrm{d}\zeta_i \, \mathrm{d}\eta_i}{(2\pi\mathrm{i})^2 \mathcal{D}(\zeta_i,\eta_i)}. \tag{3.6}$$

The circles have radii strictly less than and sufficiently close to one. This formula can be checked directly on the real physical region, as follows. At $\eta = e^{\mathrm{i}q}$ the inside root is $\zeta = \rho = e^{-\gamma(q)}$, and

$$\mathcal{D}_\zeta(\rho,\eta) = \frac{S(1-\rho^2)}{2\rho^2}, \qquad \frac{\sqrt{p}}{\mathcal{D}_\zeta(\rho,\eta)} = \sqrt{B}\,\frac{\rho}{\sinh\gamma}.$$

Subtracting two dispersion equations gives the exact canceled pair

$$a(\rho_i,\rho_j)a(\eta_i,\eta_j) = -B\frac{(\eta_i-\eta_j)^2\rho_i\rho_j}{\eta_i\eta_j(1-\rho_i\rho_j)^2} = B\left\{\frac{\sin((q_i-q_j)/2)}{\sinh((\gamma_i+\gamma_j)/2)}\right\}^2 =: P_{ij}. \tag{3.7}$$

The residues therefore give $B^{N/2}$, and the pairs give $B^{N(N-1)/2}$. This proves the full exponent and the normalization in (3.3). Reflection of either spatial coordinate is legitimate; equality under interchanging the separations while holding unequal couplings fixed is not required.

The exact product of the two spatial geometric series is

$$\sum_{m,l\in\mathbb{Z}} X^{|m|-1}Y^{|l|-1} = \frac{(1+X)(1+Y)}{XY(1-X)(1-Y)} = \frac{1}{XY} + \frac{2(X^{-1}+Y^{-1})}{(1-X)(1-Y)}. \tag{3.8}$$

The first term is the onsite coefficient. Denoting by T_N the integral with the second numerator but without its factor two gives $C_N = f_N(0,0) + 2T_N$, where C_N sums all sites. The angular form of this complete coefficient is

$$C_N = \frac{1}{N!}\int_{\sum_i q_i = 0 \pmod{2\pi}} G_N(q)\frac{1+Z}{1-Z}\prod_{i=1}^{N-1}\frac{\mathrm{d}q_i}{2\pi}, \qquad G_N = B^{N/2}\prod_i \frac{1}{\sinh\gamma_i}\prod_{i<j}P_{ij}, \quad Z = \prod_i \rho_i. \tag{3.9}$$

There are $N-1$, not N, independent normalized measures in this formula; $q_N = -\sum_{i<N} q_i$ modulo 2π.

Proposition 3.1 (Joint normal convergence)**.** *On each of the two domains $\mathcal{D}_H, \mathcal{D}_L$ specified by the independent spectral geometry, the physical germ has a single-valued jointly holomorphic continuation. It is given by*

$$\mathcal{X}_H = (1-p^2)^{1/4} \sum_{\substack{N\geq 1 \\ N \text{ odd}}} p^{(N-1)/2} F_N, \tag{3.10}$$

$$\mathcal{X}_L = (1-p^{-2})^{1/4} \sum_{\substack{N\geq 2 \\ N \text{ even}}} p^{N/2} F_N, \tag{3.11}$$

where F_N is the integral (3.6) *with its $p^{N/2}$ factor removed and with the multiplier* (3.8)*. The sums converge normally, with every fixed mixed complex derivative, on compact subsets. The fourth roots are the binomial branches supplied by $|p|<1$ on $\mathcal{D}_H$ and $|p|>1$ on $\mathcal{D}_L$.*

Proof. On a compact parameter neighborhood in $\mathbb{C}^2 \setminus \Sigma$, the polynomial $\mathcal{D}$ has a nonzero minimum modulus on the product of unit circles. By uniform continuity that gap persists on a common thin product annulus and a slightly larger parameter polydisk. Choose fixed radii $r_x, r_y < 1$ in the annulus. All Schur entries on these circles are bounded by a constant independent of N. For even N, the Schur Pfaffian identity and Hadamard's determinant inequality give

$$\left| \prod_{i<j} a(\zeta_i, \zeta_j) \right| = |\operatorname{Pf}(a(\zeta_i, \zeta_j))| \le C^{N/2} N^{N/4}.$$

For odd N, adjoin a ghost variable -1; its entries are the removable constant one and the dimension becomes $N+1$. Apply the estimate separately to the two contour-variable families. The one-body factors and contour lengths cost C^N. Also $1-X$ and $1-Y$ are bounded away from zero uniformly in N, whereas X^{-1}, Y^{-1} cost only C^N. Thus, after increasing constants,

$$|F_N| \le C_0 C_1^N \frac{N^{(N+1)/2}}{N!}. \tag{3.12}$$

The right side is summable after multiplication by any fixed exponential in N. Multivariate Cauchy estimates on the larger polydisk give the same assertion for every fixed mixed derivative. On overlaps, radii sufficiently close to one can be deformed into one another inside the common gap annulus; all pair and product denominators stay nonzero. The local integrals therefore glue. All powers of p within (3.10)–(3.11) are integers. The binomial fourth roots introduce no monodromy, and the series prove joint holomorphy, including the HT axes $S = 0$ or $T = 0$.

On a real physical compact set the same contours in (3.6) additionally give $r_x^{N|m|} r_y^{N|l|}$. Their double geometric sum is bounded uniformly in $N \ge 1$. Thus the lattice and particle sums are absolutely interchangeable, and (3.8) gives exactly (3.1)–(3.2), with the stated zero-particle subtraction. Equality on a real open physical set determines the joint germ: iterating the one-variable identity theorem over a real rectangle proves the required uniqueness. □

The canonical transfer root has an equally concrete branch description. For real q, outside Σ the equation

$$S\rho^2 - 2(A - T\cos q)\rho + S = 0$$

has a unique root of modulus less than one. For $S \neq 0$ it is the Joukowski inside root

$$\rho = W - \sqrt{W^2 - 1}, \qquad W = (A - T\cos q)/S, \qquad \sqrt{W^2-1} \sim W \quad (W \longrightarrow \infty).$$

At $S = 0$ the implicit-function theorem gives its removable value zero, since $A - T\cos q \neq 0$. This construction is jointly holomorphic locally in the couplings and in a complex angular neighborhood. On positive real noncritical parameters, $(A - S - T)(A + S + T) = (p-1)^2$ shows $W(q) > 1$, so it is the physical decaying root. We take $\gamma = -\log \rho$ only on local patches where $\rho \neq 0$; no global logarithm through a root zero is asserted.

The angular representation used below is identified with this global continuation coefficient by coefficient. Indeed $1/\sinh\gamma = 2\rho/(1-\rho^2)$ and the rational expression for P_{ij} in (3.7) contain no logarithm. After including the physical normalization, their one-body prefactors are

$$(1-p^2)^{1/4} \frac{p^{(N-1)/2}}{S^N} \quad (N \text{ odd}), \qquad (1-p^{-2})^{1/4} \frac{p^{N/2}}{S^N} \quad (N \text{ even}).$$

All powers here are integers. Hence the normalized angular coefficient is single valued and holomorphic wherever $S \neq 0$ in the corresponding physical domain; the real angular torus has

uniform root and denominator gaps on compact subsets. The set $\mathcal{D}_H \setminus \{S = 0\}$ is connected: radial contraction enters a small polydisk without crossing $z = 0$, and that polydisk minus $z = 0$ is connected. The domain $\mathcal{D}_L$ already has $S \neq 0$. The identity theorem from the real physical rectangle therefore gives equality with the double-contour coefficients throughout these sets. Near a regular edge $S, T \neq 0$, this also fixes the compatible local odd square root $\sqrt{B} = \sqrt{p}/S$. Normal convergence of the identified series follows from Proposition 3.1. Thus every boundary calculation below is a calculation of the global physical sum on its actual interior side.

Remark 3.2 (A restriction that really is removable)**.** On $K_y = 0$, the lattice splits into independent horizontal chains. For $t = \tanh K_x$ near zero its response is

$$1 + 2 \sum_{m \geq 1} t^m = \frac{1 + t}{1 - t} = e^{2K_x}.$$

This agrees with (3.10): $p = 0$, only $N = 1$ survives, and $F_1 = (A - S)^{-1} = e^{2K_x}$. The restriction is entire, although the joint domain has boundary points on this line. This exact example is why a bare joint nonextension assertion cannot be used to prove the slice part of the main theorem.

3.3 Energy-field normalization

For fixed real $J_x, J_y > 0$ and physical temperature $(K_x, K_y) = (\beta J_x, \beta J_y)$, a field of energy units enters the Hamiltonian as $-h \sum_v \sigma_v$, and $\eta = \beta h$. Finite-volume differentiation gives $\chi = \beta \mathcal{X}$. The same statement holds after taking the prescribed infinite-volume phase limit. To justify the latter without exchanging an uncontrolled field limit and infinite sum, use the GHS inequality for arbitrary nonnegative couplings and fields [6, Theorem 1.5]. Pinning removed spins by fields tending to $+\infty$ shows that plus-box covariances increase with the box. Their nonnegative sums increase to the finite sum proved above. GHS concavity gives, for the plus-box magnetization,

$$m_V(h) - m_V(0) \leq h\, m_V'(0) \leq \beta h\, \mathcal{X}, \qquad h \geq 0.$$

The plus-state limit yields $m'(0+) \leq \beta \mathcal{X}$. Conversely turn on h only on a finite set E. The resulting bounded local change of measure has derivative $\beta \sum_{v \in E} \mathrm{Cov}_{+,0}(\sigma_0, \sigma_v)$. Ferromagnetic monotonicity bounds its magnetization above by the uniform-field magnetization, with equality at $h = 0$. Increasing E proves the reverse inequality. Thus $m'(0+) = \beta \mathcal{X}$. Spin-flip symmetry gives the two-sided HT derivative; in LT the derivative is the plus-phase right derivative (or its minus-phase counterpart). This statement is made on temperature lines only.

4 Regular spectral edges: a genuinely complex-parameter theorem

The following estimates use independent complex dispersion parameters, not uniformity inferred from a collection of real ratios. On a regular edge, exchange the two lattice axes if necessary. In the local coordinates $C = A/S,\ B = T/S$ of the geometry section, one edge is $C - B = b$, $b = \cos\phi \in (-1, 1)$. Choose the side for which $\Im B_0 > 0$; conjugation handles the other side. Let $B(u)$ be any holomorphic function near zero with $B(0) = B_0$, and set

$$W(q; u) = b + u + B(u)(1 - \cos q). \tag{4.1}$$

Both a constant-B complex transversal in $\mathbb{C}^2$ and each nonramified physical temperature-line transversal have this form.

At $u = 0$, the inside root lies in the closed lower half disk, with $\rho(0) = e^{-\mathrm{i}\phi}$. Compactness gives

$$\Im\rho(q) \le -\delta < 0, \qquad |\rho(q)| \le 1.$$

There is no $W = \pm 1$ on the real angular circle. The roots, their chosen logarithms, the one-body factors, and the canceled pairs (3.7) are analytic on one fixed complex neighborhood in (u, B, q). On $u = \mathrm{i}\varepsilon$, $\varepsilon > 0$,

$$\Re\gamma(q; \mathrm{i}\varepsilon) \ge c\{\varepsilon + \mathrm{dist}(q, 0)^2\}. \tag{4.2}$$

Indeed $\Im W = \varepsilon + \Im B(\mathrm{i}\varepsilon)(1 - \cos q)$ has this lower bound; the identity $\Im W = \sinh(\Re\gamma)\sin(\Im\gamma)$ on the compact branch then gives (4.2). This is the complement side of the parallelogram edge and hence the side of the bordering physical component.

Lemma 4.1 (Strict contraction on a complex polydisk)**.** *There are $\kappa > 0$, $C < \infty$, and a fixed complex parameter polydisk about this edge such that, for every fixed mixed derivative ∂^ν in the parameters,*

$$|\partial^\nu G_N(q)| \le C_\nu \exp\{-\kappa N^2 + CN\log(N+1)\} \tag{4.3}$$

for all real angles and every $N \ge 1$.

Proof. The identity

$$|1 - vv'|^2 - |v - v'|^2 = (1 - |v|^2)(1 - |v'|^2) + 4\Im v\,\Im v'$$

gives a strict bound $|a(\rho_i, \rho_j)| \le q_0 < 1$. The denominator is separated from zero. By continuity a slightly larger constant below one still works on a common *complex* parameter polydisk; individual root moduli need not stay below one there. Every canceled pair P_{ij} remains bounded by C.

For even N, expand the Schur Pfaffian in the $\eta_i = e^{\mathrm{i}q_i}$ variables only. For each matching, combine its $N/2$ edges with the corresponding ρ factors using (3.7). The other $N(N-2)/2$ root factors retain their strict contraction. Thus

$$\left|\prod_{i<j} P_{ij}\right| \le (N-1)!!\, C^{N/2} q_0^{N(N-2)/2}.$$

For odd N, adjoining the constant ghost entry gives

$$\left|\prod_{i<j} P_{ij}\right| \le N!!\, C^{(N-1)/2} q_0^{(N-1)^2/2}.$$

These identities first hold away from reciprocal η pairs and extend by the analytic canceled formula. There is no division by a root-Schur zero. One-body factors cost C^N. Cauchy's inequalities on the larger polydisk prove all the derivative bounds. They are applied to G_N, never across a pole of $(1+Z)/(1-Z)$. □

Lemma 4.2 (The entire differentiated higher-index tail)**.** *Suppose $\phi = 2\pi a/\ell$, where ℓ is an odd prime and $1 \le a < \ell/2$. For odd parity let $n = \ell$ and $k = (n^2-1)/2$; for even parity let $n = 2\ell$ and $k = n^2/2 - 1$. For every $0 \le j \le k$,*

$$\sum_{\substack{N > n \\ N \text{ of the chosen parity}}} |\partial_u^j C_N(\mathrm{i}\varepsilon)| = O_j(1) \qquad (\varepsilon \downarrow 0).$$

All lower indices of the chosen parity are holomorphic near $u = 0$.

Proof. Write $Q_0 = \sum_i \operatorname{dist}(q_i, 0)^2$. Equation (4.2) gives

$$|1 - Z| \geq c \min\{1, N\varepsilon + Q_0\}.$$

Since all fixed derivatives of γ are bounded, differentiating the exponential and rational function gives

$$\left|\partial_u^r \frac{1+Z}{1-Z}\right| \leq C_j N^j (1 + Q_0^{-j-1}), \qquad 0 \leq r \leq j.$$

If $N \geq 2j+4$, retain the first $2j+3$ independent angle squares in Q_0. Their inverse $(j+1)$-st power is integrable in that fixed real dimension; all unused normalized angular measures have mass one. Therefore Lemma 4.1 gives

$$|\partial_u^j C_N(\mathrm{i}\varepsilon)| \leq \frac{C_j N^j}{N!} \exp\{-\kappa N^2 + CN \log(N+1)\},$$

uniformly in ε. This covers an actual infinite index range.

For each remaining fixed N, the only configuration with $|Z| = 1$ is $q_1 = \cdots = q_N = 0$. There $Z = e^{-\mathrm{i}N\phi}$; the first resonant odd index is ℓ, and the first resonant even index is 2ℓ. Lower indices have a denominator gap and hence are holomorphic. Near the collision of a higher index use $q_N = -\sum_{i<N} q_i$, $Q = \sum_i q_i^2$, and $\Delta(q) = \prod_{i<j}(q_i - q_j)$. Every fixed numerator derivative has the factor Δ^2. The worst radial bound for the j-th derivative is

$$r^{N(N-1)} r^{N-2} (\varepsilon + r^2)^{-j-1} \, \mathrm{d}r.$$

Its limiting exponent $N^2 - 2j - 4$ is greater than -1 for every larger same-parity N and $j \leq k$. This handles the finite intervening range and completes the entire tail estimate. □

Proposition 4.3 (Nonzero full-sum edge obstruction on every holomorphic transversal)**.** *Under* (4.1) *and the rational-phase hypotheses of Lemma 4.2, the complete sum of the selected parity has no holomorphic extension through $u = 0$. The assertion is locally uniform in independent complex B parameters. It remains valid after multiplication by a holomorphic nonvanishing prefactor and after a nonramified change of transversal coordinate. It holds for both endpoint signs in the even case.*

Proof. We give the first amplitude, including its constrained measure, so that noncancellation does not depend on an unspecified singular coefficient. Put $\alpha = (n^2 - 1)/2$ and

$$P_n = \prod_{r=1}^{n-1} \frac{r!}{2^r}.$$

The Gaussian Vandermonde period on the *independent-coordinate* measure is

$$\int_{q_n = -\sum_{i<n} q_i} e^{-Q} \Delta(q)^2 \, \mathrm{d}q_1 \cdots \mathrm{d}q_{n-1} = \frac{n! \, \pi^{(n-1)/2} P_n}{\sqrt{n}} =: \mathcal{G}_n. \tag{4.4}$$

To derive it, expand the ordinary n-variable Vandermonde determinant in monic Hermite polynomials. Their Gaussian norms are $\sqrt{\pi}\, r!/2^r$, so that integral is $n! \pi^{n/2} P_n$. Setting $x_i = q_i + m$, $\sum q_i = 0$, has Jacobian n, and the m integral is $\sqrt{\pi/n}$. Division proves (4.4).

On the unique collision chart,

$$\gamma(q; u) = \mathrm{i}\phi - \frac{\mathrm{i}}{\sin\phi} \left(u + \frac{B_0 q^2}{2} \right) + O(u^2 + uq^2 + q^4).$$

The leading singular integral for C_n is consequently

$$\frac{2\mathrm{i}^{1-n^2}B_0^{n^2/2}}{n!(2\pi)^{n-1}2^{n(n-1)}\sin^{n^2-1}\phi}\int\frac{\Delta(q)^2}{nu+(B_0/2)Q}\,\mathrm{d}q_1\cdots\mathrm{d}q_{n-1}. \tag{4.5}$$

For odd n the local square root of B is the one used in the physical angular coefficient. Its sign can change the displayed amplitude's sign but cannot make it zero. For even n the power in the original integrand is an integer; the square root below is fixed by the Gaussian period.

After k derivatives, the model integral is evaluated from

$$\int\frac{\Delta(q)^2}{(a+bQ)^{k+1}}\,\mathrm{d}q=\mathcal{G}_n\,b^{-\alpha}\frac{\Gamma(k+1-\alpha)}{\Gamma(k+1)}a^{\alpha-k-1},$$

initially in a common right half-plane and then by analytic continuation to $a=nu,\ b=B_0/2$. More explicitly, rotate both coefficients by $-\mathrm{i}$: $a'=n\varepsilon>0$ and $b'=-\mathrm{i}B_0/2$ has positive real part. The absolutely convergent Laplace–Gaussian calculation for a',b' fixes the continuation and its square root. No assumption $\Re B_0>0$ is made. For odd n, $k=\alpha$ is a multiple of four; for even n, $k=\alpha-\frac{1}{2}$ is odd. Substitution gives

$$\lim_{\varepsilon\downarrow0}u\,\partial_u^kC_n(u)=\frac{2^{1-k}\sqrt{B_0}\,P_nn^{k-3/2}}{\pi^{(n-1)/2}\sin^{2k}\phi}\neq0, \qquad u=\mathrm{i}\varepsilon,\quad n\text{ odd}, \tag{4.6}$$

$$\lim_{\varepsilon\downarrow0}u^{1/2}\partial_u^kC_n(u)=-\mathrm{i}\,\frac{2^{1-\alpha}\sqrt{B_0}\,P_nn^{k-1}}{\pi^{(n-2)/2}\sin^{n^2-1}\phi}\neq0, \qquad u=\mathrm{i}\varepsilon,\quad n\text{ even}. \tag{4.7}$$

In the second line $u^{1/2}=e^{\mathrm{i}\pi/4}\sqrt{\varepsilon}$.

For completeness these are limits of the actual fixed-n integral. On its collision chart write the exact integrand as $\Delta^2f(q,u)/H(q,u)$. It is even under simultaneous $q\mapsto-q$, and, with $c=-\mathrm{i}/\sin\phi$,

$$H=c\{nu+(B_0/2)Q\}+O((|u|+Q)^2),\quad H_u=cn+O(|u|+Q),\quad f=f_0+O(|u|+Q).$$

Both the exact and model denominators have modulus at least $c_1(\varepsilon+Q)$ on $u=\mathrm{i}\varepsilon$. The top derivative has leading term $(-1)^kk!fH_u^k/H^{k+1}$; all other terms have denominator power at most k. Subtracting the constant quadratic model therefore gives the pointwise bound

$$|\partial_u^k(\text{exact integrand}-\text{model integrand})|\le C\Delta^2(\varepsilon+Q)^{-k}.$$

Its radial integral is $O(\log(1/\varepsilon))$ for odd $k=\alpha$, and $O(1)$ for even $k=\alpha-1/2$. Extending the differentiated model from the chart to the whole tangent plane costs $O(1)$, since its radial tails are R^{-3} and R^{-2}, respectively. Outside the chart the exact denominator has a gap. These bounds prove the two claimed limits, including for the actual holomorphic function $B(u)$, whose contribution to the denominator remainder is $O(uQ)$.

Lemma 4.2 bounds all higher derivatives in the entire tail, and lower indices are analytic. Lower derivatives of the even first term are bounded; the next lower odd derivative may be logarithmic, but all are $o(u^{-1})$. Thus the nonzero top amplitudes cannot be canceled in the full sum or by product-rule terms from an analytic nonvanishing physical prefactor. The chain rule proves the statement for every nonramified curve parameter.

At the negative endpoint replace q_i by $\pi+q_i'$. For even N the total momentum constraint is unchanged and $B^{N^2/2}=(-B)^{N^2/2}$. Apply the preceding proof with $-B$, conjugating if necessary to have positive imaginary part. Exchanging transfer axes handles the other two edges. For odd N this shift instead makes the total reduced angle an odd multiple of π; that distinct case is addressed by the full boundary-jet theorem in the next section. □

5 The full odd sum at a regular negative spectral edge

This section proves a local theorem for an arbitrary holomorphic anisotropy along a transverse analytic curve. In particular, the anisotropy is not held constant. All estimates retain the exact particle-dependent anisotropy factor, the complete constrained angular integral, and the sum over every odd particle number.

5.1 Statement, branch, and the physical side

Fix

$$\phi = \frac{2\pi r}{p} \in (0,\pi), \qquad (r,p)=1, \qquad p \geq 3 \text{ odd}, \qquad b = \cos\phi, \qquad s_\phi = \sin\phi > 0. \tag{5.1}$$

Let B be holomorphic on a disk about zero and suppose that

$$\Im B(0) < 0. \tag{5.2}$$

Shrink this disk so that $B \neq 0$ and $\Im B < 0$ throughout it, and fix one holomorphic logarithm of B. For $q = \pi + \delta$, define

$$W(t,\pi+\delta) = b + B(t)(t - 1 + \cos\delta), \qquad g(t,\delta) = \operatorname{arcosh} W(t,\pi+\delta), \qquad g(0,0) = \mathrm{i}\phi. \tag{5.3}$$

At $t=0$, W is in the open upper half-plane except at $\delta = 0 \bmod 2\pi$, where it equals $b \in (-1,1)$. Consequently the branch in (5.3) extends to a fixed complex neighborhood of all real angles and a fixed parameter disk. It is even and 2π-periodic in δ. Put $\gamma(t,q) = g(t,q-\pi)$ and $\rho(t,q) = e^{-\gamma(t,q)}$. At $t=0$, $|\rho| \leq 1$, and the compact set of these roots lies strictly in the lower half-plane:

$$\Im\rho(0,q) \leq -\delta_0 < 0. \tag{5.4}$$

The logarithm used for γ is fixed by $0 < \Im\gamma(0,q) < \pi$. In particular all the one-body denominators and the canceled pair denominators below are nonzero on a fixed complex neighborhood.

For odd $n \geq 1$, let

$$\mathcal{T}_n = \{(q_1,\dots,q_n) \in (\mathbb{R}/2\pi\mathbb{Z})^n : q_1 + \cdots + q_n = 0 \bmod 2\pi\}, \qquad q_n = -\sum_{i<n} q_i \bmod 2\pi.$$

The measure on this torus is $\mathrm{d}q_1 \cdots \mathrm{d}q_{n-1}$. Include the normalization in the numerator:

$$H_n(t,q) = \frac{B(t)^{n^2/2}}{n!(2\pi)^{n-1}} \prod_{i=1}^{n} \frac{1}{\sinh\gamma(t,q_i)} \prod_{i<k} \left[\frac{\sin((q_i-q_k)/2)}{\sinh((\gamma(t,q_i)+\gamma(t,q_k))/2)}\right]^2, \tag{5.5}$$

$$\Gamma_n(t,q) = \sum_{i=1}^{n} \gamma(t,q_i), \tag{5.6}$$

$$C_n(t) = \int_{\mathcal{T}_n} H_n(t,q) \coth\!\left(\frac{\Gamma_n(t,q)}{2}\right) \mathrm{d}q. \tag{5.7}$$

The power of B uses the chosen common logarithm. Changing its square-root sign changes every odd coefficient by the same sign.

The relevant open side of the boundary is

$$\mathcal{U}^- = \{\, |t| < r_0 :\ \Im(B(t)t) > 0 \,\}. \tag{5.8}$$

Here

$$\Im W(t,\pi+\delta)=\Im(B(t)t)-\Im B(t)(1-\cos\delta)>0$$

for every real angle, so the chosen roots satisfy $|\rho|<1$. This is the local physical side when the regular negative endpoint has the lower-half-plane root $\rho(0,\pi)=e^{-\mathrm{i}\phi}$. It contains $t=-\epsilon$ for every sufficiently small $\epsilon>0$. When these coordinates are obtained from $W(q)=C-B\cos q$, they are exactly

$$t=\frac{C+B-b}{B}.$$

The curve hypothesis is that this t is a nonramified local parameter. Thus any transverse analytic curve through the selected endpoint is included, with its actual function $B(t)$. The opposite orientation is obtained by complex conjugating all the data.

Theorem 5.1 (Universal analytic-curve obstruction at a negative edge)**.** *Under* (5.1)–(5.2)*, the series*

$$F(t)=\sum_{\substack{n\ge 1\\ n\ \text{odd}}} C_n(t)$$

converges normally on $\mathcal{U}^-$. Every inward derivative has a finite limit

$$a_j=\lim_{\epsilon\downarrow 0}\frac{F^{(j)}(-\epsilon)}{j!},\qquad j=0,1,2,\dots. \tag{5.9}$$

Nevertheless its formal boundary Taylor series has radius zero. More precisely, as n tends to infinity through odd multiples of p, there are integers j_n such that

$$j_n=n^2\left(\log n+\tfrac{1}{2}\log\log n-c_*+o(1)\right), \tag{5.10}$$

$$|a_{j_n}|^{1/j_n}\sim\frac{4j_n}{e\pi^2\log j_n}, \tag{5.11}$$

$$c_*=-\frac{3}{4}-\frac{3}{2}\log 2+\log\pi+\frac{1}{2}\log|B(0)|-\log s_\phi. \tag{5.12}$$

It follows that F has no holomorphic extension through $t=0$ from $\mathcal{U}^-$. If P is holomorphic near zero and $P(0)\neq 0$, the same nonextension conclusion holds for PF.

The hypotheses and all estimates are locally uniform for complex holomorphic families $B(t,\eta)$: it suffices to work on a common parameter disk and a relatively compact parameter polydisk on which $\Im B(0,\eta)\le-\beta<0$ and B has a common holomorphic bound. The constants can depend on that disk, that bound, β, and the fixed rational phase. The selected integers are allowed to depend on the family parameter, $j_n=j_n(\eta)$; a common sequence of selected integers for all η is not asserted.

We prove the theorem by identifying a single particle index that dominates selected, increasingly high Taylor coefficients in modulus. The other indices have arbitrary complex phases; no positivity or phase alignment of the infinite sum is assumed.

5.2 A uniform Gaussian numerator bound and all boundary jets

Write $\mathcal{A}(v,w)=(v-w)/(1-vw)$ and $y_i=e^{\mathrm{i}q_i}$. Subtraction of two dispersion equations gives the canceled identity

$$\mathcal{A}(\rho_i,\rho_k)\mathcal{A}(y_i,y_k)=B(t)\left[\frac{\sin((q_i-q_k)/2)}{\sinh((\gamma_i+\gamma_k)/2)}\right]^2. \tag{5.13}$$

This is an identity of holomorphic canceled functions, including reciprocal y-pairs. The elementary identity

$$|1-vw|^2-|v-w|^2=(1-|v|^2)(1-|w|^2)+4\Im v\Im w \tag{5.14}$$

and (5.4) imply $|\mathcal{A}(\rho_i,\rho_k)|\le\theta<1$ at $t=0$. The same strict bound, with a slightly larger $\theta<1$, holds on a fixed complex parameter disk for all real angles. Every complete canceled pair in (5.13) is bounded there.

For completeness, the Schur identity used here is $\mathrm{Pf}(\mathcal{A}(y_i,y_k))_{i,k=1}^N=\prod_{i<k}\mathcal{A}(y_i,y_k)$ for even N. It follows by induction in N: as rational functions of y_N, the two sides have the same simple residues at $y_N=1/y_k$. In checking the residues, the other paired factors cancel by $\mathcal{A}(a,b)\mathcal{A}(a,1/b)=1$, and the remaining product is the induction hypothesis. Their difference is constant because both are bounded at infinity, and that constant is zero at $y_N=y_{N-1}$. The identity then extends from generic tuples by rational continuation.

Apply this identity only to the y-variables, adjoining the ghost variable -1 because n is odd. The entry $\mathcal{A}(y_i,-1)$ is the removable constant one. Each of the $n!!$ matchings has $(n-1)/2$ real edges. Combine each matched y-factor with the corresponding ρ-factor by (5.13). The remaining $(n-1)^2/2$ ρ-factors are at most θ in modulus. Thus, with the one-body factors and normalization included,

$$\sup_{|t|<r_0,\ q\in\mathcal{T}_n}|H_n(t,q)|\le E_n,\qquad E_n=\exp\{-\kappa n^2+Cn\log(n+1)+C\} \tag{5.15}$$

for some $\kappa>0$. Initially the Pfaffian calculation can be made off its apparent reciprocal-pair divisors; the canceled identity extends the estimate to those divisors. There is no division by a possibly zero ρ-factor. The same estimate holds when every angle is in a sufficiently small fixed complex neighborhood of π. The latter version will control the deformation of the local integration cycle.

Choose shortest lifts $q_i=\pi+\delta_i \bmod 2\pi$, with $\delta_i\in[-\pi,\pi]$, and put

$$S(q)=\sum_{i=1}^n\delta_i^2.$$

The odd momentum constraint says $\sum_i\delta_i=w\pi$, where w is odd. Therefore

$$S(q)\ge\frac{\pi^2}{n}. \tag{5.16}$$

On the inward ray, compactness of the branch and $\Im W(-\epsilon,\pi+\delta)\ge c(\epsilon+\mathrm{dist}(\delta,2\pi\mathbb{Z})^2)$ imply

$$\Re g(-\epsilon,\delta)\ge c\big(\epsilon+\mathrm{dist}(\delta,2\pi\mathbb{Z})^2\big). \tag{5.17}$$

Indeed $\Im W=\sinh(\Re g)\sin(\Im g)$, and the relevant roots remain in a fixed compact set. All parameter derivatives of g are bounded on a fixed complex neighborhood.

It follows from (5.16) that C_n is holomorphic on $|t|<c/n^2$. On this disk, $\Re\Gamma_n\ge c/n$ after decreasing c, and hence $|\coth(\Gamma_n/2)|\le Cn$. The torus volume is absorbed in the linear exponential in (5.15). Cauchy's estimate therefore gives

$$|[t^j]C_n(t)|\le(Cn^2)^j\,CnE_n. \tag{5.18}$$

More generally, the disk of radius $c(\epsilon+n^{-2})$ about $-\epsilon$ has $\Re\Gamma_n\ge c(n\epsilon+n^{-1})$. The same argument bounds the jth derivative at $-\epsilon$ by $j!(Cn^2)^jCnE_n$, uniformly for small ϵ. On compact subsets of

$\mathcal{U}^-$, the kernel is bounded and (5.15) proves normal convergence. Dominated convergence for each fixed j now proves

$$a_j = \sum_{\substack{n\geq 1\\ n \text{ odd}}} [t^j]C_n(t), \qquad \sum_n |[t^j]C_n(t)| < \infty. \tag{5.19}$$

This establishes the existence of all the jets, but does not give a positive radius for their formal Taylor series.

5.3 An exact reduction to simple-pole periods

Fix a sufficiently small $\sigma > 0$. There is a fixed $c_1 > 0$ such that on $|t| < c_1/n$ the part with $S \geq \sigma$ is analytic with a bounded kernel. To see this, use $\Re\Gamma_n(0,q) \geq cS$ and the perturbation bound $|\Gamma_n(t,q) - \Gamma_n(0,q)| \leq Cn|t|$. Choose c_1 to be a sufficiently small multiple of σ.

In the region $S < \sigma$, set

$$D_n(t,q) = \Gamma_n(t,q) - n\mathrm{i}\phi.$$

The Taylor expansion of g at $(0,0)$ gives $|D_n(t,q)| \leq C(S + n|t|)$. If $p \nmid n$, the distance of $n\phi$ from $2\pi\mathbb{Z}$ is at least $2\pi/p$. Reduce σ, c_1 so that the perturbation is less than π/p. This part is also analytic, with bounded kernel, on $|t| < c_1/n$.

It remains to consider odd multiples of p and $S < \sigma$. Here $n\phi \in 2\pi\mathbb{Z}$, and

$$\coth(\Gamma_n/2) = \frac{2}{D_n} + \mathcal{R}(D_n),$$

where $\mathcal{R}$ is holomorphic on a fixed disk containing all these values for $|t| < c_1/n$. There is a unique small zero $t = T_n(q)$ of $D_n(t,q)$. This statement has a fixed-disk quantitative form. With $a_0 = B(0)/(\mathrm{i}s_\phi) \neq 0$, one has

$$\frac{1}{n}\partial_t D_n(t,q) = a_0 + O(|t| + \sigma)$$

on a fixed disk. By reducing that disk and σ, its distance from a_0 is less than $|a_0|/4$. Rouché's theorem on a circle of radius CS/n, followed by uniqueness from this derivative estimate, gives

$$|T_n(q)| \leq CS/n, \qquad D_n(t,q) = n(t - T_n(q))A_n(t,q), \tag{5.20}$$

where A_n is holomorphic on a fixed smaller disk and bounded away from zero. One can take

$$A_n(t,q) = \frac{1}{n}\int_0^1 \partial_t D_n(T_n(q) + u(t - T_n(q)), q)\ \mathrm{d}u.$$

All such disks and bounds are independent of n and q.

Put $J_n = H_n/A_n$. The exact identity

$$\frac{2H_n(t,q)}{D_n(t,q)} = \frac{2J_n(T_n(q),q)}{n(t - T_n(q))} + \frac{2}{n}\frac{J_n(t,q) - J_n(T_n(q),q)}{t - T_n(q)} \tag{5.21}$$

separates the pole from an analytic divided difference. The second term is bounded by CE_n, by fixed-disk Cauchy estimates. The $\mathcal{R}$ term has the same bound on $|t| < c_1/n$. Combining these

observations with the exterior and nonresonant regions gives

$$a_j = \sum_{\substack{n \text{ odd} \\ p|n}} A_{n,j}^{\mathrm{pole}} + R_j, \tag{5.22}$$

$$A_{n,j}^{\mathrm{pole}} = -\int_{S<\sigma} \frac{2H_n(T_n(q),q)}{\partial_t \Gamma_n(T_n(q),q)} T_n(q)^{-j-1}\, \mathrm{d}q, \tag{5.23}$$

$$|R_j| \le \sum_{n\ge 1} (Cn)^j \exp\{-\kappa n^2 + Cn\log(n+1) + C\} \tag{5.24}$$

$$\le \exp\{\tfrac{1}{2} j\log(j+2) + Cj + C\}. \tag{5.25}$$

For each n, the coefficient identity is first justified on the smaller disk $|t| < c/n^2$; no pole is crossed there. The last inequality follows by absorbing $Cn\log(n+1)$ into half the Gaussian and maximizing $j\log n - \kappa n^2/2$. This reduction isolates all small pole locations before any local change of angular variables is made.

5.4 Bounds for every winding and every particle index

The zero in (5.20) has a useful more precise estimate. Set

$$v_i = 1 - \cos\delta_i, \qquad \bar v = \frac{1}{n}\sum_i v_i.$$

At $t = \bar v$, the linear terms in the expansion of $\operatorname{arcosh}(b + B(\bar v)(\bar v - v_i))$ cancel on averaging. The remaining average is bounded by $C\, n^{-1}\sum_i (v_i - \bar v)^2$. The uniform nonzero derivative used in (5.20) therefore yields

$$T_n = \bar v + O\left(\frac{1}{n}\sum_i (v_i - \bar v)^2\right).$$

Since $v_i \ge 0$, $\max_i v_i \le C\sigma$, and $\bar v = (S/(2n))(1 + O(\sigma))$, this implies

$$|T_n(q)| \ge (1 - C\sigma)\frac{S}{2n}, \qquad |\partial_t \Gamma_n(T_n(q),q)| \ge cn. \tag{5.26}$$

Retain the exact central numerator size

$$H_{0,n} = \frac{|B(0)|^{n^2/2}}{n!(2\pi)^{n-1} 2^{n(n-1)} s_\phi^{n^2}}. \tag{5.27}$$

Dividing the sine factors by their linear terms and all the hyperbolic sine factors by $\mathrm{i}s_\phi$ gives

$$|H_n(T_n(q),q)| \le H_{0,n}\Delta(\delta)^2 e^{CnS}, \qquad \Delta(\delta) = \prod_{i<k}(\delta_i - \delta_k). \tag{5.28}$$

Here is the relevant accounting. The logarithm of the anisotropy ratio costs $Cn^2|T_n| \le CnS$. The sum of the absolute one-body phase deviations is at most CS, so all pair denominators together cost CnS. Finally $\sum_{i<k}|\delta_i - \delta_k|^2 \le CnS$ controls the sine divided differences. Thus no undetermined constant to the power n^2 replaces the displayed normalization.

For an odd winding w, write

$$\delta_i = \frac{w\pi}{n} + \eta_i, \qquad \sum_i \eta_i = 0, \qquad Q(\eta) = \sum_i \eta_i^2, \qquad m_{n,w} = \frac{\pi^2 w^2}{n}. \tag{5.29}$$

Then $S = Q + m_{n,w}$ and $\Delta(\delta) = \Delta(\eta)$. The winding region is exactly $Q < \sigma - m_{n,w}$ when it is nonempty. In particular there are at most $C\sqrt{n}$ windings. Define

$$\alpha_n = \frac{n^2-1}{2}, \qquad P_n = \prod_{r=1}^{n-1} \frac{r!}{2^r}, \qquad G_n = \frac{n!\pi^{(n-1)/2}P_n}{\sqrt{n}}. \tag{5.30}$$

On the real zero-sum plane, with measure $\mathrm{d}\eta_1 \cdots \mathrm{d}\eta_{n-1}$,

$$\int e^{-Q(\eta)}\Delta(\eta)^2 \, \mathrm{d}\eta = G_n, \tag{5.31}$$

$$\int \frac{\Delta(\eta)^2}{(Q(\eta)+m)^{j+1}} \, \mathrm{d}\eta = G_n m^{\alpha_n - j - 1}\frac{\Gamma(j+1-\alpha_n)}{\Gamma(j+1)} \quad (j+1 > \alpha_n). \tag{5.32}$$

To verify the constants, the unrestricted Gaussian Vandermonde integral is $n!\prod_{r=0}^{n-1}(\sqrt{\pi}\, r!/2^r) = n!\pi^{n/2}P_n$. This follows by writing the Vandermonde as a determinant of monic Hermite polynomials and integrating the determinant product. The monic squared norm is $\sqrt{\pi}\, r!/2^r$, obtained by integration by parts. In the decomposition $x_i = \eta_i + u$, $\sum \eta_i = 0$, the ordinary-coordinate Jacobian is n, and the mean integral is $\int e^{-nu^2}\,\mathrm{d}u = \sqrt{\pi/n}$. This proves (5.31). Homogeneity gives radial density proportional to $Q^{\alpha_n - 1}\,\mathrm{d}Q$, proving (5.32) by the ordinary beta integral.

Let $A^{\mathrm{pole}}_{n,j,w}$ denote the contribution of the wth winding to (5.23). Equations (5.26)–(5.32) imply

$$|A^{\mathrm{pole}}_{n,j,w}| \le \frac{CH_{0,n}}{n}e^{Cn\sigma}\left(\frac{2n}{1-C\sigma}\right)^{j+1} G_n m_{n,w}^{\alpha_n - j - 1}\frac{\Gamma(j+1-\alpha_n)}{\Gamma(j+1)} \quad (j+1 > \alpha_n). \tag{5.33}$$

The dependence on w here is the explicit factor $|w|^{-2(j+1-\alpha_n)}$.

There is also a bound without any condition on j/α_n. Use $|T_n| \ge c/n^2$, and use homogeneity to write

$$\int_{Q<\sigma} \Delta(\eta)^2 \, \mathrm{d}\eta = G_n \frac{\sigma^{\alpha_n}}{\Gamma(\alpha_n+1)}.$$

The elementary factorial bounds in (5.30) then give

$$\log\left(\sum_w |A^{\mathrm{pole}}_{n,j,w}|\right) \le 2j\log n - \frac{1}{2}n^2\log n + Cn^2 + Cj + C. \tag{5.34}$$

In detail, $\log P_n = \frac{1}{2}n^2\log n + O(n^2)$ and $\log\Gamma(\alpha_n+1) = n^2\log n + O(n^2)$; the remaining factors in $H_{0,n}G_n$ have logarithm $O(n^2)$. The number of windings costs only $O(\log(n+1))$. This proves (5.34) also when n is much larger than the Taylor order.

5.5 An exact symmetric pole coordinate

We now evaluate the first windings accurately enough for comparison between adjacent admissible particle numbers. All assertions in this subsection hold for sufficiently large odd multiples of p. Put

$$d_n = \frac{\pi}{n}, \qquad \tau_n = 1 - \cos d_n, \qquad B_n = B(\tau_n), \qquad V_n = \{v \in \mathbb{C}^n : \sum_i v_i = 0\}, \tag{5.35}$$

and use the $w = 1$ chart $\delta_i = d_n + v_i$. Distinguish the holomorphic quadratic form from the norm used in estimates:

$$Q(v) = \sum_i v_i^2, \qquad q(v) = \sum_i |v_i|^2.$$

All transposes and gradients in holomorphic identities on V_n use the complex bilinear extension of its real Euclidean structure. The integration measure remains $\mathrm{d}v_1 \cdots \mathrm{d}v_{n-1}$, with $v_n = -\sum_{i<n} v_i$.

Define

$$c_n = \frac{\mathrm{i}B_n}{2s_\phi}, \qquad h_n = \cos d_n - \frac{B_n b}{s_\phi^2} \sin^2 d_n = 1 + O(n^{-2}). \tag{5.36}$$

Direct differentiation, keeping the actual function $B(t)$, gives

$$g(\tau_n, d_n) = \mathrm{i}\phi, \qquad g_t(\tau_n, d_n) = -2c_n, \qquad \frac{1}{2} g_{\delta\delta}(\tau_n, d_n) = c_n h_n. \tag{5.37}$$

For example, $W_t = B(t) + B'(t)(t - 1 + \cos\delta)$; the B' term vanishes at this center. Also $g_{\delta\delta} = W_{\delta\delta}/\sinh g - (\cosh g) W_\delta^2/\sinh^3 g$, which gives precisely (5.36).

The pole equation in this chart is

$$\sum_i \bigl(g(t, d_n + v_i) - \mathrm{i}\phi\bigr) = 0.$$

Its solution, denoted again by $T_n(v)$, is holomorphic and permutation invariant on a fixed ball $\|v\|_2 < r$, independent of n. Moreover

$$T_n(0) = \tau_n, \qquad |T_n(v) - \tau_n| \le \frac{Cq(v)}{n}, \qquad \|\nabla T_n(v)\|_2 \le \frac{C\sqrt{q(v)}}{n}, \qquad \|D^2 T_n(v)\| \le \frac{C}{n}. \tag{5.38}$$

Here and below operator norms are induced by the Euclidean norm. To prove these estimates, subtract the constant and linear terms of $g(t, d_n + v_i)$ in v_i. The linear terms sum to zero, and the remaining sum is bounded by $Cq(v)$. The t derivative of the sum is bounded below by cn, uniformly on this fixed neighborhood. Rouché's theorem, or the quantitative implicit-function theorem, gives the unique solution with $|T_n - \tau_n| \le Cq/n$. Uniqueness patches the local solutions over the whole ball. If $P = I - \mathbf{1}\mathbf{1}^T/n$, its gradient is

$$\nabla T_n = -\frac{P(g_\delta(T_n, d_n + v_i))_{i=1}^n}{\sum_i g_t(T_n, d_n + v_i)}.$$

The numerator has norm at most $C\|v\|_2$; another differentiation gives the Hessian bound. These arguments use fixed scalar derivative bounds for g, not a Cauchy radius that shrinks with the dimension.

Introduce removable divided differences

$$\mathfrak{a}_n(t) = \frac{g(t, d_n) - \mathrm{i}\phi}{t - \tau_n}, \tag{5.39}$$

$$\mathfrak{e}_n(t, z) = g(t, d_n + z) - g(t, d_n) - g_\delta(t, d_n) z, \tag{5.40}$$

$$\mathfrak{h}_n(t, z) = -\frac{2\mathfrak{e}_n(t, z)}{h_n \mathfrak{a}_n(t) z^2}. \tag{5.41}$$

The function $\mathfrak{a}_n$ is uniformly bounded away from zero. Evenness of g in its second variable implies

$$\mathfrak{h}_n(\tau_n, 0) = 1, \tag{5.42}$$

$$|\mathfrak{h}_n(t, z) - 1| \le C\bigl(|t - \tau_n| + |d_n||z| + |z|^2\bigr), \tag{5.43}$$

$$|\partial_z \mathfrak{h}_n(t, z)| \le C(|d_n| + |z|), \qquad |\partial_z^2 \mathfrak{h}_n| + |\partial_t \mathfrak{h}_n| \le C. \tag{5.44}$$

For clarity, Taylor's integral formula writes $\mathfrak{e}_n/z^2$ as an average of $g_{\delta\delta}(t, d_n + uz)$. Its first z derivative is an average of $g_{\delta\delta\delta}$, an odd function of its angular argument, and hence is $O(|d_n| + |z|)$. This proves the dimension-free bounds above.

Summing the divided differences at the pole gives the exact identity

$$T_n(v) - \tau_n = \frac{h_n}{2n} \sum_i v_i^2 \mathfrak{h}_n(T_n(v), v_i). \tag{5.45}$$

We turn its right-hand side into a quadratic form by an explicit permutation-equivariant change of variables. In particular, the construction never divides by $Q(v)$, which has a complex null cone.

First set

$$U_i(v) = v_i \sqrt{\mathfrak{h}_n(T_n(v), v_i)}, \qquad w = G(v) = PU(v), \tag{5.46}$$

using the square root near one. On a sufficiently small fixed ball,

$$\|DG - I\| \le C\big(|d_n| \|v\|_\infty + \|v\|_\infty^2 + q(v)/n\big) < \tfrac{1}{4}.$$

Thus G and its inverse are holomorphic on smaller fixed balls, with bounded derivatives. They commute with coordinate permutations. Because $T_n(v)$ is common to all components, one-variable divided differences give

$$\frac{U_i - U_k}{v_i - v_k} = 1 + \mathfrak{r}_{ik}(v), \tag{5.47}$$

$$|\mathfrak{r}_{ik}(v)| \le C\big(|T_n - \tau_n| + |d_n|(|v_i| + |v_k|) + |v_i|^2 + |v_k|^2\big). \tag{5.48}$$

The quotients extend holomorphically across $v_i = v_k$. Projection subtracts a common mean and does not change component differences. Summing logarithms of the near-one factors yields

$$\left|\log \frac{\Delta(G(v))}{\Delta(v)}\right| \le C\big(n^2 |T_n - \tau_n| + n|d_n| \sum_i |v_i| + nq(v)\big) \le C\big(\sqrt{nq(v)} + nq(v)\big). \tag{5.49}$$

The quotient in this formula means its holomorphic extension.

The Jacobian estimate has no extra dimension loss. The derivative of G is a compressed diagonal matrix plus a rank-one matrix. The trace norm of the diagonal perturbation is bounded by

$$C\big(n|T_n - \tau_n| + |d_n| \sum_i |v_i| + q(v)\big),$$

and that of the rank-one perturbation is at most $Cq(v)/n$. The convergent matrix logarithm therefore gives

$$|\log \det_{V_n} DG(v)| \le C\big(|d_n| \sqrt{nq(v)} + q(v)\big). \tag{5.50}$$

The determinant on V_n equals the change-of-variable determinant in the constrained coordinates, since the same constant linear coordinate transformation is used on the source and target.

The projection in (5.46) discarded a mean. In the w coordinates let

$$\mu(w) = \frac{1}{n} \sum_i U_i(G^{-1}(w)).$$

At fixed t,

$$U_i = A_0(t) v_i + A_1(t) v_i^2 + A_2(t, v_i) v_i^3, \qquad A_1(t) = O(d_n),$$

with uniformly bounded analytic coefficients. The first terms sum to zero. Differentiating this expansion and using (5.38) gives, with $q = q(w)$,

$$|\mu| \le \frac{C}{n}(|d_n| + \sqrt{q})q, \qquad \|\nabla\mu\| \le \frac{C}{n}(|d_n| + \sqrt{q})\sqrt{q}, \qquad \|D^2\mu\| \le \frac{C}{n}(|d_n| + \sqrt{q}). \tag{5.51}$$

For the last assertion, the second derivative of G^{-1} is bounded by $C(|d_n| + \|w\|_2)$, as follows directly by differentiating (5.46). The chain-rule terms therefore obey the same bound.

Define the holomorphic vector field

$$\ell(w) = \int_0^1 \nabla\mu(uw)\, \mathrm{d}u \in V_n. \tag{5.52}$$

Then $\mu(w) = w^T\ell(w)$, and

$$\|\ell(w)\| \le \frac{C}{n}(|d_n| + \sqrt{q})\sqrt{q}, \qquad \|D\ell(w)\| \le \frac{C}{n}(|d_n| + \sqrt{q}).$$

Equivariance under the transposition of coordinates i, k shows that

$$\ell_i(w) - \ell_k(w) = (w_i - w_k)L_{ik}(w), \qquad |L_{ik}(w)| \le \frac{C}{n}(|d_n| + \sqrt{q}). \tag{5.53}$$

Indeed compare w with the vector with these two entries exchanged and integrate the derivative along the segment. The segment remains in the ball. The corresponding one-variable divided difference also proves holomorphic extension at the coincidence hyperplane.

Now set

$$R(w) = \bigl(I + n\ell(w)\ell(w)^T\bigr)^{1/2} w = w + \frac{n\,\mu(w)\ell(w)}{1 + \sqrt{1 + n\ell(w)^T\ell(w)}}. \tag{5.54}$$

The matrix square root is the branch near the identity. Its perturbation has norm at most $C(|d_n| + \sqrt{q})^2 q/n$. The map is uniformly biholomorphic on a smaller fixed ball. Exact multiplication gives

$$Q(R(w)) = Q(w) + n\mu(w)^2 = \sum_i U_i(v)^2. \tag{5.55}$$

This identity uses bilinear transposes and remains valid on the complex null cone.

By (5.53), the relative change of each pair difference under R is at most $C(|d_n| + \sqrt{q})^2 q/n$. The total logarithmic cost for the Vandermonde is therefore at most Cnq. Differentiating (5.54) shows that the trace norm of $DR - I$ is at most $C(|d_n| + \sqrt{q})^2 q$. For example, the term involving $D\ell$ has operator norm $O((|d_n| + \sqrt{q})^2 q/n)$; its trace norm costs at most the dimension $n - 1$. All the other terms are rank one. This proves the same bound for the logarithmic Jacobian.

Combining the maps, define

$$\Psi_n(v) = \sqrt{h_n}\, R(G(v)). \tag{5.56}$$

It is a permutation-equivariant biholomorphism on fixed balls, $D\Psi_n(0) = \sqrt{h_n} I$, and it preserves all coincidence divisors. Equations (5.45) and (5.55) prove the exact normal form

$$T_n(v) = \tau_n + \frac{Q(\Psi_n(v))}{2n}. \tag{5.57}$$

After extracting the constant $h_n^{-\alpha_n}$, the Jacobian–Vandermonde ratio for its inverse has a holomorphic logarithm bounded by $C(\sqrt{nq} + nq)$. The exponent follows because the degree of Δ^2, plus the dimension of the plane, is $n(n-1) + (n-1) = 2\alpha_n$.

5.6 The complete residue density and its integration cycle

In the first-winding chart put

$$\varrho_n(v) = -\frac{2H_n(T_n(v), \pi + d_n + v)}{\partial_t \Gamma_n(T_n(v), \pi + d_n + v)}.$$

Here the arguments of H_n and Γ_n mean $q_i = \pi + d_n + v_i$, so that $\delta_i = d_n + v_i$. The quotient $\varrho_n/\Delta(v)^2$ is holomorphic and nonzero on a fixed small ball. Its value at zero is

$$\varrho_{n,0} = -\frac{2B_n^{\alpha_n - 1/2}}{n\, n!(2\pi)^{n-1} 2^{n(n-1)} s_\phi^{2\alpha_n}}. \tag{5.58}$$

To check its phase and sign, the numerator divided by Δ^2 at the center has the factor $(\mathrm{i}s_\phi)^{-n^2}$. Since n is odd, $\mathrm{i}^{-n^2} = -\mathrm{i}$, while $\partial_t \Gamma_n = -\mathrm{i}nB_n/s_\phi$. The minus sign in (5.58) also includes the minus sign from the Taylor expansion of $(t - T_n)^{-1}$.

In the coordinates $w = \Psi_n(v)$, the complete residue form is

$$\varrho_n(v)\, \mathrm{d}v = \varrho_{n,0} h_n^{-\alpha_n} \Delta(w)^2 (1 + \mathcal{E}_n(w))\, \mathrm{d}w, \qquad |\mathcal{E}_n(w)| \le e^{C(\sqrt{nq(w)} + nq(w))} - 1. \tag{5.59}$$

Here the error is a relative error, after dividing by the explicit center value. To prove it, take logarithms only after canceling Δ^2. The anisotropy ratio costs $Cn^2|T_n - \tau_n|$. Moreover

$$\sum_i |g(T_n, d_n + v_i) - \mathrm{i}\phi| \le C\big(|d_n| \sum_i |v_i| + q(v) + n|T_n - \tau_n|\big).$$

Pair denominators cost at most Cn times this sum; the one-body denominators cost only the sum itself. The sine divided differences cost $C\sum_{i<k} |v_i - v_k|^2 \le Cnq(v)$. Finally the linear terms cancel in the derivative sum, giving

$$\frac{\partial_t \Gamma_n(T_n(v), \pi + d_n + v)}{n g_t(\tau_n, d_n)} = 1 + O(q(v)/n).$$

Together with (5.38) and the Jacobian–Vandermonde estimates, these are exactly the logarithmic bound in (5.59).

We next justify replacing the original real cycle by a real cycle in the exact w coordinates. Choose a small fixed radius $a > 0$. For large n, the real ball $Q(v) < a^2$ lies strictly in the first-winding region $Q(v) < \sigma - \pi^2/n$. The omitted shell has $|T_n| \ge c/n$, by (5.26), and hence its contribution to the jth coefficient is bounded by

$$(Cn)^j \exp\{-\kappa n^2 + Cn \log(n+1) + C\}.$$

Use (5.15) for the numerator and absorb the volume and other factors into the indicated constants.

By choosing a small and then n large, one has $\|\Psi_n(v) - v\|_2 \le \eta \|v\|_2$, with a fixed $\eta < 1/4$, on a slightly larger ball. For real v, the straight homotopy $w_u = (1-u)v + u\Psi_n(v)$ satisfies

$$\Re Q(w_u) \ge (1 - 2\eta - \eta^2) Q(v) > 0 \quad (v \ne 0). \tag{5.60}$$

This follows by writing $w_u = v + e$, $\|e\| \le \eta\|v\|$, and estimating $2v^T e + e^T e$. The pole $\tau_n + Q(w)/(2n) = 0$ is therefore avoided throughout the homotopy.

Apply Stokes' theorem to the holomorphic $(n-1)$-form

$$\varrho_{n,0} h_n^{-\alpha_n} \frac{\Delta(w)^2 (1 + \mathcal{E}_n(w))}{(\tau_n + Q(w)/(2n))^{j+1}} \, \mathrm{d}w_1 \wedge \cdots \wedge \mathrm{d}w_{n-1}.$$

The difference between its integrals over the image of the real ball and over the real ball itself is the integral over the lateral boundary of the homotopy. On that boundary, (5.60) gives $|\tau_n + Q(w)/(2n)| \geq c/n$. The inverse image under Ψ_n remains in a fixed complex ball around the all-endpoint configuration. The complex-angle version of (5.15) bounds its complete numerator. The implicit derivative is at least cn; the Jacobians and the parameterized boundary volume cost at most $\exp(Cn \log(n+1) + C)$. For the latter bound it is enough to use bounded operator norms for the maps and their inverses and the elementary Euclidean ball and sphere volume bounds in dimension $n-1$. Consequently the entire boundary error has the bound

$$|b_{n,j}| \leq (Cn)^j \exp\{-\kappa n^2 + Cn \log(n+1) + C\}. \tag{5.61}$$

This argument estimates the exact residue form, not an approximate quadratic denominator.

5.7 A uniform beta evaluation of the first two windings

For $j + 1 > \alpha_n$, (5.32) yields

$$\int_{V_n \cap \mathbb{R}^n} \frac{\Delta(w)^2}{(\tau_n + Q(w)/(2n))^{j+1}} \, \mathrm{d}w = G_n (2n)^{\alpha_n} \tau_n^{\alpha_n - j - 1} \frac{\Gamma(j + 1 - \alpha_n)}{\Gamma(j+1)}. \tag{5.62}$$

The gamma quotient is retained exactly. Replacing it by $j^{-\alpha_n}$ would introduce a logarithmic error of order α_n^2/j, which is not uniformly small in the regime required below.

We prove that (5.59) has relative integrated error

$$O\left(\frac{n}{\sqrt{j}} + \frac{n^2}{j}\right) \quad \text{as } n \longrightarrow \infty, \quad n^2/j \longrightarrow 0. \tag{5.63}$$

Under the normalized positive measure in (5.62), the variable $x = Q/(2n\tau_n)$ has density proportional to

$$x^{\alpha_n - 1} (1 + x)^{-j-1} \, \mathrm{d}x.$$

Since $nQ = 2n^2 \tau_n x \asymp x$, the error in (5.59) is bounded by $e^{C(\sqrt{x} + x)} - 1$. On $0 \leq x \leq \delta$, for any fixed small $\delta > 0$, this is at most $C(\sqrt{x} + x)$. The exact mean

$$\mathbb{E}x = \frac{\alpha_n}{j - \alpha_n}$$

and Cauchy–Schwarz show that its integral is bounded by $C(\sqrt{\alpha_n/j} + \alpha_n/j)$.

It remains to estimate the tail, where the preceding exponential cannot be integrated over the whole half-line. The fixed local ball has $x \leq Ln$ for a constant L. Uniformly on $\delta \leq x \leq Ln$, when $n^2/j \to 0$,

$$C(\sqrt{x} + x) \leq \frac{j}{4} \log(1 + x).$$

Write $Z(\alpha, j) = \Gamma(\alpha)\Gamma(j + 1 - \alpha)/\Gamma(j+1)$. The tail integral divided by the original normalization is at most

$$(1 + \delta)^{-j/4} \frac{Z(\alpha_n, j/2)}{Z(\alpha_n, j)}.$$

Because α_n is an integer,

$$\frac{Z(\alpha_n, j/2)}{Z(\alpha_n, j)} = \prod_{r=0}^{\alpha_n - 1} \frac{j-r}{j/2-r} \le 4^{\alpha_n} \qquad (\alpha_n \le j/4).$$

Thus this tail is $\exp(-cj + O(\alpha_n))$. The same bound, without the exponential error factor, controls extending the main integral from the fixed real ball to the whole real plane. This proves (5.63) with constants uniform in the specified regime.

The winding $w = -1$ is identical to $w = 1$: replace all lifts by their negatives, use evenness of g, and use the squared sine factors. The ordinary real measure has the same orientation since $n - 1$ is even. The two leading terms therefore add. Substituting (5.58) and (5.30) into (5.62) gives the following exact-factor model:

$$A^{\mathrm{pole}}_{n,j,1} + A^{\mathrm{pole}}_{n,j,-1} = M(n,j)(1 + e_{n,j}) + b_{n,j}, \tag{5.64}$$

$$|e_{n,j}| \le C\left(\frac{n}{\sqrt{j}} + \frac{n^2}{j}\right), \tag{5.65}$$

$$M(n,j) = -\frac{2^{2-\alpha_n} B_n^{\alpha_n - 1/2} P_n n^{\alpha_n - 3/2}}{\pi^{(n-1)/2} s_\phi^{2\alpha_n}} \, h_n^{-\alpha_n} \tau_n^{\alpha_n - j - 1} \frac{\Gamma(j + 1 - \alpha_n)}{\Gamma(j+1)}. \tag{5.66}$$

The term $b_{n,j}$ satisfies (5.61). The power $B_n^{\alpha_n - 1/2}$, including its varying complex phase, is part of the model.

5.8 Exact finite differences and a single dominant index

Let

$$L(n,j) = \log|M(n,j)|, \qquad d = 2p.$$

The admissible indices are the odd multiples of p, so their spacing is d. Elementary Stirling bounds in (5.66) give, as $n \to \infty$ and $n^2/j \to 0$,

$$L(n,j) = 2j \log n - \frac{n^2}{2} \log j + c_* n^2 - j \log(\pi^2/2) + O(n \log n + n^4/j + j/n^2). \tag{5.67}$$

For example,

$$\log P_n = \frac{1}{2} n^2 \log n - \left(\frac{3}{4} + \frac{1}{2}\log 2\right) n^2 + O(n \log n), \qquad \log \tau_n = \log(\pi^2/2) - 2\log n + O(n^{-2}),$$

while the logarithm of the gamma quotient is $-\alpha_n \log j + O(\alpha_n^2/j)$. These formulas give exactly the value of c_* in (5.12).

We need finite differences of L, and do not obtain them by differentiating the error in (5.67). The exact factors give

$$L(n+d,j) - 2L(n,j) + L(n-d,j) = d^2\left(-\frac{2j}{n^2} - \log j + 2c_*\right) + O_d(n^2/j + j/n^4 + n^{-2}). \tag{5.68}$$

Here is a direct verification with the required uniform remainder. First,

$$\log P_{n+d} - 2\log P_n + \log P_{n-d} = \sum_{k=0}^{d-1} \sum_{\ell=1}^{d} \log(n - d + k + \ell) - d^2 \log 2 \tag{5.69}$$

$$= d^2(\log n - \log 2) + O_d(n^{-2}). \tag{5.70}$$

The offsets in this double sum have sum zero. Next, because α_n is an integer, for real $j > \alpha_{n+d}$ the exact identity is

$$\log \frac{\Gamma(j+1-\alpha_n)}{\Gamma(j+1)} = -\alpha_n \log j + \sum_{r=0}^{\alpha_n - 1} - \log(1 - r/j).$$

The upper and lower adjacent increments of α_n have lengths $dn + d^2/2$ and $dn - d^2/2$. Pair the $dn - d^2/2$ corresponding summands; their arguments differ by $O_d(n)$, and the derivative of $-\log(1 - x/j)$ is at most $1/(j - \alpha_{n+d})$. Their total difference is $O_d(n^2/j)$. The d^2 unpaired summands cost the same amount. Thus this factor contributes $-d^2 \log j + O_d(n^2/j)$.

For a precise accounting of all remaining factors, put

$$c_\tau = \pi^2/2, \qquad r_n = \log \tau_n - \log c_\tau + 2 \log n, \qquad K = -\tfrac{1}{2} \log 2 + \tfrac{1}{2} \log |B(0)| - \log s_\phi + \tfrac{1}{2} \log c_\tau.$$

Up to a constant independent of n, j, the logarithm $L(n, j)$ is exactly

$$\begin{aligned} &2j \log n - j \log c_\tau - \alpha_n \log j + \log P_n - \tfrac{1}{2} n^2 \log n + K n^2 - \tfrac{n}{2} \log \pi + \log n \\ &\quad + \sum_{r=0}^{\alpha_n - 1} - \log(1 - r/j) + (\alpha_n - \tfrac{1}{2}) \Re \log(B_n / B(0)) - \alpha_n \Re \log h_n + (\alpha_n - j - 1) r_n. \end{aligned}$$

The last three terms have convergent expansions in n^{-2}, and

$$r_n = -\frac{\pi^2}{12 n^2} + O(n^{-4}).$$

Their second differences are $O_d(n^{-2} + j/n^4)$, as is seen either term by term in these convergent expansions or by differentiating them on a fixed neighborhood of $n^{-1} = 0$. Also

$$\Delta_d^2(-\tfrac{1}{2} n^2 \log n) = -d^2(\log n + \tfrac{3}{2}) + O_d(n^{-2}), \qquad \Delta_d^2(2j \log n) = -\frac{2d^2 j}{n^2} + O_d(j/n^4).$$

Combine these with (5.70) and $2c_* = 2K - \frac{3}{2} - \log 2$. This proves (5.68) directly from the exact model.

In particular, throughout the intermediate range

$$\log j < n < \frac{\sqrt{j}}{(\log j)^{1/4}}, \tag{5.71}$$

the second difference is at most $-cd^2 \log j$ for all sufficiently large j, whenever the neighboring indices are also in that range. Indeed $n^2/j \to 0$ uniformly, and j/n^4 is smaller than j/n^2 by a factor at most $(\log j)^{-2}$. This is concavity over the entire intermediate range.

Fix a large admissible n, and temporarily let j be real of order $n^2 \log n$. Set

$$f_n(j) = L(n + d, j) - L(n - d, j).$$

The two gamma arguments differ by the integer $2dn$. Differentiating their exact product gives

$$f_n'(j) = \log \frac{\tau_{n-d}}{\tau_{n+d}} - \sum_{r=1}^{2dn} \frac{1}{j + 1 - \alpha_{n-d} - r} \tag{5.72}$$

$$= \frac{4d}{n} - \frac{2dn}{j} + O_d(n^{-3} + n^3/j^2) > 0. \tag{5.73}$$

Exact first differences also give

$$\frac{f_n(j)}{2d} = \frac{2j}{n} - n\log j + 2c_* n + O_d(\log n + n^3/j + j/n^3). \tag{5.74}$$

For detail, the quotient P_{n+d}/P_{n-d} is a product of $2d$ factorials; Stirling's bound on these individual factorials gives its centered first difference $n\log n - (1+\log 2)n + O_d(\log n)$. The centered first difference of the gamma correction sum is $O_d(n^3/j)$. The convergent n^{-2} expansions above contribute $O_d(n^{-3} + j/n^3)$. The displayed leading terms then follow exactly as in the second-difference calculation.

At

$$j = n^2\left(\log n + \tfrac{1}{2}\log\log n - c_* \pm \epsilon\right)$$

the expression in (5.74) has opposite signs for each fixed $\epsilon > 0$ and all large n. There is a unique crossing in this interval by (5.73). Round it to the nearest integer j_n. We obtain

$$j_n = n^2\left(\log n + \tfrac{1}{2}\log\log n - c_* + o(1)\right), \qquad f_n(j_n) = O_d(1/n). \tag{5.75}$$

These n lie strictly inside (5.71).

Let $\lambda_k = L(n+kd, j_n)$ for the admissible indices in that range. Their successive first differences decrease by at least $cd^2\log j_n$. Equation (5.75) gives $\lambda_1 - \lambda_{-1} = O_d(1/n)$. Consequently

$$\lambda_0 - \lambda_{\pm 1} \geq c' d^2 \log j_n, \qquad \lambda_k - \lambda_0 \leq -c'' d^2 k^2 \log j_n + O_d(|k|/n) \quad (k \neq 0).$$

Summing this discrete Gaussian proves

$$\sum_{\substack{m\neq n,\ m \text{ odd},\ p\mid m \\ m \text{ satisfies (5.71)}}} |M(m, j_n)| \leq |M(n, j_n)|\, O(j_n^{-c''' d^2}). \tag{5.76}$$

The complex phases of the different models play no role in this inequality.

5.9 Exhausting the remaining indices and windings

We compare every part of (5.19) with the selected model. First, (5.67) and (5.75) imply

$$\log|M(n, j_n)| = j_n\log j_n - j_n\log\log j_n + j_n\big(\log(4/\pi^2) - 1\big) + o(j_n), \tag{5.77}$$

$$|M(n, j_n)|^{1/j_n} \sim \frac{4j_n}{e\pi^2\log j_n}. \tag{5.78}$$

Indeed (5.75) gives

$$n^2 = \frac{j_n}{\frac{1}{2}\log j_n - c_* + o(1)}.$$

Substituting this in the four leading terms of (5.67) gives (5.77); each error there is $o(j_n)$.

The following estimates cover all the complementary terms.

Small particle indices. For $m \leq \log j$, the full coefficient bound (5.18) gives

$$\sum_{\substack{m\leq \log j \\ m \text{ odd}}} |[t^j]C_m(t)| \leq \exp\{2j\log\log(j+2) + Cj + C\}. \tag{5.79}$$

This is negligible compared with (5.77) when $j = j_n$. Equivalently, after the pole reduction one can apply the same bound to the poles plus their analytic remainders, since their difference has the smaller bound (5.25).

All analytic and cycle remainders. The sum of the regular remainders in (5.25) and all the terms (5.61) is bounded by

$$\exp\{\tfrac{1}{2}j\log(j+2)+Cj+C\}. \tag{5.80}$$

This too is negligible at the selected orders. This estimate includes the entire nonresonant particle class $p \nmid m$ outside the small-index range.

All sufficiently large particle indices. For

$$m \geq \frac{\sqrt{j}}{(\log j)^{1/4}},$$

use (5.34), which has no restriction on m^2/j. With $x = m^2$, its exponent is at most

$$j\log x - \frac{1}{4}x\log x + Cx + Cj + C.$$

Its derivative in x is negative on $x \geq j/\sqrt{\log j}$, for all large j, and at the lower endpoint its value is

$$j\log j - \frac{1}{2}j\log\log j - \frac{1}{4}j\sqrt{\log j} + o(j\sqrt{\log j}). \tag{5.81}$$

For example its derivative is at most $-c\log j$ there for large j, since $j/x \leq \sqrt{\log j}$ and $\frac{1}{4}\log x \sim \frac{1}{4}\log j$. Summing the resulting decreasing exponential over the discrete values $x = m^2$ costs at most a harmless polynomial factor. Thus (5.81) bounds the logarithm of the whole infinite large-index tail up to $o(j\sqrt{\log j})$. Its gap below (5.77) tends to infinity faster than j.

Higher odd windings in the intermediate range. Let $M_0(m,j) > 0$ be the modulus of the expression (5.66) with B_m replaced by $B(0)$, h_m by 1, and τ_m by $\pi^2/(2m^2)$. The exact factors give

$$\left|\log\frac{M_0(m,j)}{|M(m,j)|}\right| \leq C(1 + j/m^2). \tag{5.82}$$

For the τ factor this follows from $\log(\tau_m/(\pi^2/(2m^2))) = O(m^{-2})$; the B and h factors cost $O(1)$. Comparison of (5.33) with this explicit M_0 yields, for each odd $|w| \geq 3$,

$$|A^{\mathrm{pole}}_{m,j,w}| \leq CM_0(m,j)e^{Cm\sigma}(1-C\sigma)^{-j-1}|w|^{-2(j+1-\alpha_m)}.$$

The constants C here do not depend on m, j, w. Choose the fixed σ still smaller, if necessary, so that $-\log(1-C\sigma) < \frac{1}{2}\log 3$. Throughout (5.71),

$$\alpha_m = o(j), \qquad m = o(j), \qquad j/m^2 = o(j)$$

uniformly. The number of possible windings is at most $C\sqrt{m}$. Hence for some $c > 0$,

$$\sum_{\substack{w \text{ odd} \\ |w|\geq 3}} |A^{\mathrm{pole}}_{m,j,w}| \leq |M(m,j)|e^{-cj} \tag{5.83}$$

uniformly in that range. The choice of a smaller σ merely changes the fixed constants in the earlier arguments.

The first windings of every intermediate index. Equation (5.65) tends uniformly to zero in (5.71); it is at most $C(\log j)^{-1/4}$. At $j = j_n$, (5.76) shows that the sum of the moduli of all intermediate models other than $M(n, j_n)$ is $o(|M(n, j_n)|)$. The sum of all their relative errors is bounded by

$$o(1) \sum_m |M(m, j_n)| = o(|M(n, j_n)|).$$

The same is true of all higher windings by (5.83).

These classes exhaust every odd particle number and every winding, together with all analytic and cycle remainders. Combining them in (5.19) proves the actual full-sum coefficient asymptotic

$$a_{j_n} = M(n, j_n)(1 + o(1)). \tag{5.84}$$

In particular, this is a statement about the sum of the complex coefficients, not merely a comparison of individual singularity sets.

5.10 Nonextension and uniformity

Equations (5.78) and (5.84) prove (5.11), so $\limsup_j |a_j|^{1/j} = \infty$. If F admitted a holomorphic extension to a disk about zero from the physical side $\mathcal{U}^-$, its derivatives on the inward negative ray would agree with those of that extension. By (5.9), its Taylor coefficients at zero would therefore be the a_j. Cauchy's estimate on any smaller extension disk would bound $|a_j| \le CR^{-j}$, a contradiction. If $P(0) \neq 0$, a holomorphic extension of PF would, after division by the nonvanishing holomorphic factor P, give an extension of F. Thus a local physical normalization by such a factor cannot remove the obstruction.

Finally, all the constants used above arise from a common holomorphic neighborhood for B, a lower bound for $-\Im B(0)$, and fixed scalar bounds for the dispersion and its derivatives. A holomorphic family with $\Im B(0, \eta) \le -\beta < 0$ on a relatively compact parameter polydisk supplies these uniformly by Cauchy's estimate. The square roots and logarithms near one used in the normal form also have common neighborhoods. The exact coefficient model continues to use the actual values $B(\tau_n, \eta)$; nothing in the argument freezes the anisotropy. The crossing argument and all range estimates are uniform on each such compact family, with $c_* = c_*(\eta)$ and the selected orders $j_n(\eta)$. This completes the proof of Theorem 5.1.

6 The isotropic restriction: a unified even–odd proof

This section proves the isotropic restriction directly from the complete form-factor integrals. In particular, singularity of a function of two variables is not used to infer singularity on the isotropic line. The particle parity is the only phase-dependent ingredient in the analytic estimates below.

Write ζ for the auxiliary exterior variable, and put

$$E(\zeta) = \zeta + \zeta^{-1}, \qquad D(x, y; \zeta) = E(\zeta) - \tfrac{1}{2}(x + x^{-1} + y + y^{-1}).$$

For the low-temperature restriction $\zeta = s$; for the high-temperature restriction $\zeta = s^{-1}$. This is a change of variable, not a duality identification between the two susceptibilities. Define

$$T_N^\sigma(\zeta) = \frac{1}{N!} \oint \cdots \oint \frac{Z^{-1} + Y^{-1}}{(1 - \sigma Z)(1 - \sigma Y)} \prod_{i<j} P_{ij} \prod_i R(y_i; \zeta) \frac{\mathrm{d}y_i}{2\pi \mathrm{i}}, \tag{6.1}$$

$$f_N(\zeta) = \frac{1}{N!} \oint \cdots \oint (ZY)^{-1} \prod_{i<j} P_{ij} \prod_i R(y_i; \zeta) \frac{\mathrm{d}y_i}{2\pi \mathrm{i}}, \tag{6.2}$$

where $\sigma \in \{1,-1\}$, $Y = \prod_i y_i$, $Z = \prod_i z_i$, and

$$W = E(\zeta) - \tfrac{1}{2}(y + y^{-1}), \quad z = W - \sqrt{W^2 - 1}, \quad R = \frac{2z^2}{1-z^2}, \quad P_{ij} = -\frac{(y_i - y_j)^2 z_i z_j}{y_i y_j (1 - z_i z_j)^2}.$$

On real $\zeta > 1$ the root satisfies $0 < z < 1$ on the angular circle; elsewhere it is the continued interior root. Put $T_N = T_N^1$ and $C_N = f_N + 2T_N$.

These expressions retain the full-site normalization. Their double-contour lift, with $X = \prod_i x_i$, is

$$T_N^\sigma = \frac{1}{N!} \oint \cdots \oint \frac{X^{-1} + Y^{-1}}{(1-\sigma X)(1-\sigma Y)} \prod_{i<j} \frac{x_i - x_j}{1 - x_i x_j} \frac{y_i - y_j}{1 - y_i y_j} \prod_i \frac{\mathrm{d}x_i \, \mathrm{d}y_i}{(2\pi \mathrm{i})^2 D(x_i, y_i; \zeta)}. \tag{6.3}$$

For f_N replace the first rational factor by $(XY)^{-1}$. Choose a common radius $r < 1$ with $|z(y;\zeta)| < r$ on the y circle. The only enclosed dispersion pole is then $x = z$, with residue $2z^2/(1-z^2)$. At $x = 0$, $D^{-1} = -2x + O(x^2)$ cancels the apparent numerator pole. All pair and product poles remain outside after each successive residue. Subtracting two dispersion equations gives

$$(z_i - z_j)(1 - z_i z_j) = -\frac{z_i z_j}{y_i y_j}(y_i - y_j)(1 - y_i y_j),$$

and hence the canceled expression for P_{ij}. The full lattice multiplier is exactly

$$\frac{(1+X)(1+Y)}{XY(1-X)(1-Y)} = \frac{1}{XY} + \frac{2(X^{-1} + Y^{-1})}{(1-X)(1-Y)}. \tag{6.4}$$

Thus f_N is the onsite coefficient, T_N is the indicated half of the offsite coefficient, and C_N is the full lattice coefficient. Equivalently,

$$C_N(\zeta) = \frac{1}{N!} \int_{\sum q_i = 0 \ (2\pi)} \prod_i \frac{1}{\sinh \gamma_i} \frac{1+Z}{1-Z} \prod_{i<j} \left[\frac{\sin((q_i - q_j)/2)}{\sinh((\gamma_i + \gamma_j)/2)} \right]^2 \prod_{i<N} \frac{\mathrm{d}q_i}{2\pi}, \tag{6.5}$$

where $\cosh \gamma_i = E(\zeta) - \cos q_i$ and $z_i = e^{-\gamma_i}$. The independent measure has $N-1$ coordinates, with $q_N = -\sum_{i<N} q_i$ modulo 2π. For completeness, this equality can be checked before any boundary limit. On real $\zeta > 1$ let $q_N(m,n)$ be $1/N!$ times the N-angle integral with multiplier $Z^m Y^n$, measure $\prod_i \mathrm{d}q_i/(2\pi \sinh \gamma_i)$, and the displayed squared pairs. Angle reversal gives $q_N(m,n) = q_N(m,-n)$; exchanging the two coordinate sets in (6.3) gives $q_N(m,n) = q_N(n,m)$ for $m, n \geq 0$. Any permutation sign of the top form is canceled by the same permutation of the product orientation, for either parity. A common small angle strip has $\Re \gamma_i \geq c > 0$, so simultaneous angle shifts give, at fixed N, ζ, a bound $C_{N,\zeta} e^{-N(cm + \eta|n|)}$. Consequently the spatial sum is absolutely convergent. Summing $n \in \mathbb{Z}$ enforces $\sum q_i = 0$ modulo 2π, and summing the reflected m coordinate gives $(1+Z)/(1-Z)$. Alternatively,

$$\frac{Y+Z}{(1-Y)(1-Z)} = \sum_{m \geq 1, n \geq 0} Y^m Z^n + \sum_{m \geq 0, n \geq 1} Y^m Z^n$$

counts each positive axis once and each open positive quadrant twice. Thus $f_N + 2T_N$ counts exactly every lattice site, establishing (6.5) without omitting the onsite term.

6.1 Compact convergence and a dense arithmetic family

For even N the Schur identity states that

$$\operatorname{Pf}\left[\frac{v_i - v_j}{1 - v_i v_j}\right] = \prod_{i<j} \frac{v_i - v_j}{1 - v_i v_j}.$$

For odd N, append $v_{N+1} = -1$; its last upper column is the removable constant one, and the same identity gives the product over the N real indices. One proof puts $a_i = (1+v_i)/(1-v_i)$ and compares residues at $a_N = -a_i$ inductively. Both rational functions have the same residues, are bounded at infinity, and vanish at $a_N = a_{N-1}$.

On a compact exterior neighborhood choose fixed radii $r < 1$ with all interior roots bounded by $q < r$ and bounded away from zero. The global inverse products cost only C^N. Hadamard's inequality gives $|\operatorname{Pf} A| \le C^{m/2}(m-1)^{m/4}$ for an even matrix of size m. Applying this to both matrices, including the ghost index if necessary, gives

$$|T_N^\sigma| + |f_N| + |C_N| \le C_0 \frac{C_1^N N^{(N+1)/2}}{N!}. \tag{6.6}$$

The bound is summable. Cauchy's estimate on a larger compact neighborhood proves local normal convergence of every fixed derivative. The constants in this compact statement are not asserted uniform at the unit circle.

Lemma 6.1 (The selected first orders)**.** *Let $p \ge 11$ be prime, choose distinct $a, b \in \{1, \dots, (p-1)/2\}$, and set*

$$\alpha = 2\pi a/p, \quad \beta = 2\pi b/p, \quad E_* = \cos\alpha + \cos\beta = 2\cos\theta, \quad \zeta_* = e^{\mathrm{i}\theta}.$$

For the even family choose $\alpha, \beta \in (0, \pi/2)$ and put $N_0 = 2p$, $k = N_0^2/2 - 1$. For the odd family choose both angles in $(0, \pi/2)$ or both in $(\pi/2, \pi)$ and put $N_0 = p$, $k = (p^2-1)/2$. The first order of the respective parity satisfying a Nickel equation is N_0, and its unordered cosine pair is uniquely $\{\cos\alpha, \cos\beta\}$. No endpoint configuration is resonant at any order. The even family is dense in the upper-right arc; the odd families are dense in both upper quadrants.

Proof. Let $\omega = e^{2\pi\mathrm{i}/p}$, $d = (p-1)/2$, $u_j = \omega^j + \omega^{-j}$, and $F = \mathbb{Q}(\omega + \omega^{-1})$. Every nonzero-index u_j has trace -1. Thus $v = u_a + u_b$ has trace -2 and is irrational: if rational it would be an integer, whereas its trace would force $v = -2/d$. If $p \nmid n$, then $\mathbb{Q}(\omega) \cap \mathbb{Q}(e^{2\pi\mathrm{i}/n}) = \mathbb{Q}$. Indeed the compositum has degree $\varphi(pn) = \varphi(p)\varphi(n)$, so the intersection has degree one. A lower order of the required parity would contradict this fact.

The only rational relation between $1, u_1, \dots, u_d$ is a multiple of $1 + \sum_j u_j = 0$: lift the relation to a polynomial of degree at most $p-1$ and use the minimal polynomial $1 + X + \cdots + X^{p-1}$. At order p all cosines are $u_j/2$ or 1; at order $2p$ they are $\pm u_j/2$ or ± 1. An equality of two cosine pairs has at most four nonconstant positions. Since $d \ge 5$, a zero coefficient forces the multiple of the relation to vanish, proving uniqueness.

For $E_* > 0$, an endpoint resonance requires a root of unity ξ with $\xi + \xi^{-1} = v - 2$. Its trace is $-2 - 2d < -2d$, impossible because every conjugate of a root-of-unity trace belongs to $[-2, 2]$. For $E_* < 0$ the possible endpoint equation is $\xi + \xi^{-1} = v + 2$. Direct expansion gives $\operatorname{Tr}_F(v^2) = 4d - 6$. If all conjugates of $v + 2$ were in $[-2, 2]$, every conjugate of v would be in $[-4, 0]$, where $v^2 \le -4v$. Taking traces would imply $4d - 6 \le 8$, contrary to $d \ge 5$. Approximating any angle in either relevant half interval by two distinct adjacent prime-grid angles proves density. □

Lemma 6.2 (Exponential endpoint separation)**.** *If a fixed algebraic unit-modulus number ξ is not a root of unity, there is $A > 0$ such that $|1 - \xi^N|, |1 + \xi^N| \ge e^{-AN}$ for every positive integer N.*

Proof. Write its primitive irreducible polynomial as $a_d \prod_{\nu=1}^{d}(X-\xi_\nu)$ with $\xi_1=\xi$. Its nonzero integer resultant with X^N-1 has modulus at least one. Bounding every other conjugate by $|\xi_\nu^N-1| \le 2\max(1,|\xi_\nu|)^N$ proves the first assertion. The inequality $|1+\xi^N| \ge |1-\xi^{2N}|/2$ proves the second after increasing A. □

6.2 The first complete singular coefficient

Put $\zeta_\varepsilon=(1+\varepsilon)\zeta_*$ and keep the integration radius $r=e^{-c_0\varepsilon}$ fixed during every complex derivative. On a sufficiently small disk of radius $c\varepsilon$, $\Im W \ge c'\varepsilon$ for every angle. The interior roots belong to the lower unit half-disk and satisfy $|z|<r$ if c_0 is small enough. These statements hold for either upper quadrant.

Lemma 6.3 (Fixed-order complements)**.** *At each point in Lemma 6.1, every fixed derivative of each lower coefficient of the selected parity is bounded on the exterior radial approach. At order N_0, the complement of the two lower double-contour configurations*

$$(x_i,y_i)=(e^{-\mathrm{i}\beta},e^{-\mathrm{i}\alpha}) \quad \textit{for all } i, \qquad \textit{and their exchange}$$

has bounded derivatives of every fixed order. The onsite coefficient has bounded fixed derivatives for every fixed N.

Proof. The potentially singular factors on the unit double torus have gradient vectors consisting of the two positive all-ones product vectors, the positive reciprocal-pair vectors, and the dispersion vectors with entries $(\Im x_i, \Im y_i)$. The sign of dispersion damping is fixed by $\Im E(\zeta_\varepsilon)>0$. A nonnegative zero combination cannot contain a pair vector: its two positive entries would require both corresponding imaginary parts to be negative, whereas reciprocal circle coordinates have opposite imaginary parts.

If both product vectors occur, every coordinate is lower and the ratios $\sin A/\sin B$ agree along $\cos A+\cos B=E_*$. Along that curve

$$\frac{\mathrm{d}}{\mathrm{d}A}\frac{\sin A}{\sin B}=\frac{E_*(1-\cos A\cos B)}{\sin^3 B}$$

has a strict constant sign. All pairs therefore coincide, and both common coordinates are Nth roots of unity. Lemma 6.1 leaves exactly the specified configurations at N_0 and none below it. With only one product vector, the other coordinates must all be $+1$ when $E_*>0$, or all -1 when $E_*<0$; this is the excluded endpoint resonance. With neither product vector, a nonzero dispersion vector cannot cancel because $E_*\notin\{-2,0,2\}$.

On the complement, a finite smooth partition supplies a real direction strictly separating all active vectors from zero on each chart. Write its ℓ singular factors as damped positive-half-line exponential integrals, and put $R=\sum_{i=1}^{\ell}\xi_i$. The exponent g has $\Re g\le 0$, $|e\cdot\nabla g|\ge cR$, and fixed angular jets $O(R)$. A fixed jth parameter derivative costs $O(R^j)$. Integrating by parts $\ell+j+1$ times with $(e\cdot\nabla)/(e\cdot\nabla g)$ leaves radial majorant CR^{-2} for $R\ge 1$, including all coefficient and cutoff derivatives. The small-R part is bounded directly. Omitting both product factors proves the onsite assertion. This argument is at fixed N,j only. □

Proposition 6.4 (Nonzero first amplitude)**.** *There is $L_*\neq 0$ such that*

$$T_{N_0}^{(k)}(\zeta_\varepsilon)=\begin{cases}2L_*\varepsilon^{-1/2}+o(\varepsilon^{-1/2}), & N_0 \textit{ even},\\ 2L_*\varepsilon^{-1}+o(\varepsilon^{-1}), & N_0 \textit{ odd}.\end{cases}$$

For even N_0 all lower derivatives are bounded. For odd N_0 the $(k-1)$st derivative is $O(1+|\log\varepsilon|)$ and all smaller derivatives are bounded. The corresponding leading coefficient for C_{N_0} is $4L_*$.

Proof. Use fixed smooth weights depending only on the y angles around the two specified configurations, with disjoint supports. All x integrations remain unweighted, so the residues giving (6.1) apply to each chart exactly. Lemma 6.3 controls the full complementary double integral.

In the first chart write $y_i = re^{\mathrm{i}(-\alpha+u_i)}$, $v = \sum_i u_i$, $u_i = v/N + \eta_i$, $\sum_i \eta_i = 0$. The absolute real Jacobian in coordinates $(v, \eta_1, \dots, \eta_{N-1})$ is one. We use positive real product measures here; equivalently, the induced domain orientation compensates the determinant $(-1)^{N-1}$ of this coordinate change. Write $z_i = e^{-\mathrm{i}\phi_i}$. At the center,

$$b = \phi_u = \frac{\sin\alpha}{\sin\beta} > 0, \quad d = -\tfrac{1}{2}\phi_{uu} = \frac{H_{\alpha\beta}}{2\sin^3\beta} \neq 0, \qquad H_{\alpha\beta} = E_*(1 - \cos\alpha\cos\beta).$$

Also

$$\phi = \beta - \mathrm{i}A_0\varepsilon + bu - du^2 + O(\varepsilon^2 + \varepsilon|u| + |u|^3), \quad A_0 = \frac{2\sin\theta - c_0\sin\alpha}{\sin\beta} > c_0.$$

Replace the smooth mean cutoff by a fixed hard segment; its difference has separated product phases if the shape radius is sufficiently small. Move that segment downward in a fixed rectangle. The $Y = 1$ pole is at $v = -\mathrm{i}Nc_0\varepsilon$; downward motion increases the damping of Z. The vertical sides have a fixed phase gap and the bottom fixed damping, so all their fixed derivatives are bounded. The clockwise closure and $\partial_v(1 - Y) = -\mathrm{i}$ give the actual residue multiplier $+2\pi$. No nonholomorphic weight is moved through a complex contour.

At the residue the y_i lie on the unit circle and

$$1 - Z = Q\varepsilon - \mathrm{i}d\sum_i \eta_i^2 + O(\varepsilon^2 + \varepsilon|\eta| + |\eta|^3), \quad Q = \frac{2N\sin\theta}{\sin\beta} > 0, \tag{6.7}$$

with nonlinear bound $|1 - Z| \geq c(\varepsilon + \sum_i \eta_i^2)$. To see this bound, modulus damping is at least $c\varepsilon$; permutation symmetry cancels the linear shape term, and the real phase Hessian has a strict sign determined by $-H_{\alpha\beta}$. Its integral remainder controls the phase outside $\sum \eta_i^2 = O(\varepsilon^2)$, while damping controls the remaining region.

The exact numerator is $\Delta(\eta)^2 A(\zeta, \eta)$ with A analytic and nonzero at the center. Fixed parameter derivatives preserve the Vandermonde. Each pair contributes $-(\eta_i - \eta_j)^2/(4\sin^2\beta)$, and each one-body angle measure contributes $y_0 z_0/(2\pi\mathrm{i}\sin\beta)$. Since $y_0^N z_0^N = 1$, their combined phase is $(-1)^{N(N-1)/2}\mathrm{i}^{-N} = \mathrm{i}^{-N^2}$. After the actual mean residue the leading constrained value is

$$\frac{2\mathrm{i}^{-N^2}}{N!(2\pi)^{N-1}2^{N(N-1)}\sin^{N^2}\beta} \frac{\Delta(\eta)^2\ \mathrm{d}\eta_1 \cdots \mathrm{d}\eta_{N-1}}{Q\varepsilon - \mathrm{i}d\sum_i \eta_i^2}.$$

After k derivatives and $\eta = \sqrt{\varepsilon}\, w$, only the top pole power contributes at scale $\varepsilon^{-1/2}$ for even N and ε^{-1} for odd N. The radial majorants have tails R^{-2} and R^{-3} respectively. Lower pole powers vanish after multiplication by the indicated singularity scale; splitting at $R = \sqrt{\varepsilon}$ gives the logarithm at the odd lower derivative. This is dominated convergence after the mean residue, not of the unintegrated absolute mean–shape integral.

For $a = (N^2 - 1)/2$, let

$$c_N = \frac{1}{\sqrt{N}} \int_{\sum \omega_i = 0,\ |\omega| = 1} \Delta(\omega)^2\ \mathrm{d}\omega > 0.$$

The remaining radial period is

$$\int \frac{\Delta(w)^2\ \mathrm{d}w_1 \cdots \mathrm{d}w_{N-1}}{(Q - \mathrm{i}d\sum_i w_i^2)^{k+1}} = \frac{c_N}{2} Q^{a-k-1}(-\mathrm{i}d)^{-a} B(a, k + 1 - a),$$

where the quadratic coefficient is continued from positive real part. Here $k+1-a$ equals $1/2$ or 1, and every factor is nonzero. For odd $N=p$, $a=k\equiv 0 \pmod 4$, so $d^k=|d|^k$ also on the negative-E_* family. The pole derivative is $Qe^{-\mathrm{i}\theta}$. In particular the complete one-chart coefficient is

$$L_* = \frac{\mathrm{i}^{-N^2}(-1)^k k!\, c_N B(a,k+1-a)\, 2^a (2N\sin\theta)^{a-1} e^{-\mathrm{i}k\theta}}{N!(2\pi)^{N-1} 2^{N(N-1)} (-\mathrm{i}H_{\alpha\beta})^a}. \tag{6.8}$$

Indeed the sine exponent is $-N^2-(a-1)+3a=0$. For even N, $H_{\alpha\beta}>0$ and the last power uses argument $-\pi/2$. For odd N, $a=k$ is divisible by four, so that power is $|H_{\alpha\beta}|^k>0$ for either sign of E_*. The factor i^{-N^2} equals one for even N and $-\mathrm{i}$ for odd N. Every remaining factor is symmetric in α,β. This checks the equality of the two complex coefficients, including orientation and phase, rather than just equality of their moduli. The two charts therefore add. Finally $C_N=f_N+2T_N$ and Lemma 6.3 give the factor four. □

6.3 A uniform boundary-tail theorem

Negating (ζ,y,z) gives, directly from the forms and their measures,

$$T_N^{\sigma}(-\zeta) = T_N^{\sigma(-1)^N}(\zeta), \qquad f_N(-\zeta) = (-1)^N f_N(\zeta). \tag{6.9}$$

Conjugation follows from the real germ. Consequently it suffices to estimate either signed kernel at a selected point with $0<E_*<2$. Set $b_*=\arccos(E_*-1)$ and $\xi=e^{-\mathrm{i}b_*}$. Lemma 6.1, also applied after the reflection of the negative family, and Lemma 6.2 give the needed $|1\pm\xi^N|\ge e^{-AN}$.

Theorem 6.5 (The entire differentiated tail)**.** *For either parity, its selected first order N_0, and every fixed $0\le j\le k$,*

$$\sum_{\substack{N>N_0\\ N \text{ of the selected parity}}} |T_N^{(j)}(\zeta_\varepsilon)| = o(\varepsilon^{-1/2}), \qquad \sum_{\substack{N>N_0\\ N \text{ of the selected parity}}} |C_N^{(j)}(\zeta_\varepsilon)| = o(\varepsilon^{-1/2}).$$

The estimates include all particle orders and both upper quadrants needed by the odd family.

The rest of this subsection proves the theorem, including the parameter disks, real-domain differentiation, and coarea estimates it requires. Fix the point and k first. Put

$$H=\log(1/\varepsilon), \qquad 0<\alpha_0<\frac{1}{2(k+2)}, \qquad \lambda_*=\varepsilon^{\alpha_0}.$$

All contour functions below are fixed smooth periodic functions, with bounded jets of every order required through k.

Choose m supported near the lower branch angle $-b_*$ and equal to one on a neighborhood of every lower branch-cutoff support. Choose a supported in $[\delta,\pi-\delta]$, equal to one on $[2\delta,\pi-2\delta]$, and choose p supported in $(0,\pi)$ and equal to one on a neighborhood of $\operatorname{supp} a$. The upper inverse branch is inside $a=p=1$, and the supports of m,p are disjoint. For a small fixed $\tau>0$ define

$$P=\sum_i p(\theta_i), \quad h_i=-2\tau p(\theta_i)+\frac{\tau P}{2N}m(\theta_i), \quad y_i(\lambda,\theta)=re^{\mathrm{i}\theta_i+\lambda h_i}, \quad \psi=\prod_i(1-a(\theta_i)).$$

Write $U_{\lambda,\zeta}$ for the pullback of the complete normalized form (6.1), including $1/N!$.

Lemma 6.6 (Exact weighted deformation)**.** *For every N and either σ,*

$$T_N^\sigma = F_N + K_N + S_N,$$
$$F_N = \int (1-\psi)U_{1,\zeta}, \qquad K_N = \int \psi U_{0,\zeta},$$
$$S_N = -2\mathrm{i}\tau \int_0^1 \sum_q \int (\partial_{\theta_q}\psi) U_{\lambda,\zeta}\, \mathrm{d}\lambda.$$

Every current term has one named upper index q with $p_q = 1$, $p'_q = m_q = m'_q = 0$; all other indices are in $\operatorname{supp}(1-a)$.

Proof. The total radial shift satisfies $\sum h_i \le -3\tau P/2$, so $|Y| < 1$. Moreover,

$$\partial_{\log|y|} \Im W = -\cosh(\log|y|)\sin\theta.$$

Both inward upper motion and outward lower motion increase $\Im W$. All roots remain interior; the product and canceled-pair denominators remain nonzero throughout the closed homotopy.

After dividing row i of the angular Jacobian by y_i, its matrix is

$$M_{ij} = d_i\delta_{ij} + \frac{\lambda\tau}{2N} m_i p'_j, \quad d_i = \mathrm{i} + \lambda\left(-2\tau p'_i + \frac{\tau P}{2N} m'_i\right).$$

Since $m_i p'_i = 0$, its determinant is $\prod_i d_i$. At a named current index its row and column are diagonal with $d_q = \mathrm{i}$; the homotopy replacement column multiplies the determinant by $2\mathrm{i}\tau$. Cartan's formula on the closed torus gives $\partial_\lambda \int \psi U_\lambda = -2\mathrm{i}\tau \sum_q \int (\partial_q\psi) U_\lambda$. Combine this with invariance of the unweighted closed integral. □

Support geometry and pair suppression. Here and below "branch" means the lower branch, never its excluded upper inverse. Choose nested branch cutoffs: an outer one supported in $|u| \le \delta_B$, equal to one on $|u| \le \delta_B/2$, and an inner one supported in $|u| \le 2r_0$, equal to one on $|u| \le r_0$, where $u = \theta + b_*$ and $r_0 = \delta_B/(2L^2)$ with L sufficiently large. All these supports lie in $m = 1$. Telescope the product of the outer cutoffs. A complementary term has an outer compact index with $|u_q| \ge \delta_B/2$, including all cutoff jets. Split the other coordinates by the inner cutoff and compact left/right labels. The latter can be made periodic by returning their transition inside the upper $a = 1$ plateau, where the full K and current weights vanish with all jets. Thus no cutoff differentiation introduces a new support type.

The partition has at most C^N labels. Its types are: all branch; a compact left root with a modulus gap; a separated compact/true-branch pair; and all compact right. A named current has no all-branch sector. If it has a branch index, choose the named upper index and a fixed branch index as the separated pair. The nested widths ensure that the branch slope dominates the compact slope by at least four; preliminary upper endpoint neighborhoods fix the latter bound before their final shrinking.

For clarity the quadratic pair estimate used throughout is proved next. At $\varepsilon = \lambda = 0$, lower angles lie in $[-\pi, 0]$; their roots are real in $[z_0, 1]$ on $[-\pi, -b_*]$ and are $e^{-\mathrm{i}v}$, $0 \le v \le b_*$, on $[-b_*, 0]$, where $z_0 = E_* + 1 - \sqrt{(E_*+1)^2 - 1} > 0$. The identity

$$|1 - vw|^2 - |v - w|^2 = (1 - |v|^2)(1 - |w|^2) + 4\Im v \Im w \tag{6.10}$$

and its angular version imply $|P_{ij}| \le 1$. Equality is possible only for $y_i = 1$ paired with a left angle, or $y_i = -1$ paired with a right angle, with either ordering. Choose fixed overlapping groups

$$B = [-b_* - w, -b_* + w], \quad L = [-\pi, -b_* - \ell_L], \quad R = [-b_* + \ell_R, 0], \qquad 0 < \ell_L, \ell_R < w.$$

There is no equality configuration in a same-group square. For B^2 the angular Schur factor is at most $\sin w/\min(\sin(b_*-w),\sin(b_*+w))<1$ after shrinking w; the other groups have a strict root Schur bound. At the double branch the apparent singularity is removable: the inverse root coordinate is $y_\zeta(\phi)=G_\zeta(\phi^2)$ and

$$P_{ij}=(\phi_i-\phi_j)^2A_\zeta(\phi_i,\phi_j),\qquad A_{\zeta_*}(0,0)=-\frac{1}{4\sin^2 b_*},\tag{6.11}$$

with analytic bounded fixed jets. This follows by factoring both y_i-y_j and $1-e^{-\mathrm{i}(\phi_i+\phi_j)}$; the common $\phi_i+\phi_j$ factor cancels.

On the closed lower support, $1-z_iz_j=0$ forces the already resolved double root $z_i=z_j=1$: the alternative $z_i=z_j=-1$ is excluded by $W_0+1\ge E_*>0$. Away from a smaller double-branch chart, this denominator therefore has a fixed positive minimum. The regular expression (6.11) and the original rational expression agree on their overlaps, and give a finite cover with bounded fixed jets. This also proves continuity at the common endpoints, where the canceled pair is zero. No reciprocal-root pole remains on that cover.

Choose a strict same-group bound $q<q_1<q'<1$ and cross-group slack $\eta>0$ satisfying $\log(1+\eta)<-\log(q')/4$. Project the upper slivers $[0,2\delta]$ and $[\pi-2\delta,\pi]$ to their respective endpoints, assigning them to R,L. If $\bar\theta_i$ denotes this projection (and equals θ_i for lower angles), then uniformly in the entire independent box $0\le\lambda,P/N\le1$,

$$|y_i-e^{\mathrm{i}\bar\theta_i}|+|W_i-W_0(\bar\theta_i)|\le C(\varepsilon+\tau+\delta)+C|\zeta-\zeta_\varepsilon|.$$

Compact-root changes are Lipschitz, and branch changes are one-half Hölder by the square-root factorization and (6.11). The finite cover therefore gives a complete-pair change at most

$$C\sqrt{\varepsilon+\tau+\delta+|\zeta-\zeta_\varepsilon|}.$$

Choose $\delta,\tau,\varepsilon_0$ to make the radial-center change smaller than $\min(q_1-q,\eta/2)$; reserve $q'-q_1$ and $\eta/2$ for the parameter disks. This proves both same-group and cross-group bounds on the slivers, uniformly in the actual coupled occupancy.

The named current index needs only bounded pairs. At its right limiting endpoint the root is $\xi=e^{-\mathrm{i}b_*}$, and for every limiting lower root z, $|1-\xi z|=|e^{\mathrm{i}b_*}-z|\ge\sin b_*>0$. At the left endpoint the root is $z_0<1$ and $|1-z_0z|\ge1-z_0$. These margins persist on the chosen neighborhoods. The order of choices is essential: the branch/upper-sliver limiting moduli are

$$\frac{\sin((b_*+a)/2)}{\sin((b_*-a)/2)}=1+a\cot(b_*/2)+O(a^2),\qquad \frac{\cos((b_*-a)/2)}{\cos((b_*+a)/2)}=1+a\tan(b_*/2)+O(a^2).$$

They can exceed one. Keeping the compact-group margins fixed before shrinking the slivers is what preserves the strict same-group bound; no claim that every sliver pair contracts is used.

For $n=N$ lower variables in K, or $n=N-1$ in a current, the number of same-group pairs is at least $n^2/6-n/2$ and the number of cross pairs at most $n^2/3$. After deleting at most j factors, apply the bounds directly to the remaining product; no division by a possibly zero pair is allowed. The result is

$$\prod_{(i,l)\notin\mathcal{E}}|P_{il}|\le C_j^Ne^{-\kappa N^2},\qquad|\mathcal{E}|\le j,\tag{6.12}$$

for some fixed $\kappa>0$. Named pairs cost only C^N.

The selected contour, including arbitrarily large N. Telescope $1-\psi$ so that a selected index has $p_q = 1$ and hence $P \geq 1$. At $\lambda = 1$ it has a fixed interior root gap, and $\sum h_i \leq -3\tau P/2$ supplies a fixed Y gap. Every true branch has $\Im W \geq cP/N$, so its residue factor is bounded by $C\sqrt{N}$. These facts persist on a common disk $|\zeta - \zeta_\varepsilon| \leq c/N$. Branch roots remain in the lower unit half-disk; compact roots move by $O(1/N)$, and the fixed anchor absorbs their total inflation $e^{CN(c/N)}$ when c is small.

Let M be the number of compact indices. Divide their radial-center roots into a strict interior group and a compact lower-arc group. Same-group Schur factors are at most a fixed number below one. For a branch root w on the actual disk and a compact radial-center root z_0, the denominator $|1 - z_0 w|$ has a fixed positive lower bound, and

$$\frac{z-w}{1-zw} - \frac{z_0 - w}{1 - z_0 w} = \frac{(z - z_0)(1 - w^2)}{(1 - zw)(1 - z_0 w)}.$$

Thus mixed factors are at most $1 + C/N$; only the compact root has been displaced in this comparison. It follows, even with the weaker $1 + C/\sqrt{N}$ budget, that the entire root Pfaffian product is at most $C^N e^{-aM^2}$.

Every y-pair involving a true branch is bounded: its reciprocal upper partner has inward shift -2τ, dominating the outward shift at most $\tau/2$. Compact pairs have the occupancy-independent majorant

$$\left| \frac{y_i - y_j}{1 - y_i y_j} \right| \leq \frac{C}{\varepsilon + \operatorname{dist}(\theta_i + \theta_j, 2\pi\mathbb{Z})}.$$

Indeed the sum of the two shifts is nonpositive at reciprocity and is $O(\operatorname{dist}(\theta_i + \theta_j, 2\pi\mathbb{Z}))$ nearby. A matched two-angle integral therefore costs $O(H + 1)$. Expand the angular Pfaffian into bounded minors and a matching on an even compact subset. Hadamard bounds the minor by $C^N N^{(N+1-2r)/4}$, the subset and matching count by N^{3r}, and the disjoint singular pair integrals by $(C(H+1))^r$. Including the one-body cost $C^N N^{N/2}$ and $N!$, the complete selected integral is bounded, for each assignment, by

$$C^N N^{-N/4+1/4} (CN^3(H+1))^M e^{-aM^2}.$$

The C^N assignments and named-index choices are absorbed in C^N. Completing the square in M, and applying Cauchy only to the complete integral with its fixed angular weights, gives

$$\sum_N |F_N^{(j)}(\zeta_\varepsilon)| \leq C_j \exp(C_j \log^2(H+2)). \tag{6.13}$$

Here $e^{C_j \log^2(N+1)} N^{-N/4+j+1/4}$ is summable. Pointwise assignments need not be holomorphic, because none is separately differentiated. The argument applies to every N, not just a moving finite window, and also with the product poles deleted.

The original and protected current disks. On $|\zeta - \zeta_\varepsilon| \leq c\varepsilon$, the undeformed imaginary margin and its monotone radial improvement keep all roots lower and interior. The two signed product gaps are at least $c'\varepsilon$. Near a true branch set $\mu = \lambda P/N$ and $u = \theta + b_*$. For $y = e^{v + \mathrm{i}\theta}$, $v = -c_0\varepsilon + \tau\mu/2$, the exact formulas

$$\Re W = \Re E - \cosh v \cos\theta, \qquad \Im W = \Im E - \sinh v \sin\theta$$

give an imaginary margin $c(\varepsilon + \mu)$ and real branch-center displacement $O(\varepsilon^2 + \mu^2 + |\zeta - \zeta_\varepsilon|)$. Splitting at $|u| = K(\varepsilon + \mu)$, first using the imaginary margin and then the linear real term, proves

the occupancy-independent majorant

$$|R(y;\zeta)| \le C(|u|+\varepsilon)^{-1/2}. \tag{6.14}$$

Its integral is bounded. Compact factors and Jacobians cost C^N. Branch-coordinate motion on the disk is $O(\sqrt{\varepsilon})$; choosing ε_0 small retains the reserved pair slack, so (6.12) holds throughout the disk. Cauchy therefore gives

$$|K_N^{(j)}| + |J_{N,j}(\lambda)| \le C_j^N \varepsilon^{-j-2} e^{-\kappa N^2}, \qquad 0 \le \lambda \le 1, \tag{6.15}$$

where $J_{N,j}$ is the already differentiated, angularly integrated current. This bound will be used only for large N.

For a named current and $\lambda \ge \lambda_*$, $P \ge 1$ protects a larger disk $|\zeta - \zeta_\varepsilon| \le c\lambda_*/N^2$. Its branch margin is at least $c(\varepsilon + \lambda P/N)$. Choose its disk constant after τ so that the parameter displacement uses at most one quarter of this margin. Compact roots move by $O(\lambda_*/N^2)$; the named upper root has modulus at most $e^{-c\lambda}$ and absorbs the total compact inflation. Both product gaps are therefore at least $c'\lambda$. The same split proving (6.14), now including the disk displacement, gives the same integrable majorant. Pair-coordinate motion uses the remaining fixed slack in (6.12). Cauchy and then integration in λ yield

$$\sum_N \int_{\lambda_*}^1 |J_{N,j}(\lambda)| \, \mathrm{d}\lambda \le C_j \lambda_*^{-j-2} = C_j \varepsilon^{-\alpha_0(j+2)}. \tag{6.16}$$

The convergent sum $\sum_N C^N N^{2j} e^{-\kappa N^2}$ includes all orders. No large-current estimate is used to justify the original disk.

6.4 Real-domain differentiation in the intermediate window

The remaining orders satisfy $N_0 + 2 \le N < D\sqrt{H}$, with fixed D. For the small current $0 \le \lambda \le \lambda_*$, all finite compact jet errors $N^{C_j}(\varepsilon + \lambda_*)$ tend uniformly to zero in this window. We prove the required bound

$$|K_N^{(j)}| + \int_0^{\lambda_*} |J_{N,j}(\lambda)| \, \mathrm{d}\lambda \le e^{C_j N \log(N+1)} (H+N)^{C_j}. \tag{6.17}$$

Lemma 6.7 (Differentiation with simple kernels)**.** *Let w be a fixed smooth real angular weight, and let V be a parameter-dependent complex vector field. Put $\mathcal{D} = \partial_\zeta - \mathcal{L}_V$ on top forms. If all puncture fluxes vanish at every preceding stage, then*

$$\partial_\zeta^j \int w\Omega_\zeta = \int \mathcal{D}^j (w\Omega_\zeta).$$

If $(\partial_\zeta - V\cdot\nabla)Y = (\partial_\zeta - V\cdot\nabla)Z = 0$, both signed product kernels remain simple at every order.

Proof. The Lie derivative of a top form is a divergence. Integrate it on the fixed real domain, with punctures removed, and use the vanishing flux. Repeat on the entire preceding top form, including all derivatives of V, its divergence, and the previous coefficients. Complex coefficients are permitted by linearity of real integration by parts; no flow of a smooth cutoff into a complex domain is assumed. □

On an original branch chart set $\phi = \arccos W$ and $\chi_\zeta : u \mapsto \phi$. The branch one-body form is regular:

$$\frac{\mathrm{d}y}{2\pi\mathrm{i}} \frac{z}{\sinh\gamma} = -\frac{z}{\pi(1-y^{-2})} \, \mathrm{d}\phi.$$

Together with (6.11), the regular numerator is $\Delta(\phi)^2 A_\zeta(\phi)$ with analytic bounded fixed jets. Put $a_i = E'/W_{u_i}$, $b_i = \partial_{u_i}\phi_i$, $A = \sum_i a_i$. For a selected pair p, q use

$$V_i = a_i \ (i \neq p, q), \qquad V_p = a_p + \frac{Ab_q}{b_p - b_q}, \qquad V_q = a_q - \frac{Ab_p}{b_p - b_q}.$$

Then $\sum V_i = 0$ and $\sum b_i V_i = \partial_\zeta \sum \phi_i$. The residual regular-coordinate field $B = \partial_\zeta \chi_\zeta - (\chi_\zeta)_* V$ has only

$$B_p = -\frac{Ab_p b_q}{b_p - b_q}, \qquad B_q = -B_p.$$

Because $b(\phi) = B_\zeta(\phi^2)/\phi$ with B_ζ analytic and $B_\zeta(0) \neq 0$, one has the exact analytic factorization

$$\phi_q B_\zeta(\phi_p^2) - \phi_p B_\zeta(\phi_q^2) = (\phi_q - \phi_p)\{B_\zeta(0) + O(|\phi_p \phi_q|)\}.$$

The expression in braces is bounded away from zero on a smaller fixed chart. Thus $b_p b_q/(b_p - b_q)$ is an analytic bounded coefficient divided by $\phi_q - \phi_p$, with bounded required jets. All fixed coefficient jets have polynomial N-cost apart from this explicit selected difference pole. For smooth real w the exact pullback identity is

$$(\partial_\zeta - \mathcal{L}_V)(w\chi_\zeta^* \Omega) = w\chi_\zeta^*(\partial_\zeta + \mathcal{L}_B)\Omega - (V \cdot \nabla_u w)\chi_\zeta^* \Omega. \tag{6.18}$$

It follows by differentiating the Jacobian determinant and subtracting the two divergence formulas. The real cutoff is never holomorphically extended.

After j operations, each term has a derivative $\partial_u^\mu w$, a numerator derivative $\partial_\zeta^\ell \partial_\phi^\nu(\Delta^2 A)$, the unchanged simple kernels, and a coefficient of selected pole order m satisfying

$$m + |\mu| + |\nu| \leq 2j, \qquad \ell + |\mu| + |\nu| \leq j. \tag{6.19}$$

Here is the full induction: a fixed-ϕ parameter derivative of a coefficient adds no difference pole; $B \cdot \nabla$ on a coefficient, or multiplication by $\operatorname{div} B$, adds at most two; a numerator-coordinate derivative adds one derivative and at most one pole; a numerator-parameter derivative adds only one derivative; and a cutoff derivative adds one cutoff derivative and at most one pole. These are all product-rule terms in (6.18). There are $N^{O(j)}$ descendants. At most j complete pair factors are changed in each descendant, so (6.12) survives on the others. One must not divide by the original Vandermonde, which would create poles at unselected collisions.

To select pairs smoothly, fix an integer $M > 2k + 2$ and use

$$w_{pq} = \frac{(u_p - u_q)^{2M}}{\sum_{i<l}(u_i - u_l)^{2M}}, \quad d_u = \max_{i<l} |u_i - u_l|.$$

Off full equality these sum to one and obey

$$|\partial_u^\mu w_{pq}| \leq C_k N^{C_k} \frac{|u_p - u_q|^{2M - |\mu|}}{d_u^{2M}}.$$

The analytic inverse branch is $u = U_\zeta(\phi^2)$, with its fixed contour constants included. Its bounded derivative gives

$$|u_p - u_q| \leq C|\phi_p^2 - \phi_q^2| \leq C'|\phi_p - \phi_q|.$$

The rational weights therefore absorb the selected poles, losing at most $2j$ powers of shape radius. A further power is reserved for a transition or flux estimate. At fixed ε, N, a selected-diagonal

tube of radius h has flux $O(h^{2M+1-2j})$ on a separated support. At full equality, the shape surface measure gives flux exponent

$$N(N-1)-2j-1+N-2=N^2-2j-3>0 \quad (N\ge N_0+2,\ j\le k).$$

Thus all puncture fluxes vanish, in this order, at every preceding Lie stage. These checks justify the identities before taking boundary-uniform bounds.

Branch geometry used in coarea. For the original contour, near $u=0$,

$$1-W=a_*u-\mathrm{i}b'_*\varepsilon+O(u^2+\varepsilon|u|+\varepsilon^2),\quad a_*=\sin b_*>0,\quad b'_*=2\sin\theta-c_0a_*>0.$$

The exact factorization $\phi=\sqrt{2(1-W)}\,h(1-W)$, $h(0)=1$, gives

$$|\phi'|\asymp(|u|+\varepsilon)^{-1/2},\quad \int|\phi'|\,\mathrm{d}u\le C,\quad -\Im\phi\ge c\sqrt{|u|+\varepsilon}\quad (u\le 0).$$

For $u\ge 0$, $x'=\partial_u\Re\phi\asymp(u+\varepsilon)^{-1/2}$, and $-x''\ge cu^{-3/2}$ for $u\ge C_0\varepsilon$. If $M_1/\varepsilon\to\infty$ and $0\le m_1\le M_1/2$, then

$$x'(m_1)-x'(M_1)\ge c(m_1+\varepsilon)^{-1/2},\quad \frac{|\phi'(m_1)\phi'(M_1)|}{x'(m_1)-x'(M_1)}\le CM_1^{-1/2}.$$

For $M_1/2<m_1<M_1$, with $m_1\gg\varepsilon$, the last ratio is at most $C\sqrt{M_1}/(M_1-m_1)$. These follow from the leading square-root derivative, using its $u^{-3/2}$ curvature in the comparable case. On a true branch plateau of the small current the same factorization holds with scale $\varepsilon+\lambda P/N$. In particular $\Re\phi_u\ge(2/3)|\phi_u|$ for $u\ge 0$, and

$$\frac{|\phi'_u|}{|1-\sigma Z|}\le\frac{C}{|u|+\varepsilon+\lambda P/N}\quad (u<0). \tag{6.20}$$

The possible real branch-center displacement is quadratic in this scale; it is retained in deriving these bounds. A plateau variation fixes P.

All-branch microcore. Choose $b_N=c_*e^{-(A+4)N}/N$ and localize by $\prod_i\chi(u_i/b_N)$, with $\chi=1$ on $[-1/2,1/2]$ and supported in $[-1,1]$. Uniformly for $N<D\sqrt{H}$, $N\varepsilon e^{AN}\to 0$ and $\varepsilon/b_N^m\to 0$ for each fixed m. Therefore $|1-\sigma Y|\ge\frac12 e^{-AN}$ on the core and all cutoff jets. The one-phase field $V_i=a_i$ freezes every regular ϕ_i, hence Z, and preserves the Vandermonde polynomial. Its jth derivative has only $e^{O_j(N)}$ cutoff and Y-kernel losses and $e^{O_j(N^2)}$ analytic product cost. The branch image has radius $C\sqrt{b_N}$ in a fixed cone; choosing c_* small makes $N\sqrt{b_N}$ small and $|1-\sigma Z|\ge c\sum_i|\phi_i|$ (a weaker bound also valid for $\sigma=-1$). The Laplace identity for the reciprocal of this sum and

$$\int_{\mathrm{core}}e^{-t|\phi|}|\,\mathrm{d}\phi|\le C\min(\sqrt{b_N},t^{-1})$$

bound the core derivative by

$$e^{C_jN^2}b_N^{(N^2-1)/2}. \tag{6.21}$$

Its logarithm is $-(A+4)N^3/2+O_j(N^2\log(N+1))$, and the series is bounded independently of ε.

All-branch complement. Telescope the complement of the microcore. A named anchor satisfies $|u_a| \ge b_N/2$ on every cutoff jet. Put $\bar{u} = N^{-1}\sum u_i$, $\rho^2 = \sum(u_i - \bar{u})^2$ and split smoothly at $\rho_N = e^{-BN}$, where B is fixed large enough that $2\rho_N \le b_N/8$ for every $N \ge N_0 + 2$.

On $\rho \ge \rho_N$, maximal separation is at least $\rho_N/\sqrt{N}$. The Lie losses and cutoff jets cost $e^{O_j(N\log(N+1))}$ and unchanged pairs retain quadratic suppression. If all coordinates are nonnegative, select the actual minimum m_1 and maximum M_1 only in this final absolute integral. For fixed other variables the phase map

$$(m_1, M_1) \longmapsto (m_1 + M_1 + \text{constant},\ x(m_1) + x(M_1) + \text{constant})$$

has positive Jacobian $x'(m_1) - x'(M_1)$ and multiplicity one on each extreme-index cell. The preceding branch estimates absorb the two branch measures at cost $e^{O(N)}N^{O(1)}$; here $M_1 \ge b_N/2$, and in the comparable case $M_1 - m_1 \ge \rho_N/\sqrt{N}$. The two unwrapped phase intervals have bounded length for fixed other variables.

If some negative coordinate has magnitude at least $\delta_N = e^{-C_1 N} \le b_N/16$, then Z has gap $e^{-O(N)}$; integrate the remaining Y kernel in the named anchor, whose measure costs at most $e^{O(N)}$. If every negative coordinate has magnitude below δ_N, the anchor is positive; use one negative coordinate with (6.20) and use the anchor for the Y kernel. Both integrations cost at most $e^{O(N)}(H + N)$.

On $\rho \le 2\rho_N$, the anchor gives $|\bar{u}| \ge 3b_N/8$, so all coordinates have the same sign and $|u_i| \ge b_N/4$. The branch measures and the conversion of the Vandermonde to the angular shape radius cost at most

$$C^{N^2} b_N^{-N^2/2} = \exp\left(\tfrac{A+4}{2}N^3 + \tfrac{1}{2}N^2\log N + O(N^2)\right).$$

The remaining shape power after Lie losses and coarea is

$$L_N = N^2 - 2j - 4 \ge \begin{cases} 4N_0 + 2, & N_0 \text{ even}, \\ 4N_0 + 1, & N_0 \text{ odd}. \end{cases}$$

For negative mean the Z gap is $e^{-O(N)}$ and a single mean phase suffices. For positive mean the phase map $(\bar{u}, \rho) \mapsto (N\bar{u} + \text{constant}, \sum_i x(u_i))$ has Jacobian at least $cN\rho$, by strict concavity and the zero first shape derivative. It has multiplicity one on each shape ray. Bound ρ^{L_N} by $(2\rho_N)^{L_N}$ before integrating the phase kernels. Both unwrapped phase lengths are $O(N)$, giving an explicit factor $CN^2(H + N)^2$. The logarithm of the total bound is

$$\left(\tfrac{A+4}{2} - B\right)N^3 + \tfrac{1}{2}N^2\log N + O_j(N^2) + B(2j+4)N,$$

apart from that phase factor. Choose $B > A + 4$ after fixing the anchor-neighborhood constants. This is summable; finitely many small orders have fixed, possibly large, constants. The same positive power already proved the required full-equality puncture fluxes vanish.

For the simple-kernel integrations just used, the elementary inequality

$$|1 - e^{-a+\mathrm{i}t}|^{-1} \le C\big(\min(1, a) + \operatorname{dist}(t, 2\pi\mathbb{Z})\big)^{-1}$$

gives $C(1 + \ell)(1 + |\log a|)$ on an interval of length ℓ. For two phases, divide out the specified real Jacobian, enlarge to the containing rectangle, and apply this estimate twice. Periodic wrapping is accounted for by interval lengths, separately from the stated unwrapped multiplicity. A coarea radius is never integrated a second time after being exchanged for a phase.

Mixed and compact sectors. Use regular ϕ_i on true branch indices J, and original angles on compact indices. Let q be the named compact anchor, choose $j_0 \in J$, and put

$$A = \sum_{i \in J} a_i, \quad D_0 = \sum_{i \notin J} \partial_\zeta \phi_i, \quad X_0 = \frac{D_0 + b_q A}{b_{j_0} - b_q}.$$

Set $V_i = a_i$ for $i \in J \setminus \{j_0\}$, $V_{j_0} = a_{j_0} + X_0$, $V_q = -A - X_0$, and all others zero. This freezes Y, Z and every unselected branch phase. The sole residual branch velocity is $-(D_0 + b_q A)/(1 - b_q/b_{j_0})$. The nested support has $|b_q/b_{j_0}| \le 1/4$, so this coefficient and all fixed jets have polynomial N-cost in regular coordinates; there is no selected-difference pole. On current supports the chosen indices are on true plateaus, giving $V \cdot \nabla P = V \cdot \nabla \sum m_i = 0$ identically, including parameter derivatives. The hybrid Jacobian is triangular; its branch measure is precisely the regular one displayed above, and its compact factor retains the full $\prod_i d_i$ determinant.

For positive branch coordinate, the phase Jacobian in the compact anchor and selected branch is $\Re(b_{j_0} - b_q) \ge (2/3 - 1/4)|b_{j_0}|$; its fixed-first-phase map is injective and absorbs the selected branch measure. If a branch coordinate is negative, use (6.20) instead. With a compact left root, Z has a fixed gap: take $V_i = a_i$ on branches and $V_q = -A$, freezing Y and each branch root, and differentiate the separated Z kernel at polynomial cost. These alternatives prove (6.17) for the mixed and left sectors using (6.12) and the simple phase estimates.

On an all compact right interval put $g(\theta) = \arccos(E_* - \cos\theta)$. Direct differentiation gives

$$g''(\theta) = -\frac{E_*(1 - \cos\theta \cos g)}{\sin^3 g} \le -c_R < 0.$$

The coupled-contour finite jets differ from their limiting values by $N^{O(j)}(\varepsilon + \lambda)$, uniformly small in this window. Thus $G = \Re \sum_i \phi_i$ retains a strictly negative Hessian. Let $f_i = \partial_{\theta_i} \log Y$, $g_i = \partial_{\theta_i} \sum \phi_i$, and $Q_0 = \partial_\zeta \sum \phi_i$. By permutation symmetry and smooth division,

$$f_p g_q - f_q g_p = (\theta_p - \theta_q) c_{pq}, \qquad |c_{pq}| \ge c_R/2.$$

The field $V_p = -Q_0 f_q/(f_p g_q - f_q g_p)$, $V_q = Q_0 f_p/(f_p g_q - f_q g_p)$ freezes both kernels. The same five-case induction and rational weights lose at most two shape powers per Lie order. Coupled occupancy is included: keep each complete pair as a scalar function of all angles. A derivation changes at most one new pair per product-rule step, even when its occupancy derivative moves many roots. Thus each descendant changes at most j pairs, has $N^{O(j)}$ terms, and retains (6.12) on the others.

For maximal separation at least N^{-K_0} all selected losses are polynomial. The corresponding real coarea assertion also holds for the genuinely coupled contour. Fix the other angles and the mean of an ordered extreme pair. By permutation symmetry $G(\dots, m + d, \dots, m - d, \dots)$ is even in d, and its second derivative is at most $-2c'_R$ for a fixed $c'_R > 0$. Integrating from $d = 0$ gives

$$|\partial_p G - \partial_q G| \ge c'_R |\theta_p - \theta_q|.$$

Consequently the phase map $(\theta_p, \theta_q) \mapsto (\theta_p + \theta_q, G)$ has multiplicity one on each ordered extreme-index cell and Jacobian at least c'_R times the maximal separation. Its coarea cost is polynomial in N; unwrapped phase lengths $O(N)$ give a further $N^{O(1)}(H + \log N)^2$ factor by the simple-kernel estimates. Below that scale use $\theta_i = \mu + \rho\omega_i$, $\sum \omega_i = 0$, $\sum \omega_i^2 = 1$. Then $\rho \le \sqrt{N}\, N^{-K_0}$, not merely N^{-K_0}. The phase map $(\mu, \rho) \mapsto (N\mu, G)$ has Jacobian at least $c_R N \rho/2$ and multiplicity one on each ray. The numerator costs $C^{N^2} \rho^{N(N-1)}$. After the same conservative losses its remaining

coarea power is $L_N > 0$ above. A fixed sufficiently large $K_0 > 1$ absorbs C^{N^2} and the polynomial field costs. The phase rectangle has side lengths $O(N)$ and attenuation floors $cN\varepsilon$, so its integral costs $CN^2(H + \log N)^2$. Rational-weight and full-equality fluxes vanish as already checked. This proves (6.17) in the last support type.

Onsite terms and completion of the tail. For f_N repeat the exact weighted deformation with both product poles deleted. The selected estimate (6.13) remains valid. For the other pieces use only the one-phase branch field $V_i = (\partial_\zeta \phi_i)/(\partial_{\theta_i} \phi_i)$ and zero velocity on compact indices. It freezes each branch phase, has bounded analytic coefficients, and preserves occupancy on the branch plateaus. There are no selected poles or punctures. Every fixed derivative changes at most j pairs; regular branch lengths are uniformly integrable by (6.14). Thus the remaining pieces are bounded by $C_j^N N^{C_j} e^{-\kappa N^2}$ for every N and $0 \le \lambda \le 1$. Consequently

$$\sum_N |f_N^{(j)}(\zeta_\varepsilon)| \le C_j \exp(C_j \log^2(H+2)) = o(\varepsilon^{-\delta}) \quad \text{for every } \delta > 0. \tag{6.22}$$

Differentiate the locally normally convergent series first. Only afterwards split its already differentiated terms and current at $N = D\sqrt{H}$ and λ_*. Choose a common κ in the pair estimates and then D with $\kappa D^2 > k + 2$. The selected sum is $e^{O_j(\log^2(H+2))}$. The large current is $O_j(e^{\alpha_0(j+2)H}) = o(e^{H/2})$. The entire remaining high-order tail from (6.15) is at most

$$\exp((j + 2 - \kappa D^2)H + O_j(\sqrt{H})) = o(e^{H/2}).$$

Summing (6.17) over all intermediate orders costs $\exp(O_j(\sqrt{H}\log(H+2))) = o(e^{H/2})$. This proves the assertion for signed T_N^σ at positive E_*; (6.9) and conjugation cover the other odd family. Equation (6.22) and $C_N = f_N + 2T_N$ prove the full-site assertion of Theorem 6.5.

Order of constants and quantifiers. The point, N_0, k, radial margin and all required derivative orders are fixed first. Next fix branch and compact margins and the nested slope separation; then the same-group pair bound, its cross slack, and only then the endpoint slivers and τ. Choose disk constants after these data, preserving the branch and named-anchor gaps. Next fix α_0, D, M, K_0 and the algebraic separation constant A. Choose the microcore prefactor and determine its anchor-neighborhood costs before choosing B, C_1 large enough for the inclusions at every $N \ge N_0 + 2$ and the negative cubic coefficient. Finally choose one ε_0 small enough for all the preceding estimates. The elementary limits $\varepsilon^r e^{CN} N^m \to 0$ for $N \le D\sqrt{H}$ and fixed $r > 0, C, m$ make this single choice uniform in that window. Finitely many small-order constants may be large but remain independent of ε. No particle or current window endpoint is differentiated.

6.5 Dense local obstructions and the exact physical identities

The full-site form-factor identification gives, on the respective real physical intervals,

$$\mathcal{X}_L(s) = (1 - s^{-4})^{1/4} \sum_{N \ge 2,\ N \text{ even}} C_N(s), \qquad s > 1, \tag{6.23}$$

$$\mathcal{X}_H(s) = \frac{(1 - s^4)^{1/4}}{s} \sum_{N \ge 1,\ N \text{ odd}} C_N(1/s), \qquad 0 < s < 1. \tag{6.24}$$

The low-temperature formula has the plus-phase subtraction already performed: its omitted zero-particle term is M_+^2, and its onsite contribution is $1 - M_+^2$. Equation (6.4) therefore also gives the

exact alternative $\mathcal{X}_L = 1 - M_+^2 + 2M_+^2 \sum_{N\geq 2,\ N \text{ even}} T_N$. The high-temperature onsite sum is one, and $C_1 = 1/(E-2)$ checks the normalization. The branches in (6.23)–(6.24) are positive on their stated real intervals.

Proposition 6.8 (Dense obstructions for the physical isotropic sums)**.** *The full normalized even exterior sum in* (6.23) *and the full normalized odd interior sum in* (6.24) *have no local holomorphic extension at a dense set of points of* $|s| = 1$*. At their selected points the nonzero leading derivatives are those of Proposition 6.4, multiplied by the physical nonvanishing factors. These assertions use the newly proved full-tail bounds of Theorem 6.5.*

Proof. At every selected point, Proposition 6.4, the bounded lower orders, and Theorem 6.5 give a nonzero leading derivative of the complete physical sum. All lower derivatives are smaller than that leading scale, so multiplication by its local analytic nonzero phase prefactor preserves it by the product rule. A holomorphic extension through such a point would bound that fixed finite derivative on the short radial segment, a contradiction. The selected points are dense on the required arcs. For the even sum, the contour transformation $(s, x, y) \mapsto (-s, -x, -y)$ gives evenness, and conjugation completes all four quadrants. For the odd sum both signs were treated explicitly; conjugation completes a dense subset of the circle. The inversion $s = 1/\zeta$ is locally biholomorphic at each unit-circle point, so the odd conclusion concerns the physical interior germ. This proves the stated local obstructions. Their passage to all boundary points and their role in the exact slice and isotropic corollaries are established in Section 7. □

The high-temperature value at $s = 0$ is the removable value one. This can also be checked without the joint representation: uniformly in the real angular variables, $z = s/2 + O(s^2)$, each one-body factor is $O(s)$ and each squared pair is $O(s^2)$, giving $C_N(1/s) = O(s^{N^2})$ with a bound $C^{N^2}|s|^{N^2}/N!$ for sufficiently small $|s|$. The $N = 1$ coefficient is $s/(1-s)^2$, and every higher odd order vanishes after division by s. The exterior normalized low-temperature germ is single valued; the energy-field susceptibility is interpreted on the physical continuation of $\beta_{\text{phys}} = \operatorname{arsinh}(s)/(2J)$. At every selected nonaxis point this factor is analytic and nonzero, so the same dense obstructions hold for $\chi = \beta_{\text{phys}}\mathcal{X}$. No global single-valued inverse temperature on the exterior is asserted.

Finally put $s = \sinh z$ on the equal-coupling restriction. The central cell

$$|\Im z| < \pi/2, \qquad |\Re z| < \operatorname{arsinh}(\cos \Im z)$$

maps biholomorphically onto $|s| < 1$. The right cell

$$\Re z > \operatorname{arsinh}(|\cos \Im z|)$$

is the universal cover of $|s| > 1$ under the same map, with deck translations $z \mapsto z + 2\pi\mathrm{i}$. One direct verification writes $w = e^z$ and solves $w - w^{-1} = 2s$: for $|s| > 1$ exactly one root has $|w| > 1$, since a unit root would give $s \in \mathrm{i}[-1, 1]$. Exponentiation supplies the stated covering. At the selected nonramified boundary points the local isotropic obstructions pull back directly. These identities fix the entire even/odd sums and the physical factor $\beta = z/(2J)$ used in the proof of the slice theorem and its isotropic descendants below.

7 Closure of the joint theorem and its exact descendants

7.1 The complete joint obstruction

Proof of Theorem 1.1. Theorem 2.1 determines the spectral set at every coupling, including collinear and zero-real-part degeneracies. Theorem 2.2 identifies the physical components and their con-

tractibility and p-bounds. Proposition 2.18 gives an explicit plurisubharmonic exhaustion, independent of the susceptibility. Proposition 3.1, with the physical interface and its coefficientwise angular identification, proves joint holomorphy and normal convergence of the actual physical sums.

Take a regular face point whose free coefficient is $b = \cos(2\pi a/\ell)$, with ℓ an odd prime and $0 < a < \ell/2$. By Proposition 2.10 the variables (C, B), after exchanging transfer axes if needed, are independent holomorphic local coordinates, $\Im B \neq 0$, and the physical domain is the full complementary side of that edge. At such a point $S, T \neq 0$, and the edge identity (2.18) implies $p^2 \neq 1$. Thus the locally chosen physical prefactor is analytic and nonzero.

At a positive endpoint, restrict to any holomorphic transversal with nonramified edge coordinate $u = C - B - b$. Its actual anisotropy is a holomorphic function $B(u)$. Conjugating the data when needed gives $\Im B(0) > 0$, and the physical side contains $u = \mathrm{i}\varepsilon$. Proposition 4.3 proves a nonzero singular derivative of the entire odd or even sum, including the complete differentiated higher tail and physical prefactor. The same proposition handles a negative endpoint for the even sum by the exact angular shift and, if necessary, conjugation.

For a negative endpoint of the odd sum use

$$t = \frac{C + B - b}{B}.$$

This is also nonramified on the stipulated transversal, because $B(0) \neq 0$. In its lower-half-root orientation the actual $B(t)$ has $\Im B(0) < 0$, and the local physical side is $\Im(B(t)t) > 0$. Theorem 5.1 gives all finite boundary jets of the complete odd sum and a selected sequence whose coefficient roots diverge. Its exact full-sum asymptotic (5.84) excludes cancellation by every other particle index and winding. Division by the nonvanishing physical prefactor proves the same nonextension for the physical response. The opposite orientation is its complex conjugate. The local uniform complex-parameter hypotheses and constants are stated and proved in Sections 4 and 5; no estimate for real fixed ratios is promoted to a joint estimate without proof.

In particular choose a constant-B transversal. If the joint physical function extended to a neighborhood of the selected point, its restriction to that disk would give the forbidden one-variable extension. This is a valid implication from a proved transversal obstruction to ambient nonextension. The selected points are dense on regular faces by Proposition 2.10, and regular faces are dense in the entire physical boundaries by Proposition 2.9. Corollary 2.16 then excludes componentwise extension at every boundary point, including corners, collinear points and vanishing transfer-coordinate charts. This proves the joint assertion and its transverse-curve strengthening. Proposition 6.8, proved from the full sums in Section 6, supplies the additional local assertion on the collinear equal-coupling restriction.

Finally consider analytic continuation in an unramified Riemann domain over $\mathbb{C}^2$ from one of these physical germs. As long as the projected continuation path remains in its domain, uniqueness of analytic continuation makes every continued germ the germ of the single-valued function already constructed there. If a path first leaves that domain, its first exit is a boundary point. A local biholomorphic chart of the Riemann domain at the continued point would yield a holomorphic function on a base neighborhood agreeing with the physical function on an accumulating domain-side component. The proved componentwise obstruction forbids this. Compactness of the finite path interval guarantees a finite first exit whenever the path reaches outside. Thus neither a monodromy construction nor a different unramified physical sheet bypasses the boundary. This uses the standard notion of holomorphic continuation, not a definition tied to the spectral set. □

7.2 All physical ratios, including the collinear one

Lemma 7.1 (Actual complex-ratio neighborhoods)**.** *At any regular selected point of a positive-ratio slice where its edge coordinate has nonzero derivative, the transverse representation and the boundary estimates hold in a neighborhood with λ complex. The uniformity is on common compact parameter polydisks, not merely pointwise on the real-ratio family.*

Proof. The boundary temperature coordinate z_0 is nonzero. Thus $(z, \lambda) \mapsto (z, w = \lambda z)$ has determinant z_0 and is a local biholomorphism. Its composition with the regular (C, B) coordinate map is holomorphic. If the positive-edge parameter is $u = C - B - b$, its derivative in z is nonzero by hypothesis; the holomorphic implicit-function theorem gives $z = z(u, \lambda)$ and hence $B = B(u, \lambda)$ on a product polydisk. At a negative edge use $t = (C + B - b)/B$ in the same way. Shrinking the polydisk preserves the strict sign of $\Im B$, the nonvanishing physical prefactor and the other strict face conditions. Cauchy estimates on a slightly larger polydisk give the common holomorphic bounds required in Proposition 4.3 and Theorem 5.1. Their proofs therefore apply to the actual complex-ratio family. In the negative case the selected integers may depend on the family parameter, as explicitly stated there. No boundary estimate is claimed uniform through the collinear degeneration at $\lambda = 1$; that restriction is treated by its own proved full-sum estimates. Joint holomorphy on interior compact sets already includes complex-ratio neighborhoods at $\lambda = 1$. □

Proof of Theorem 1.2. The exact open-set equalities are Theorem 2.6. The restrictions are jointly constructed physical functions by Section 3.1, and are holomorphic on precisely the stated pullbacks. A boundary point of either pullback maps to the corresponding joint boundary: it is a limit of domain points and cannot map to the joint interior by the same open-set equality.

First fix $\lambda \neq 1$. Proposition 2.12 proves, separately on this actual complex line, density of regular boundary points at which the local edge map has nonzero derivative and its free phase is one of the chosen prime-grid values. This proof includes every finite positive ratio, whether rational or irrational. In particular it does not rely on intersecting an arbitrary ambient dense set with the line. At each such point the temperature line is a holomorphic curve $B = B(u)$ or $B = B(t)$ of exactly the kind treated in the proof of Theorem 1.1. Apply Proposition 4.3 or Theorem 5.1 to this actual curve, retaining its varying anisotropy. These prove nonextension of the restricted complete physical sum. The one-variable collar in Lemma 2.14 and Corollary 2.16 give every boundary point.

Now let $\lambda = 1$. The exact coefficient and normalization identities in (6.23)–(6.24) identify the restriction with the full even and odd integrals of Section 6; its local obstructions were proved in Proposition 6.8, not assumed from either source manuscript. Lemma 2.7 maps the two slice domains to the exterior and disk, respectively. At every boundary point with $\cosh \zeta \neq 0$ this is a local biholomorphism. The selected obstructing points pull back densely on the nonramified boundary arcs. The remaining points with $\cosh \zeta = 0$ are limits of those arcs. The same local collar therefore proves nonextension everywhere on both slice boundaries. This argument uses no limit as $\lambda \to 1$.

All nonzero even orders were retained after subtraction of the zero-particle M^2, and the exact full-site multiplier retains the origin and both axes. Section 3 proves the physical energy-field equality $\chi = \beta \mathcal{X}$ with the plus phase fixed in LT. Here $\beta = \zeta/(2J_x)$ is an entire function of the slice coordinate and is nonzero on both boundaries: zero lies in the HT interior and is strictly outside the LT closure. Multiplication or division by it cannot change local boundary extendability. The result consequently recovers both response conventions and every clause of the supplied anisotropic Theorem 1.1. □

7.3 Derivation of the complete isotropic theorems from the slice theorem

Proof of Corollary 1.3. Apply the now proved Theorem 1.2 with $\lambda = 1$. The identities $S = T = s$, $p = s^2$, $B = 1$, and

$$\mathcal{D}(\zeta_1, \eta_1; z, z) = s\left[s + s^{-1} - \tfrac{1}{2}(\zeta_1 + \zeta_1^{-1} + \eta_1 + \eta_1^{-1})\right]$$

identify the parent coefficients term by term with the full isotropic coefficients. On the real LT interval $\sqrt{p} = s$, so each factor $\sqrt{p}/\mathcal{D}$ is exactly its isotropic counterpart. The even prefactor becomes $(1 - s^{-4})^{1/4}$. On the real HT interval the dispersion is $s + s^{-1} - \cos q$ and the odd prefactor is $(1 - s^4)^{1/4}/s$, giving the exterior auxiliary variable $1/s$. Normal convergence and the identity theorem extend these identities to their physical complex domains.

For HT, the biholomorphism in Lemma 2.7 transports the restricted holomorphic function to the disk. The removable value at zero is one, independently checked at the end of Section 6. For LT, the normalized termwise identities show invariance under every deck translation $z \mapsto z + 2\pi i k$; the full normalized function therefore descends from $\Omega_{L,1}$ to a single-valued holomorphic function on $|s| > 1$. It is the positive real LT germ and has the stated connected onsite contribution.

Suppose one of these descended functions extended through a point s_0 of the unit circle. Choose a boundary preimage z_0 on its physical slice. Composing the proposed extension with the entire map sinh would extend the slice function near z_0, in agreement on the physical side. This contradicts Theorem 1.2. This composition argument also works at the ramified boundary values $s_0 = \pm i$; no local inverse at those values is required. Hence the entire unit circle is a natural boundary for both normalized germs.

Finally $\beta = z/(2J)$ on the prescribed physical temperature continuation yields the energy-field claims. In the disk it is the branch $\operatorname{arsinh}(s)/(2J)$. In the exterior it is interpreted on the proved covering. At all selected nonaxis points it is analytic and nonzero, so the local singular derivative remains nonzero after the product rule; density gives all remaining boundary points. This proves exactly the mathematical HT and connected LT descendants, with distinct germs and the full odd/even sums. No old natural-boundary theorem was appended as an assumption. □

8 Dependency map and verification scope

ID	Obligation	Load-bearing derivation
P0	Physical full observable and phase	Equations (3.1)–(3.5); uniform cylinder limit; Gibbs-state identification; connected subtraction and GHS argument.
G1	Independent joint domains	Theorems 2.1, 2.2; complete exceptional-case algebra; Proposition 2.18.
A1	Joint holomorphy and branches	Double contours, Proposition 3.1, integer powers at the HT axes, coefficientwise angular gluing.
N1	Positive-edge full-sum obstruction	Lemma 4.2, Proposition 4.3; exact constrained Gaussian amplitudes, fixed-index remainder and infinite differentiated tail.
N2	Negative odd-edge full-sum obstruction	Theorem 5.1; exact pole reduction, dimension-uniform quadratic coordinates, beta error, exact finite differences, all-index equation (5.84).

ID	Obligation	Load-bearing derivation
N3	Every joint boundary point and physical sheet	Regular joint density, nonzero coordinate Jacobian, Lemma 2.14, Corollary 2.16, first-exit continuation argument.
R1	Every fixed positive ratio, exact domains	Theorem 2.6; separate slice-density proof; N1/N2 on the actual analytic curve, including nonconstant B.
I0	Collinear local analytic mechanism	Proposition 6.4, Theorem 6.5, Proposition 6.8; two complete chart phases, all orders and onsite terms.
I1	Exact isotropic descendants	The proved R1 theorem at $\lambda = 1$, exact termwise identities, Lemma 2.7, descent and composition at ramified boundary values.

All estimates involving an infinite number of particles in this paper are analytic bounds with their index ranges specified. The numerical diagnostics accompanying the manuscript do not replace any such bound. The exact checks use standard-library polynomial reduction and rational Gaussian moments: they verify the Cramer factorizations, the regular-face identity, the local coordinate Jacobian, and the constrained Gaussian normalization for dimensions two, three and four. A saved randomized spectral-set diagnostic tests generic and degenerate real-part cases. A separate floating-point diagnostic checks the exact negative-edge finite-difference model for a nonconstant complex $B(t)$. Its finite output is a consistency check on formulas, not an asymptotic certificate.

Two destructive tests are useful to state explicitly. First the decoupled restriction $K_y = 0$ is entire, as proved in Remark 3.2; it refutes an unrestricted principle that ambient nonextension passes to every complex line. The positive-ratio slice theorem instead uses its explicit transverse-curve argument and separate collinear proof. Second, Σ cannot be replaced by $\{|p| = 1\}$: at

$$z = \mathrm{i}\pi/2, \qquad w = \operatorname{arsinh}(1/2)$$

one has $A = 0, S = \mathrm{i}, T = 1/2$, so $a = b = 0$ represents a spectral point, while $|p| = 1/2$. In fact $F_1 = -1/4, F_2 = 1$, so this point lies in the interior of Σ. This test refutes the naive modulus geometry without asserting a physical full-sum singularity at that point.

The mathematical verification recorded with this manuscript consists of the displayed derivations, separate initial investigator records, subsequent cross-reconstruction, and the stated exact or finite diagnostics. It is internal verification in one research campaign, not an external proof certificate. No publication-priority conclusion or S4 significance rating is inferred from it. The physical exact-solution and Gibbs-state imports are the named published results; the supplied manuscripts are retained as source provenance.

References

[1] A. I. Bugrij and O. Lisovyy, *Correlation function of the two-dimensional Ising model on a finite lattice. II*, arXiv:0708.3643 (2007), especially (18), (20), (25), and (35)–(39). arXiv:0708.3643.

[2] N. Iorgov and O. Lisovyy, *Ising correlations and elliptic determinants*, arXiv:1012.2856 (2010), Theorem 6.1 and Remark 6.2. arXiv:1012.2856.

[3] W. P. Orrick, B. G. Nickel, A. J. Guttmann and J. H. H. Perk, *The susceptibility of the square lattice Ising model: New developments*, arXiv:cond-mat/0103074 (2001), especially (1.8)–(1.11) and Section 3.2. arXiv:cond-mat/0103074.

[4] S. Boukraa, S. Hassani and J.-M. Maillard, *Holonomic functions of several complex variables and singularities of anisotropic Ising n-fold integrals*, arXiv:1207.1784v2 (2012), Section 6, especially the unsquared equation (54). arXiv:1207.1784.

[5] A. Raoufi, *Translation-invariant Gibbs states of Ising model: General setting*, arXiv:1710.07608 (2017), Theorem 1 and assumptions C1–C4. arXiv:1710.07608.

[6] F. Camia, C. Garban and C. M. Newman, *The Ising magnetization exponent on* $\mathbb{Z}^2$ *is* $1/15$, arXiv:1205.6612v4 (2013), Theorem 1.5, the GHS inequality for nonnegative interactions and site fields. arXiv:1205.6612v4.

[7] User-supplied complete low-temperature manuscript, *Low-temperature square-lattice Ising bulk susceptibility: natural-boundary argument*, `MANUSCRIPT_LT.pdf` and editable source, in the fixed input package dated 21 October 2026.

[8] User-supplied complete high-temperature manuscript, *High-temperature bulk Ising susceptibility: a natural-boundary proof*, `MANUSCRIPT_HT.pdf` and editable source, in the same fixed package.

[9] User-supplied complete anisotropic manuscript, *Natural boundaries of the full bulk susceptibility for the fixed-ratio anisotropic square-lattice Ising model*, `MANUSCRIPT_ANISO.pdf` and editable source, in the same fixed package.