Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Thu Jul 07 14:12:32 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
50
40
30
20
10
0
0
2
4
6
8
Age (yr)


Age (yr)
0
2
4
6
8
CV of lengths
0
0.04
0.08
Length (cm, beginning of the year)
50
40
30
20
10
0
SD of lengths
0.0
1.0
2.0
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


50
40
30
20
10
0
Length
0
2
4
6
8
Age


50
40
30
20
10
0
Length
0
2
4
6
8
Age


Mean weight (kg) in last year
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
Female weight (kg)


2.0
1.5
1.0
0.5
0.0
Fecundity
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
Female length (cm)


Spawning output
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
Length (cm)


Spawning output
0.00
0.05
0.10
0.15
0.20
0
2
4
6
8
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)
FISHERY
SURVEY2


Selectivity
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
0
2
4
6
8
Age (yr)
1
0.8
0.6
0.4
0.2


Length (cm)
0
10
20
30
40
0
2
4
6
8
Age (yr)
1
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
100
200
300
400
500
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
200
400
600
800
1000
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
2.0
1.0
0.0
−1.0
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
3
2
1
0
−1
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1 - SE(\hat{r}_y)^2/\sigma_R^2$
1.0
0.5
0.0
−0.5
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
100
200
300
400
500
0.0
0.5
1.0
1.5
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
100
200
300
400
500
0.0
0.5
1.0
1.5
Spawning biomass (mt)
1985
1986
1998
1994
1999
2006
2005
2011
1983
2007
1992
1993
1997
1990
2021
2015
2014


Log recruitment deviation
Spawning output (relative to B_0)
2
1
0
−1
−2
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1985
1986
1998
1994
2006
1999
2005
1992
1990
2011
1993
1983
2007
1997
1967
2021
2019
2018
2017
2016
2014
2015


Landings (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
2
4
6
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
Relative spawning output: B/B_MSY
2021
1967


Index
0
10
20
30
40
50
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
Observed index
0
5
10
15
20
25
30
35


Log index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Log observed index


Residual
4
2
0
−2
−4
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
Year


Expected index
0
5
10
15
0
5
10
15
20
25
Observed index


Log index
4
2
0
−2
−4
1990
1995
2000
2005
Year


Log index
4
2
0
−2
−4
1990
1995
2000
2005
Year


Log expected index
0.5
1.0
1.5
2.0
2.5
−1
0
1
2
3
Log observed index


Effective catchability
5.290260
5.290264
5.290268
0.5
1.0
1.5
2.0
2.5
3.0
Vulnerable biomass


Residual
4
2
0
−2
−4
1990
1995
2000
2005
Year


Deviation
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
Year
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020


100
300
500
Age
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


100
300
Age
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


100
300
500
Length
50
40
30
20
10
0
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


50
150
250
Length
0
10
20
30
40
50
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
10
20
30
40
50
60
0
2
4
6
8
Age


FISHERY
Sum of N adj.=9727.2
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
5
10
15
20
25
30
35
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
35
30
25
20
15
10
5
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=247.8
2010
N adj.=153.7
2016
N adj.=352
2005
N adj.=155.4
2011
N adj.=815.6
2017
N adj.=169.7
2006
N adj.=125.2
2012
N adj.=1337.3
2018
N adj.=162.1
2007
N adj.=275.5
2013
N adj.=2574.6
2019
N adj.=220.9
2008
N adj.=297.4
2014
N adj.=1048.3
2020
N adj.=165.5
2009
N adj.=472.1
2015
N adj.=1062.6
2021
N adj.=91.6
Proportion
0.00
0.10
0.20
0.30
0 5 15 25 35
Length (cm)


0.01
0.2
0.4
35
30
25
20
15
10
5
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
23.0
22.5
22.0
21.5
21.0
20.5
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=9727.2
Sum of N eff.=1509.2
Proportion
0.30
0.25
0.20
0.15
0.10
0.05
0.00
0
5
10
15
20
25
30
35
Length (cm)


FISHERY
−1000
−600
200
600
1000
35
30
25
20
15
10
5
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=247.8
N eff.=168.3
2010
N adj.=153.7
N eff.=38.9
2016
N adj.=352
N eff.=87.3
2005
N adj.=155.4
N eff.=71.7
2011
N adj.=815.6
N eff.=80.6
2017
N adj.=169.7
N eff.=39.2
2006
N adj.=125.2
N eff.=71.3
2012
N adj.=1337.3
N eff.=70.9
2018
N adj.=162.1
N eff.=42.8
2007
N adj.=275.5
N eff.=124.7
2013
N adj.=2574.6
N eff.=76.5
2019
N adj.=220.9
N eff.=97
2008
N adj.=297.4
N eff.=62.6
2014
N adj.=1048.3
N eff.=87.9
2020
N adj.=165.5
N eff.=24.3
2009
N adj.=472.1
N eff.=99.2
2015
N adj.=1062.6
N eff.=248.6
2021
N adj.=91.6
N eff.=17.2
Proportion
0.00
0.10
0.20
0.30
0
5
15
25
35
Length (cm)


−1000
−600
200
600
1000
35
30
25
20
15
10
5
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


Effective sample size
0
50
100
200
0
500
1000
1500
2000
2500
Observed sample size
2015
2004
2007
2019
2009
2016
2011
2014
2012
2013
2003
2020
2021
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
23.0
22.5
22.0
21.5
21.0
20.5
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.4
0.8
1.2
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.4
0.8
1.2
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0
2
4
6
0
5
10
15
Total biomass (mt)


Yield per recruit (kg)
0.000
0.005
0.010
0.015
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_inflection_FISHERY(1)
15
20
25
30
35
40
InitF_seas_1_flt_1FISHERY
0.0
0.5
1.0
1.5
2.0
Size_95%width_FISHERY(1)
0
1
2
3
4
5
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Density
Parameter value