Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Thu Jul 28 10:08:48 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
120
100
80
60
40
20
0
0
10
20
30
40
50
Age (yr)


Age (yr)
0
10
20
30
40
50
120
100
80
60
40
20
0
Length (cm, beginning of the year)
CV of lengths
0.04
0.02
0
SD of lengths
4
2
0
Age (yr)
CV of lengths
0
0.02
0.04
SD of lengths
0
1
2
3
4
Length (cm, beginning of the year)


Age (yr)
0
10
20
30
40
50
120
100
80
60
40
20
0
Length (cm, beginning of the year)
Mean weight
0
5
10
15
20
Frac. mature
0.0
0.5
1.0
Age (yr)


Length
120
100
80
60
40
20
0
0
10
20
30
40
50
60
Age


Length
120
100
80
60
40
20
0
0
10
20
30
40
50
60
Age


Mean weight (kg) in last year
25
20
15
10
5
0
0
20
40
60
80
100
120
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
20
40
60
80
100
120
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
20
Female weight (kg)


25
20
15
10
5
0
Fecundity
0
5
10
15
20
Female weight (kg)


Fecundity
0
5
10
15
20
25
0
20
40
60
80
100
120
Female length (cm)


Spawning output
25
20
15
10
5
0
0
20
40
60
80
100
120
Length (cm)


Spawning output
0
2
4
6
8
0
10
20
30
40
50
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
120
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
120
Length (cm)


Length (cm)
0
20
40
60
80
100
0
10
20
30
40
50
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
10
20
30
40
50
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
20
40
60
80
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
20
40
60
80
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.1
0.2
0.3
0.4
0.5
0.6
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
30
35
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.4
0.2
0.0
−0.2
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0
1
2
3
4
0
5
10
15
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0
1
2
3
4
0
5
10
15
Spawning biomass (mt)
2021
1967
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Log recruitment deviation
0.4
0.2
0.0
−0.2
−0.4
2021
1967
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)


Landings (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.00
0.05
0.10
0.15
0.20
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Relative spawning output: B/B_MSY
1980
2010
2020
2000
1990
2021
1967
1970


Index
0
20
40
60
80
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
20
40
60
80
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
Observed index
0
1
2
3
4
0
2
4
6
8
10
12
14


Log index
4
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
4
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
1.1
1.2
1.3
1.4
−2
−1
0
1
2
Log observed index


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
1995
2000
2005
2010
2015
2020
Year


0.1
2
60
50
40
30
20
10
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
60
50
40
30
20
10
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
120
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
120
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
0
10
20
30
40
50
Age


FISHERY
Sum of N adj.=285
Proportion
0.14
0.12
0.10
0.08
0.06
0.04
0.02
0.00
20
40
60
80
100
Length (cm)


FISHERY
0.01
0.2
Length (cm)
110
100
90
80
70
60
50
40
30
20
2008
2011
2013
2015
2017
Year


2007
N adj.=26.6
2014
N adj.=43.3
2008
N adj.=31.2
2015
N adj.=25.9
2009
N adj.=34.9
2016
N adj.=23.2
2011
N adj.=24.5
2017
N adj.=15.1
2012
N adj.=19.2
2018
N adj.=20.8
2013
N adj.=20.2
0.00
0.10
0.20
20
40
60
80
100
Proportion
Length (cm)


0.01
0.2
Length (cm)
110
100
90
80
70
60
50
40
30
20
2008
2011
2013
2015
2017
Year


FISHERY (whole catch)
Mean length
70
65
60
55
50
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=285
Sum of N eff.=646.6
Proportion
0.14
0.12
0.10
0.08
0.06
0.04
0.02
0.00
20
40
60
80
100
Length (cm)


FISHERY
−250
−150
50
150
250
Length (cm)
110
100
90
80
70
60
50
40
30
20
2008
2011
2013
2015
2017
Year


2007
N adj.=26.6
N eff.=46.8
2014
N adj.=43.3
N eff.=98.9
2008
N adj.=31.2
N eff.=135
2015
N adj.=25.9
N eff.=65.3
2009
N adj.=34.9
N eff.=120.4
2016
N adj.=23.2
N eff.=28
2011
N adj.=24.5
N eff.=15.4
2017
N adj.=15.1
N eff.=22.5
2012
N adj.=19.2
N eff.=40.4
2018
N adj.=20.8
N eff.=48.9
2013
N adj.=20.2
N eff.=24.9
Proportion
0.00
0.10
0.20
0.30
20
40
60
80
100
Length (cm)


−250
−150
50
150
250
110
100
90
80
70
60
50
40
30
20
Length (cm)
2008
2011
2013
2015
2017
Year


Effective sample size
0
20
60
100
140
0
10
20
30
40
Observed sample size
harmonic mean
arithmetic mean
2008
2009
2014
2015
2018
2007
2012
2016
2017
2013
2011


FISHERY (whole catch)
Mean length
70
65
60
55
50
2008
2010
2012
2014
2016
2018
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.5
1.0
1.5
2.0
0
20
40
60
80
Total biomass (mt)


Yield per recruit (kg)
0.0
0.4
0.8
1.2
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max_likelihood
2
4
6
8
Size_inflection_FISHERY(1)
20
40
60
80
100
InitF_seas_1_flt_1FISHERY
0.00
0.05
0.10
0.15
0.20
0.25
0.30
Size_95%width_FISHERY(1)
0
10
20
30
40
50
60
70
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value