American Samoa Model Checks

Meg Oshima

2022-08-16

This is a summary report for the ETCO base model run.

Model Output

Input Data





Convergence Check

```
##   Converged     MaxGrad
## 1      TRUE 4.41683e-06
```

```
## [1] "1 NOTE:  Max data length bin: 90  < max pop len bins: 100; so will accumulate larger pop len bi
## [2] " N parameters are on or within 1% of min-max bound: 1; check results, variance may be suspect"
## [3] "N warnings: 1"
```

Fit to Model

CPUE

```
##
##  Running Runs Test Diagnosics for Index
## Plotting Residual Runs Tests
```




```
##
## Runs Test stats by Index:
## Plotting JABBA residual plot
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : Chernobyl! trL>n 6
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : Chernobyl! trL>n 6
```

```
## Warning in sqrt(sum.squares/one.delta): NaNs produced
```




```
##
## RMSE stats by Index:
```

Length Comp

#Factor	Fleet	New_Var_adj	Type	Name
4	1	0.286047	len	FISHERY

```
##
##  Running Runs Test Diagnosics for Mean length
## Plotting Residual Runs Tests

##
## Runs Test stats by Mean length:

##     Index runs.p   test  sigma3.lo sigma3.hi type
## 1 FISHERY  0.287 Passed -0.1868255 0.1868255  len

## Plotting JABBA residual plot
```




```
##
## RMSE stats by Index:

##    indices RMSE.perc nobs
## 1  FISHERY       6.1   10
## 2 Combined       6.1   10
```


FISHERY (whole catch)
Mean length
70
65
60
55
50
45
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=709
Sum of N eff.=665.4
Proportion
0.15
0.10
0.05
0.00
20
30
40
50
60
70
80
90
Length (cm)



Retrospective and Hindcasting

Retrospective

```
## Plotting Retrospective pattern
```

```
##
## Mohn's Rho stats, including one step ahead forecasts:
## Plotting Retrospective pattern
```




```
##
## Mohn's Rho stats, including one step ahead forecasts:

##   type     peel          Rho  ForecastRho
## 1    F     2020 -0.003510976 -0.003441188
## 2    F     2019 -0.004791569 -0.004718431
## 3    F     2018 -0.002512613 -0.002504181
## 4    F Combined -0.003605053 -0.003554600
```

Hindcasting

```
## Plotting Hindcast Cross-Validation (one-step-ahead)
##
##  Computing MASE with only 2 of 3  prediction residuals for Index FISHERY
##
##  Warning:  Unequal spacing of naive predictions residuals may influence the interpretation of MASE
```




```
##
## MASE stats by Index:
## Plotting Hindcast Cross-Validation (one-step-ahead)
##
## No observations in evaluation years to compute prediction residuals for Index FISHERY
##
## MASE stats by Index:
```

Recruitment Deviations

```
## Skipped SSplotrecdevs - no rec devs estimated
```

Likelihood Profile

```
## [1] "SR_LN"
```

```
## Parameter matching profile.string=SR_LN: SR_LN(R0)
```

```
## Parameter values (after subsetting based on input 'models'): 0.5, 0.7, 0.9, 1.1, 1.3, 0.780824
```

```
##
## Likelihood components showing max change as fraction of total change.
## To change which components are included, change input 'minfraction'.
```

```
##                       frac_change include                           label
## TOTAL                      1.0000    TRUE                           Total
## Catch                      0.0000   FALSE                           Catch
## Equil_catch                0.0017   FALSE                 Equilibrium catch
## Survey                     0.0173    TRUE                         Index data
## Length_comp                1.0104    TRUE                        Length data
## Recruitment                0.0000   FALSE                        Recruitment
## InitEQ_Regime              0.0000   FALSE Initital equilibrium recruitment
## Forecast_Recruitment       0.0000   FALSE              Forecast recruitment
## Parm_priors                0.0000   FALSE                             Priors
## Parm_softbounds            0.0001   FALSE                        Soft bounds
## Parm_devs                  0.0000   FALSE              Parameter deviations
## Crash_Pen                  0.0000   FALSE                     Crash penalty
```

```
## Parameter matching profile.string = 'SR_LN': 'SR_LN(R0)
## Parameter values (after subsetting based on input 'models'): 0.5, 0.7, 0.9, 1.1, 1.3, 0.780824,
```

```
## Fleet-specific likelihoods showing max change as fraction of total change.
## To change which components are included, change input 'minfraction'.
##                        frac_change include
## prof.table....c.1.3..            1    TRUE
```





Management Quantities

```
##
##  starter.sso with Bratio: SSB/SSBMSY and F: _abs_F
##
```




```
##
## Plot Comparison of stock
```




```
##
## Plot Comparison of harvest
```




```
##
## Plot Comparison of SSB
```




```
##
## Plot Comparison of F
```


MVLN
Fishing Mortality (F)
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year

```
## RStudioGD
##         2
```

Jitter




model 1
model 2
model 3
Spawning biomass (t)
0
10
20
30
1970
1980
1990
2000
2010
2020
Year


model 1
model 2
model 3
abs F; with F=sum(full Fs)
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


model 1
model 2
model 3
Age−0 recruits (1,000s)
3.0
2.5
2.0
1.5
1.0
0.5
0.0
1970
1980
1990
2000
2010
2020
Year