Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Sun Aug 28 10:44:37 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
100
80
60
40
20
0
Length (cm, beginning of the year)
CV of lengths
0.04
0.02
0
SD of lengths
3.0
1.5
0.0
0.0
1.0
2.0
3.0
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Age (yr)
Frac. mature
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Mean weight (kg) in last year
0
2
4
6
8
10
12
0
20
40
60
80
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
14
0
20
40
60
80
Female length (cm)


Spawning output
0
2
4
6
8
10
12
0
20
40
60
80
Length (cm)


Spawning output
0
2
4
6
8
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY
FISHERY
FISHERY


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Length (cm)
0
20
40
60
80
1
0.8
0.6
0.4
0.2
0
5
10
15
20
25
30
Age (yr)


Spawning biomass (mt)
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
0
1
2
3
4
5
6
7
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
0
1
2
3
4
5
6
7
Spawning biomass (mt)
2021
1967
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Landings (mt)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
Relative spawning output: B/B_MSY
2000
2020
1990
2010
1980
2021
1968
1970


Index
0
5
10
15
2016
2017
2018
2019
2020
2021
Year


Index
0
5
10
15
2016
2017
2018
2019
2020
2021
Year


Expected index
0
2
4
6
8
0
2
4
6
8
10
Observed index


Log index
1.6
1.8
2.0
2.2
2.4
2.6
2.8
2016
2017
2018
2019
2020
2021
Year


Log index
1.6
1.8
2.0
2.2
2.4
2.6
2.8
2016
2017
2018
2019
2020
2021
Year


Log expected index
2.06
2.10
2.14
2.18
1.9
2.0
2.1
2.2
2.3
Log observed index


Residual
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
2021
Year


Deviation
0.2
0.1
0.0
−0.1
−0.2
2016
2017
2018
2019
2020
2021
Year


FISHERY
Standardized index
0.0
0.2
0.4
0.6
0.8
1.0
1.2
2015
2016
2017
2018
2019
2020
2021
2022
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
0
5
10
15
20
25
30
Age


FISHERY
Sum of N input=1412.1
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
0.01
0.2
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2014
2016
2018
2020
Year


2004
N input=69.2
2012
N input=61
2019
N input=80.6
2005
N input=50.4
2014
N input=165.1
2020
N input=89
2007
N input=125.2
2015
N input=125.5
2008
N input=136.2
2016
N input=163
2009
N input=152
2017
N input=83.2
2011
N input=60.5
2018
N input=51.2
Proportion
Length (cm)


0.01
0.2
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2014
2016
2018
2020
Year


FISHERY (whole catch)
55
50
45
40
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N input=1412.1
Sum of N adj.=1015.7
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
−10
−6
2
6
10
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2014
2016
2018
2020
Year


2004
N input=69.2
N adj.=49.9
2012
N input=61
N adj.=44
2019
N input=80.6
N adj.=58.1
2005
N input=50.4
N adj.=36.4
2014
N input=165.1
N adj.=118.6
2020
N input=89
N adj.=64.1
2007
N input=125.2
N adj.=90
2015
N input=125.5
N adj.=90.2
2008
N input=136.2
N adj.=97.8
2016
N input=163
N adj.=117
2009
N input=152
N adj.=109.2
2017
N input=83.2
N adj.=59.9
2011
N input=60.5
N adj.=43.6
2018
N input=51.2
N adj.=37
Proportion
0.00
0.10
0.20
0.30
10
30
50
70
90
Length (cm)


−10
−6
2
6
10
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2014
2016
2018
2020
Year


FISHERY (whole catch)
Mean length
55
50
45
40
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.5
1.0
1.5
0
5
10
15
20
25
30
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max_likelihood
0.5
1.0
1.5
2.0
ln(DM_theta)_1
−4
−2
0
2
4
6
8
10
LnQ_base_FISHERY(1)
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
Size_inflection_FISHERY(1)
20
25
30
35
40
45
50
Size_95%width_FISHERY(1)
5
10
15
20
Density
Parameter value