American Samoa Model Checks

2022-09-21

This is a summary report for the LUKA base model run.

Model Output

Input Data





Convergence Check

```
  Converged     MaxGrad
1      TRUE 0.000346525
```

```
[1] "1 NOTE:  Max data length bin: 28  < max pop len bins: 31; so will accumulate larger pop
[2] "2 warning: poor convergence in Fmsy, final dy/dy2= 0.0016082"
[3] "3 Forecast F capped by max possible F from control file: 2.9"
[4] "4 Final gradient: 0.000346525 is larger than final_conv: 0.0001"
[5] "5 Forecast F capped by max possible F from control file: 2.9"
[6] "N warnings: 5"
```

Fit to Model

CPUE

```
Residual Runs Test (/w plot) stats by Index:
```

```
RMSE stats by Index:
```


Index
0
1
2
3
4
2016
2018
2020
Year
Index residuals
−0.6
−0.2
0.2
0.4
0.6
FISHERY
2016
2018
2020
Year


Index residuals
−0.4
−0.2
0.0
0.2
0.4
RMSE = 22.9%
FISHERY
Loess
2017
2018
2019
2020
2021
Year

Length Comp

```
Residual Runs Test (/w plot) stats by Mean length:

    Index runs.p   test   sigma3.lo  sigma3.hi type
1 FISHERY  0.065 Passed -0.03023546 0.03023546  len

RMSE stats by Index:

# A tibble: 2 x 3
  Fleet    RMSE.perc  Nobs
  <chr>        <dbl> <int>
1 FISHERY        1.2    17
2 Combined       1.2    17
```



Retrospective and Hindcasting


FISHERY (whole catch)
Mean length
22.5
22.0
21.5
21.0
20.5
20.0
2005
2010
2015
2020
Year


FISHERY
Sum of N input=9660.9
Sum of N adj.=2283.7
Proportion
0.4
0.3
0.2
0.1
0.0
14
16
18
20
22
24
26
28
Length (cm)



Retrospective

`Mohn's Rho stats, including one step ahead forecasts:`



```
Mohn's Rho stats, including one step ahead forecasts:

  type     peel          Rho  ForecastRho
1    F     2020  0.001687926  0.001657997
2    F     2019 -0.009289515 -0.009109606
3    F     2018 -0.002570552 -0.002440897
4    F Combined -0.003390714 -0.003297502
```

Hindcasting

```
Plotting Hindcast Cross-Validation (one-step-ahead)

 Computing MASE with only 2 of 3  prediction residuals for Index FISHERY

 Warning:  Unequal spacing of naive predictions residuals may influence the interpretation o
```




```
MASE stats by Index:
Plotting Hindcast Cross-Validation (one-step-ahead)

 Computing MASE with only 2 of 3  prediction residuals for Index FISHERY
```

```
 Warning:  Unequal spacing of naive predictions residuals may influence the interpretation o
```

```
MASE stats by Index:
```



Recruitment Deviations

Likelihood Profile

```
[1] "SR_LN"
```

	frac_change	include	label
TOTAL	1.0000	TRUE	Total
Catch	0.9416	TRUE	Catch
Equil_catch	0.0000	FALSE	Equilibrium catch
Survey	0.1929	TRUE	Index data
Length_comp	1.5966	TRUE	Length data
Recruitment	0.0000	FALSE	Recruitment
InitEQ_Regime	0.0000	FALSE	Initital equilibrium recruitment
Forecast_Recruitment	0.0000	FALSE	Forecast recruitment
Parm_priors	0.0150	TRUE	Priors

```
Parm_softbounds          0.0000   FALSE                    Soft bounds
Parm_devs                0.0000   FALSE          Parameter deviations
Crash_Pen                0.0000   FALSE                  Crash penalty
```





Management Quantities

starter.sso with Bratio: SSB/SSBMSY and F: _abs_F




SSB/SSB$_{MSY}$
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year
MVLN


F/F$_{MSY}$
0.0
0.2
0.4
1970
1980
1990
2000
2010
2020
Year
MVLN






```
null device
          1
```

Jitter


383.525
383.500
383.475
383.450
Likelihood
1.0
1.5
2.0
2.5
3.0
Run


Spawning biomass (t)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


abs F; with F=Z−M; for ages 3 6
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


Age−0 recruits (1,000s)
0
100
200
300
400
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3