Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Sun Jul 31 11:53:36 2022
Data_File: data.ss
Control_File: control.ss


Females
Males
Length (cm, beginning of the year)
0
10
20
30
40
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Females
Males
Length (cm, beginning of the year)
40
30
20
10
0
CV of lengths
0
0.01
0.03
0.0
0.5
1.0
1.5
SD of lengths
CV of lengths
0.04
0.02
0
SD of lengths
1.5
1.0
0.5
0.0
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Females
Males
Frac. mature
Age (yr)


Length
0
10
20
30
40
0
5
10
15
20
25
30
Age


Length
0
10
20
30
40
0
5
10
15
20
25
30
Age


Length
0
10
20
30
40
0
5
10
15
20
25
30
Age


Length
0
10
20
30
40
0
5
10
15
20
25
30
Age


Females
Males
Mean weight (kg) in last year
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
Length (cm)


Females
Males
Mean weight (kg) in last year
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
Female weight (kg)


2.0
1.5
1.0
0.5
0.0
Fecundity
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
Female length (cm)


2.0
1.5
1.0
0.5
0.0
Spawning output
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Spawning output
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY (f)
FISHERY (m)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
0
5
10
15
20
25
30
Age (yr)
0.8
0.6
0.4


Length (cm)
0
10
20
30
40
0
5
10
15
20
25
30
Age (yr)
0.8
0.6
0.4


Spawning biomass (mt)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.4
0.2
0.0
−0.2
−0.6
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.5
0.5
−0.5
−1.5
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
1.0
0.5
0.0
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
0.0
0.5
1.0
1.5
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
Spawning biomass (mt)
0.0
0.5
1.0
1.5
1967
2021
1997
2000
1999
2006
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Log recruitment deviation
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)
1967
2021
1997
2000
1999
2006


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Relative spawning output: B/B_MSY
1980
1990
2000
2010
2020
2021
1967
1970


Index
0
1
2
3
4
5
6
7
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
1
2
3
4
5
6
7
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0.0
0.5
1.0
1.5
2.0
Observed index


Log index
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.2
0.0
−0.2
−0.4
−0.6
−2.5
−2.0
−1.5
−1.0
−0.5
0.0
0.5
Log observed index


Effective catchability
0.5452870
0.5452874
0.5452878
1.0
1.5
2.0
Vulnerable biomass


Residual
1
0
−1
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
1990
1995
2000
2005
2010
2015
2020
Year


0.1
2
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


Mean age (yr)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year
Females
Males


200
600
1000
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


200
600
1000
Age
30
25
20
15
10
5
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


Mean age (yr)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year
Females
Males


Age
Year
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
0.5
0.502
0.504
0.506
0.508
0.51
0.512
0.514
0.516
0.518
0.52
0.522


100
300
500
Length
50
40
30
20
10
0
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


100
300
500
Length
Year
0
10
20
30
40
50
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021


Mean length (cm)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year
Females
Males


100
300
500
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


100
300
500
Length
Year
0
10
20
30
40
50
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021


Mean length (cm)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year
Females
Males


Length
Year
10
20
30
40
1970
1980
1990
2000
2010
2020
0.1
0.2
0.5
0.4
0.3
0.6
0.5
0.6
0.5
0.7
0.5
0.6
0.6
0.7
0.7
0.5
0.6
0.7
0.8
0.6
0.5
0.8
0.7
0.3
0.4
0.2
0.1
0.1
0.2


Numbers at age at equilibrium
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0
5
10
15
20
25
30
Age
Females
Males


FISHERY
Sum of N adj.=148.8
Proportion
0.00
0.05
0.10
0.15
0.20
15
20
25
30
35
40
Length (cm)


FISHERY
0.01
0.15
0.25
Length (cm)
40
38
36
34
32
30
28
26
24
22
20
18
16
2012
2017
Year


2007
N adj.=48
2012
N adj.=42
2017
N adj.=58.8
0.25
0.20
0.15
0.10
0.05
0.00
15
20
25
30
35
40
Proportion
Length (cm)


0.01
0.15
0.25
40
36
32
28
24
20
16
Length (cm)
2007
2012
2017
Year


FISHERY (whole catch)
30
29
28
27
Mean length
2008
2010
2012
2014
2016
Year


FISHERY
Sum of N adj.=148.8
Sum of N eff.=520.1
Proportion
0.20
0.15
0.10
0.05
0.00
15
20
25
30
35
40
Length (cm)


FISHERY
−2
0.1
2
Length (cm)
40
38
36
34
32
30
28
26
24
22
20
18
16
2012
2017
Year


2007
N adj.=48
N eff.=88.7
2012
N adj.=42
N eff.=111
2017
N adj.=58.8
N eff.=320.4
0.00
0.10
0.20
15
20
25
30
35
40
Proportion
Length (cm)


−2
0.1
2
40
36
32
28
24
20
16
Length (cm)
2007
2012
2017
Year


Effective sample size
0
50
150
250
350
0
10
20
30
40
50
60
Observed sample size
harmonic mean
arithmetic mean
2017
2012
2007


FISHERY (whole catch)
30
29
28
27
Mean length
2008
2010
2012
2014
2016
Year


Equilibrium yield (mt)
0.00
0.10
0.20
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.00
0.10
0.20
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.00
0.10
0.20
0.30
0
1
2
3
4
Total biomass (mt)


Yield per recruit (kg)
0.00
0.10
0.20
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
5
Size_inflection_FISHERY(1)
16
18
20
22
24
26
28
30
InitF_seas_1_flt_1FISHERY
0.00
0.05
0.10
0.15
0.20
0.25
0.30
Size_95%width_FISHERY(1)
0
2
4
6
8
10
12
14
LnQ_base_FISHERY(1)
−4
−2
0
2
4
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value