Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Wed Aug 24 16:10:44 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
20
40
60
80
100
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Length (cm, beginning of the year)
100
80
60
40
20
0
CV of lengths
0.06
0.04
0.02
0
SD of lengths
4
2
0
Age (yr)
CV of lengths
0
0.02
0.04
0.06
SD of lengths
0
1
2
3
4
5
Length (cm, beginning of the year)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Mean weight (kg) in last year
0
5
10
15
20
0
20
40
60
80
100
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
100
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
20
Female weight (kg)


Fecundity
0
5
10
15
20
0
5
10
15
20
Female weight (kg)


Fecundity
0
5
10
15
20
0
20
40
60
80
100
Female length (cm)


Spawning output
0
5
10
15
20
0
20
40
60
80
100
Length (cm)


Spawning output
15
10
5
0
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY
FISHERY
FISHERY


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Length (cm)
0
20
40
60
80
100
1
0.8
0.6
0.2
0
5
10
15
20
25
30
Age (yr)


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.00
0.05
0.10
0.15
0.20
0.25
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
2
4
6
8
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
2021
1967
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
2
4
6
8
Spawning biomass (mt)


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.00
0.04
0.08
0.12
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Relative spawning output: B/B_MSY
2010
2000
1990
1980
1968
1970


Index
0
5
10
15
20
2016
2017
2018
2019
2020
2021
Year


Index
0
5
10
15
20
2016
2017
2018
2019
2020
2021
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0
1
2
3
4
5
6
Observed index


Log index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
2016
2017
2018
2019
2020
2021
Year


Log index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
2016
2017
2018
2019
2020
2021
Year


Log expected index
1.20
1.21
1.22
1.23
1.24
1.25
0.8
1.0
1.2
1.4
1.6
1.8
Log observed index


Residual
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
2021
Year


Deviation
0.4
0.2
0.0
−0.2
−0.4
2016
2017
2018
2019
2020
2021
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2015
2016
2017
2018
2019
2020
2021
2022
Year


0.1
2
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=339.8
Proportion
0.00
0.05
0.10
0.15
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
0.01
0.2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


2007
N adj.=32.7
2013
N adj.=17.5
2008
N adj.=28.3
2014
N adj.=25.1
2009
N adj.=38.6
2015
N adj.=34.7
2010
N adj.=14.4
2016
N adj.=28.2
2011
N adj.=21.5
2017
N adj.=36.5
2012
N adj.=32.6
2019
N adj.=29.7
0.00
0.10
0.20
20
40
60
80
Proportion
Length (cm)


0.01
0.2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


FISHERY (whole catch)
60
55
50
45
Mean length
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=339.8
Sum of N eff.=1102.8
Proportion
0.15
0.10
0.05
0.00
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
−2
0.1
2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


2007
N adj.=32.7
N eff.=49.4
2013
N adj.=17.5
N eff.=78.8
2008
N adj.=28.3
N eff.=126.2
2014
N adj.=25.1
N eff.=18.6
2009
N adj.=38.6
N eff.=277.9
2015
N adj.=34.7
N eff.=238.8
2010
N adj.=14.4
N eff.=75.6
2016
N adj.=28.2
N eff.=31.3
2011
N adj.=21.5
N eff.=33.2
2017
N adj.=36.5
N eff.=58.1
2012
N adj.=32.6
N eff.=58.7
2019
N adj.=29.7
N eff.=56.3
0.30
0.20
0.10
0.00
20
40
60
80
Proportion
Length (cm)


−2
0.1
2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


Effective sample size
0
50
150
250
0
10
20
30
40
Observed sample size
harmonic mean
arithmetic mean
2009
2015
2008
2010
2013
2019
2012
2017
2007
2011
2016
2014


FISHERY (whole catch)
Mean length
60
55
50
45
2008
2010
2012
2014
2016
2018
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.4
0.8
0
10
20
30
40
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
5
LnQ_base_FISHERY(1)
−4
−2
0
2
Size_inflection_FISHERY(1)
20
25
30
35
40
45
50
55
Size_95%width_FISHERY(1)
5
10
15
20
25
30
Density
Parameter value