Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Mon Aug 08 13:30:45 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Length (cm, beginning of the year)
0
20
40
60
80
100
CV of lengths
0
0.02
0.04
SD of lengths
0
1
2
3
4


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
Age


Mean weight (kg) in last year
0
5
10
15
20
0
20
40
60
80
100
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
100
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
20
Female weight (kg)


Fecundity
0
5
10
15
20
0
5
10
15
20
Female weight (kg)


Fecundity
0
5
10
15
20
0
20
40
60
80
100
Female length (cm)


Spawning output
0
5
10
15
20
0
20
40
60
80
100
Length (cm)


Spawning output
0
5
10
15
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY
FISHERY
FISHERY
FISHERY
FISHERY


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Length (cm)
0
20
40
60
80
100
1
0.8
0.6
0.2
0
5
10
15
20
25
30
Age (yr)


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
30
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.00
0.05
0.10
0.15
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.6
0.2
−0.2
−0.6
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
1
2
3
4
5
6
0
2
4
6
8
10
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
2012
2021
1960
2010
Recruitment (1,000s)
0
1
2
3
4
5
6
0
2
4
6
8
10
Spawning biomass (mt)


2012
2021
2010
1967
0.5
0.0
−0.5
0.0
0.2
0.4
0.6
0.8
1.0
Log recruitment deviation
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.00
0.02
0.04
0.06
0.08
0.10
1970
1980
1990
2000
2010
2020
Year


SPR
1.0
0.8
0.6
0.4
0.2
0.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Relative spawning output: B/B_MSY
2010
2000
1980
1960


Index
0
5
10
15
20
2016
2017
2018
2019
2020
2021
Year


Index
0
5
10
15
20
2016
2017
2018
2019
2020
2021
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0
1
2
3
4
5
6
Observed index


Log index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
2016
2017
2018
2019
2020
2021
Year


Log index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
2016
2017
2018
2019
2020
2021
Year


Log expected index
1.200
1.210
1.220
0.8
1.0
1.2
1.4
1.6
1.8
Log observed index


Residual
1.0
0.5
0.0
−0.5
−1.0
2016
2017
2018
2019
2020
2021
Year


Deviation
0.6
0.4
0.2
0.0
−0.2
−0.6
2016
2017
2018
2019
2020
2021
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2015
2016
2017
2018
2019
2020
2021
2022
Year


0.1
4
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
4
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=975.8
0.15
0.10
0.05
0.00
Proportion
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
0.01
0.2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


2007
N adj.=93.4
2013
N adj.=50
2008
N adj.=80.7
2014
N adj.=71.7
2009
N adj.=110.2
2015
N adj.=99.3
2010
N adj.=41.2
2016
N adj.=80.6
2011
N adj.=61.3
2017
N adj.=104.2
2012
N adj.=93.2
2019
N adj.=89.9
0.00
0.10
0.20
20
40
60
80
Proportion
Length (cm)


0.01
0.2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


FISHERY (whole catch)
60
55
50
45
Mean length
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=975.8
Sum of N eff.=1290.9
Proportion
0.15
0.10
0.05
0.00
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
−2
0.1
2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


2007
N adj.=93.4
N eff.=58.3
2013
N adj.=50
N eff.=90.2
2008
N adj.=80.7
N eff.=115
2014
N adj.=71.7
N eff.=25.4
2009
N adj.=110.2
N eff.=282
2015
N adj.=99.3
N eff.=302.4
2010
N adj.=41.2
N eff.=72.3
2016
N adj.=80.6
N eff.=43.7
2011
N adj.=61.3
N eff.=39.4
2017
N adj.=104.2
N eff.=91.4
2012
N adj.=93.2
N eff.=75.8
2019
N adj.=89.9
N eff.=94.9
Proportion
0.00
0.10
0.20
0.30
20
40
60
80
Length (cm)


−2
0.1
2
Length (cm)
90
80
70
60
50
40
30
20
2008
2010
2012
2014
2016
2019
Year


Effective sample size
0
50
150
250
0
20
40
60
80
100
120
Observed sample size
harmonic mean
arithmetic mean
2015
2009
2008
2019
2017
2012
2007
2016
2013
2010
2011
2014


FISHERY (whole catch)
Mean length
60
55
50
45
2008
2010
2012
2014
2016
2018
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.4
0.8
1.2
0
10
20
30
40
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
Abundance indices
Length compositions
FISHERY
FISHERY
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max_likelihood
1
2
3
4
5
Size_inflection_FISHERY(1)
20
25
30
35
40
45
50
55
InitF_seas_1_flt_1FISHERY
0.00
0.05
0.10
0.15
0.20
0.25
0.30
Size_95%width_FISHERY(1)
5
10
15
20
25
30
LnQ_base_FISHERY(1)
−4
−2
0
2
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value