Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Thu Jul 07 08:12:29 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Length (cm, beginning of the year)
0
20
40
60
80
100
CV of lengths
0.04
0.08
0.12
SD of lengths
2
4
6
8
10


Age (yr)
0
5
10
15
20
25
30
100
80
60
40
20
0
Length (cm, beginning of the year)
Mean weight
10
5
0
1.0
0.5
0.0
Frac. mature
Age (yr)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Mean weight (kg) in last year
15
10
5
0
0
20
40
60
80
100
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
100
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
Female weight (kg)


15
10
5
0
Fecundity
0
5
10
15
Female weight (kg)


Fecundity
15
10
5
0
0
20
40
60
80
100
Female length (cm)


Spawning output
15
10
5
0
0
20
40
60
80
100
Length (cm)


Spawning output
0
1
2
3
4
5
6
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Length (cm)
0
20
40
60
80
100
0
5
10
15
20
25
30
Age (yr)
1
0.8
0.6
0.4
0.2


Length (cm)
0
20
40
60
80
100
0
5
10
15
20
25
30
Age (yr)
1
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
5
10
15
1985
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0
5
10
15
20
1985
1990
1995
2000
2005
2010
2015
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.1
0.2
0.3
0.4
1985
1990
1995
2000
2005
2010
2015
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.1
0.2
0.3
0.4
0.5
1985
1990
1995
2000
2005
2010
2015
2020
Year


Total biomass (mt)
0
10
20
30
40
1985
1990
1995
2000
2005
2010
2015
2020
Year


Summary biomass (mt)
0
10
20
30
40
1985
1990
1995
2000
2005
2010
2015
2020
Year


Age−0 recruits (1,000s)
0.0
1.0
2.0
3.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
1985
1990
1995
2000
2005
2010
2015
2020
Year


Summary Fishing Mortality
0
1
2
3
4
5
6
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0
5
10
15
20
1990
1995
2000
2005
2010
2015
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.5
0.0
−0.5
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−1.0
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0.0
0.2
0.4
0.6
0.8
1.0
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
2010
2011
2012
2021
1987
2016
1996
2008
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0.0
0.2
0.4
0.6
0.8
1.0
Spawning biomass (mt)


2010
2011
2012
1987
2021
2016
1996
2008
0.5
0.0
−0.5
Log recruitment deviation
0.00
0.02
0.04
0.06
0.08
0.10
Spawning output (relative to B_0)


Landings (mt)
0
1
2
3
4
1985
1990
1995
2000
2005
2010
2015
2020
Year


Landings (mt)
0
1
2
3
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
1
2
3
4
1985
1990
1995
2000
2005
2010
2015
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1990
1995
2000
2005
2010
2015
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1990
1995
2000
2005
2010
2015
2020
Year


1987
2021
Fishing intensity: 1−SPR
Relative spawning output: B/B_MSY
0.0
0.2
0.4
0.6
0.8
1.0


Index
0
5
10
15
20
25
30
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
30
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0
2
4
6
8
10
12
0
5
10
15
Observed index


Log index
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
1.6
1.8
2.0
2.2
2.4
1.0
1.5
2.0
2.5
Log observed index


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
Year


Expected index
0
1
2
3
4
5
6
7
0
2
4
6
8
Observed index


Log index
0
1
2
3
1990
1995
2000
2005
Year


Log index
0
1
2
3
1990
1995
2000
2005
Year


Log expected index
1.0
1.2
1.4
1.6
1.8
1.0
1.5
2.0
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
Year


Deviation
0.5
0.0
−0.5
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0.0
0.5
1.0
1.5
2.0
1990
1995
2000
2005
2010
2015
2020
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


0.1
2
100
80
60
40
20
0
Length
1987
1991
1995
1999
2003
2007
2011
2015
2019
Year


200
600
1000
100
80
60
40
20
0
Length
1987
1991
1995
1999
2003
2007
2011
2015
2019
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=2147.9
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
20
40
60
80
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
90
80
70
60
50
40
30
20
10
2004
2007
2009
2011
2013
2015
2017
2019
Year


2004
N adj.=68.9
2013
N adj.=441.8
2019
N adj.=80.6
2007
N adj.=119.3
2014
N adj.=190.7
2020
N adj.=89
2008
N adj.=134.4
2015
N adj.=180.6
2009
N adj.=152
2016
N adj.=163
2011
N adj.=135.2
2017
N adj.=83.2
2012
N adj.=257.9
2018
N adj.=51.2
Proportion
Length (cm)


0.01
0.2
0.4
Length (cm)
90
80
70
60
50
40
30
20
10
2004
2007
2009
2011
2013
2015
2017
2019
Year


FISHERY (whole catch)
55
50
45
40
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=2147.9
Sum of N eff.=877.2
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
20
40
60
80
Length (cm)


FISHERY
−25
−15
5
15
25
Length (cm)
90
80
70
60
50
40
30
20
10
2004
2007
2009
2011
2013
2015
2017
2019
Year


2004
N adj.=68.9
N eff.=61.7
2013
N adj.=441.8
N eff.=92.6
2019
N adj.=80.6
N eff.=50.3
2007
N adj.=119.3
N eff.=65.7
2014
N adj.=190.7
N eff.=170.8
2020
N adj.=89
N eff.=39.2
2008
N adj.=134.4
N eff.=55.3
2015
N adj.=180.6
N eff.=90.6
2009
N adj.=152
N eff.=53.4
2016
N adj.=163
N eff.=81
2011
N adj.=135.2
N eff.=10
2017
N adj.=83.2
N eff.=69.9
2012
N adj.=257.9
N eff.=19.8
2018
N adj.=51.2
N eff.=17.1
Proportion
0.0
0.1
0.2
0.3
0.4
20
40
60
80
Length (cm)


−25
−15
5
15
25
90
80
70
60
50
40
30
20
10
Length (cm)
2004
2007
2009
2011
2013
2015
2017
2019
Year


Effective sample size
0
50
100
150
0
100
200
300
400
500
Observed sample size
2014
2015
2016
2013
2017
2004
2007
2008
2009
2019
2020
2018
2011
2012
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
55
50
45
40
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
2.0
2.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
2.0
2.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
1.0
2.0
0
1
2
3
4
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
0.8
1.0
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_inflection_FISHERY(1)
15
20
25
30
35
40
45
InitF_seas_1_flt_1FISHERY
0.0
0.5
1.0
1.5
2.0
Size_95%width_FISHERY(1)
0
2
4
6
8
10
12
14
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Density
Parameter value