American Samoa Model Checks

Meg Oshima

2022-08-11

This is a summary report for the CALU base model run.

Model Output

Input Data





Convergence Check

```
##   Converged     MaxGrad
## 1      TRUE 7.83285e-05
```

```
## [1] "1 NOTE:  Max data length bin: 65  < max pop len bins: 72; so will accumulate larger pop len bins
## [2] "2 warning: poor convergence in Fmsy, final dy/dy2= -0.0218076"
## [3] "N warnings: 2"
```

Fit to Model

CPUE

```
##
##  Running Runs Test Diagnosics for Index
## Plotting Residual Runs Tests
```




```
##
## Runs Test stats by Index:
## Plotting JABBA residual plot
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : span too small. fe
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : pseudoinverse used
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : neighborhood radius

## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : reciprocal conditio

## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : There are other nea
```




```
##
## RMSE stats by Index:
```

Length Comp

#Factor	Fleet	New_Var_adj	Type	Name
4	1	0.33815	len	FISHERY

```
##
##  Running Runs Test Diagnosics for Mean length
## Plotting Residual Runs Tests

##
## Runs Test stats by Mean length:

##     Index runs.p   test  sigma3.lo sigma3.hi type
## 1 FISHERY  0.358 Passed -0.1458023 0.1458023  len
```

```
## Plotting JABBA residual plot
```




```
##
## RMSE stats by Index:

##    indices RMSE.perc nobs
## 1  FISHERY       4.1    7
## 2 Combined       4.1    7
```


FISHERY (whole catch)
Mean length
48
46
44
42
40
38
36
34
2008
2010
2012
2014
2016
2018
Year


FISHERY
Sum of N adj.=474.4
Sum of N eff.=477.8
Proportion
0.30
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
Length (cm)



Retrospective and Hindcasting

Retrospective

```
## Plotting Retrospective pattern
```

```
##
## Mohn's Rho stats, including one step ahead forecasts:
## Plotting Retrospective pattern
```




```
##
## Mohn's Rho stats, including one step ahead forecasts:

##   type     peel          Rho ForecastRho
## 1    F     2020 -0.002777689 0.001789017
## 2    F     2019  0.024777045 0.015005907
## 3    F     2018  0.000000000 0.000000000
## 4    F Combined  0.007333119 0.005598308
```

Hindcasting

```
## Plotting Hindcast Cross-Validation (one-step-ahead)
##
##  Computing MASE with only 2 of 3  prediction residuals for Index FISHERY
##
##  Warning:  Unequal spacing of naive predictions residuals may influence the interpretation of MASE
```




```
##
## MASE stats by Index:
## Plotting Hindcast Cross-Validation (one-step-ahead)
##
## No observations in evaluation years to compute prediction residuals for Index FISHERY
##
## MASE stats by Index:
```

Recruitment Deviations





Likelihood Profile

```
## [1] "SR_LN"
```

```
## Parameter matching profile.string=SR_LN: SR_LN(R0)
```

```
## Parameter values (after subsetting based on input 'models'): 2, 2.2, 2.4, 2.6, 2.8, 3, 3.2, 3.4, 3.6
```

```
##
## Likelihood components showing max change as fraction of total change.
## To change which components are included, change input 'minfraction'.
```

```
##                       frac_change include                            label
## TOTAL                      1.0000    TRUE                            Total
## Catch                      0.1322    TRUE                            Catch
## Equil_catch                0.0000   FALSE                Equilibrium catch
## Survey                     0.0331    TRUE                       Index data
## Length_comp                1.2264    TRUE                      Length data
## Recruitment                0.1273    TRUE                      Recruitment
## InitEQ_Regime              0.0000   FALSE Initital equilibrium recruitment
## Forecast_Recruitment       0.0000   FALSE             Forecast recruitment
## Parm_priors                0.0000   FALSE                           Priors
## Parm_softbounds            0.0000   FALSE                      Soft bounds
## Parm_devs                  0.0000   FALSE             Parameter deviations
## Crash_Pen                  0.0000   FALSE                    Crash penalty
```

```
## Parameter matching profile.string = 'SR_LN': 'SR_LN(R0)
## Parameter values (after subsetting based on input 'models'): 2, 2.2, 2.4, 2.6, 2.8, 3, 3.2, 3.4, 3.6
```

```
## Fleet-specific likelihoods showing max change as fraction of total change.
## To change which components are included, change input 'minfraction'.
##                         frac_change include
## prof.table....c.1.3..           1    TRUE
```


Change in −log−likelihood
0
1
2
3
4
1.5
2.0
2.5
3.0
3.5
SR_LN(R0)
Total
Length data
Catch
Recruitment
Index data



Management Quantities

```
##
##  starter.sso with Bratio: SSB/SSBMSY and F: _abs_F
##
```




```
##
## Plot Comparison of stock
```




```
##
## Plot Comparison of harvest
```




```
##
## Plot Comparison of SSB
```




```
##
## Plot Comparison of F
```


MVLN
Fishing Mortality (F)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year

```
## RStudioGD
##         2
```

Jitter




Spawning biomass (t)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year
model 1
model 2
model 3


model 1
model 2
model 3
abs F; with F=sum(full Fs)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


model 1
model 2
model 3
Age−0 recruits (1,000s)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year