Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Tue Jul 12 07:59:16 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
2
4
6
8
10
12
Age (yr)


Age (yr)
0
2
4
6
8
10
12
Length (cm, beginning of the year)
0
20
40
60
80
100
CV of lengths
0
0.04
0.08
SD of lengths
0
2
4
6
8


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
2
4
6
8
10
12
Age


Length
100
80
60
40
20
0
0
2
4
6
8
10
12
Age


Mean weight (kg) in last year
20
15
10
5
0
0
20
40
60
80
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
20
40
60
80
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
Female weight (kg)


Fecundity
20
15
10
5
0
0
5
10
15
Female weight (kg)


Fecundity
0
5
10
15
20
0
20
40
60
80
Female length (cm)


Spawning output
20
15
10
5
0
0
20
40
60
80
Length (cm)


Spawning output
10
8
6
4
2
0
0
2
4
6
8
10
12
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
12
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Length (cm)
0
20
40
60
80
0
2
4
6
8
10
12
Age (yr)
0.8
0.6
0.4
0.2


Length (cm)
0
20
40
60
80
0
2
4
6
8
10
12
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
1.0
0.5
0.0
−0.5
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
5
10
15
0
1
2
3
4
5
6
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
1985
1984
1983
2021
2007
Recruitment (1,000s)
0
5
10
15
0
1
2
3
4
5
6
Spawning biomass (mt)


1985
1984
1983
2007
2021
7
1.0
0.5
0.0
−0.5
−1.0
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Log recruitment deviation
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
Relative spawning output: B/B_MSY
1967
2021


Index
0
5
10
15
20
25
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0
2
4
6
8
10
12
Observed index


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.6
0.8
1.0
1.2
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
2.5
Log observed index


Residual
4
2
0
−2
−4
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.5
1.0
0.5
−0.5
−1.5
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
1990
1995
2000
2005
Year


Index
0
5
10
15
1990
1995
2000
2005
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
1
2
3
4
5
6
7
Observed index


Log index
3
2
1
0
−1
−2
−3
−4
1990
1995
2000
2005
Year


Log index
3
2
1
0
−1
−2
−3
−4
1990
1995
2000
2005
Year


Log expected index
0.5
0.6
0.7
0.8
0.9
1.0
1.1
−2
−1
0
1
2
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


5
15
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


2
6
10
14
Age
12
11
10
9
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


5
15
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
1
2
3
4
5
0
2
4
6
8
10
12
Age


FISHERY
Sum of N adj.=244.2
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
20
40
60
80
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
0
10
20
30
40
50
60
70
80
2009
2012
2014
2017
Year


2007
N adj.=38.4
2017
N adj.=24.1
2009
N adj.=35.9
2012
N adj.=50.6
2013
N adj.=25.9
2014
N adj.=27
2015
N adj.=42.3
0.30
0.20
0.10
0.00
0
20
40
60
80
Proportion
Length (cm)


0.01
0.2
0.4
Length (cm)
0
10
20
30
40
50
60
70
80
2007
2009
2012
2014
2017
Year


FISHERY (whole catch)
Mean length
50
45
40
35
2008
2010
2012
2014
2016
Year


FISHERY
Sum of N adj.=244.2
Sum of N eff.=241.7
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
20
40
60
80
Length (cm)


FISHERY
−300
100
300
Length (cm)
0
10
20
30
40
50
60
70
80
2009
2012
2014
2017
Year


2007
N adj.=38.4
N eff.=35.8
2017
N adj.=24.1
N eff.=46.7
2009
N adj.=35.9
N eff.=12.2
2012
N adj.=50.6
N eff.=13.8
2013
N adj.=25.9
N eff.=18.9
2014
N adj.=27
N eff.=88.9
2015
N adj.=42.3
N eff.=25.5
Proportion
0.00
0.10
0.20
0.30
0
20
40
60
80
Length (cm)


−300
100
300
Length (cm)
0
10
20
30
40
50
60
70
80
2007
2009
2012
2014
2017
Year


Effective sample size
0
20
40
60
80
100
0
10
20
30
40
50
Observed sample size
2014
2017
2007
2015
2013
2009
2012
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
50
45
40
35
2008
2010
2012
2014
2016
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


MSY
B target
SPR target
Current
Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Surplus production (mt)
0
1
2
3
4
5
0
5
10
15
20
25
Total biomass (mt)


Yield per recruit (kg)
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_inflection_FISHERY(1)
20
30
40
50
60
InitF_seas_1_flt_1FISHERY
0.0
0.5
1.0
1.5
2.0
Size_95%width_FISHERY(1)
0
5
10
15
20
25
Density
LnQ_base_FISHERY(1)
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−4
−2
0
2
4
Parameter value