Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Fri Jul 08 10:43:32 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Length (cm, beginning of the year)
0
20
40
60
80
100
CV of lengths
0.04
0.08
0.12
SD of lengths
2
4
6
8
10


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Mean weight (kg) in last year
15
10
5
0
0
20
40
60
80
100
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
100
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
Female weight (kg)


15
10
5
0
Fecundity
0
5
10
15
Female weight (kg)


Fecundity
0
5
10
15
0
20
40
60
80
100
Female length (cm)


Spawning output
15
10
5
0
0
20
40
60
80
100
Length (cm)


Spawning output
0
1
2
3
4
5
6
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
100
Length (cm)


Length (cm)
0
20
40
60
80
100
0
5
10
15
20
25
30
Age (yr)
1
0.8
0.6
0.4
0.2


Length (cm)
0
20
40
60
80
100
0
5
10
15
20
25
30
Age (yr)
1
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
−1.0
0.0
0.5
1.0
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1
0
−1
−2
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1 - SE(\hat{r}_y)^2/\sigma_R^2$
1.0
0.5
0.0
−1.0
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
0
2
4
6
8
10
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
0
2
4
6
8
10
011
2010
2005
2002
12
021
1967
2004
2008


Log recruitment deviation
Spawning output (relative to B_0)
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
0.0
0.2
0.4
0.6
0.8
1.0
011
2010
12
2005
2002
021
1967
1996
2004
2008


Landings (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Relative spawning output: B/B_MSY
021
1969


Index
0
5
10
15
20
25
30
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
30
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0
2
4
6
8
10
12
0
5
10
15
Observed index


Log index
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
1.4
1.6
1.8
2.0
2.2
2.4
1.0
1.5
2.0
2.5
Log observed index


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1.0
0.5
0.0
−0.5
−1.0
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
Year


Expected index
Observed index
0
1
2
3
4
5
6
0
2
4
6
8


Log index
0
1
2
3
1990
1995
2000
2005
Year


Log index
0
1
2
3
1990
1995
2000
2005
Year


Log expected index
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.0
1.5
2.0
Log observed index


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
Year


Deviation
0.5
0.0
−0.5
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0.0
0.5
1.0
1.5
2.0
1990
1995
2000
2005
2010
2015
2020
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


100
300
500
700
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=650.8
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
20
40
60
80
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
90
80
70
60
50
40
30
20
10
2004
2007
2009
2011
2013
2015
2017
2019
Year


2004
N adj.=20.9
2013
N adj.=133.9
2019
N adj.=24.4
2007
N adj.=36.1
2014
N adj.=57.8
2020
N adj.=27
2008
N adj.=40.7
2015
N adj.=54.7
2009
N adj.=46.1
2016
N adj.=49.4
2011
N adj.=41
2017
N adj.=25.2
2012
N adj.=78.1
2018
N adj.=15.5
Proportion
Length (cm)


0.01
0.2
0.4
Length (cm)
90
80
70
60
50
40
30
20
10
2004
2007
2009
2011
2013
2015
2017
2019
Year


FISHERY (whole catch)
55
50
45
40
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=650.8
Sum of N eff.=429.7
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
20
40
60
80
Length (cm)


FISHERY
−15
5
15
Length (cm)
90
80
70
60
50
40
30
20
10
2004
2007
2009
2011
2013
2015
2017
2019
Year


2004
N adj.=20.9
N eff.=34.9
2013
N adj.=133.9
N eff.=50.4
2019
N adj.=24.4
N eff.=26.9
2007
N adj.=36.1
N eff.=24.8
2014
N adj.=57.8
N eff.=30.6
2020
N adj.=27
N eff.=19.8
2008
N adj.=40.7
N eff.=31.9
2015
N adj.=54.7
N eff.=41.3
2009
N adj.=46.1
N eff.=18.6
2016
N adj.=49.4
N eff.=62.4
2011
N adj.=41
N eff.=10.6
2017
N adj.=25.2
N eff.=41.4
2012
N adj.=78.1
N eff.=18.2
2018
N adj.=15.5
N eff.=18.1
Proportion
Length (cm)
0.0
0.1
0.2
0.3
0.4
20
40
60
80


15
5
15
90
80
70
60
50
40
30
20
10
Length (cm)
2004
2007
2009
2011
2013
2015
2017
2019
Year


Effective sample size
Observed sample size
0
10
30
50
70
0
20
40
60
80
100
120
140
2016
2013
2017
2015
2004
2008
2014
2019
2007
2020
2009
2012
2018
2011
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
55
50
45
40
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
−0.5
0.5
1.5
2.5
0
10
20
30
40
Total biomass (mt)


Yield per recruit (kg)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_95%width_FISHERY(1)
0
2
4
6
8
10
12
14
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
15
20
25
30
35
40
45
Density
Parameter value