Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Fri Jul 01 13:20:48 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
2
4
6
8
10
Age (yr)


Age (yr)
0
2
4
6
8
10
Length (cm, beginning of the year)
100
80
60
40
20
0
CV of lengths
0.08
0.04
0
SD of lengths
8
4
0
CV of lengths
0.04
0.08
SD of lengths
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
2
4
6
8
10
12
Age


Length
100
80
60
40
20
0
0
2
4
6
8
10
12
Age


Mean weight (kg) in last year
20
15
10
5
0
0
20
40
60
80
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
20
40
60
80
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
Female weight (kg)


Fecundity
20
15
10
5
0
0
5
10
15
Female weight (kg)


Fecundity
0
5
10
15
20
0
20
40
60
80
Female length (cm)


Spawning output
20
15
10
5
0
0
20
40
60
80
Length (cm)


Spawning output
0
2
4
6
8
0
2
4
6
8
10
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
10
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Length (cm)
0
20
40
60
80
0
2
4
6
8
10
Age (yr)
0.8
0.6
0.4
0.2


Length (cm)
0
20
40
60
80
0
2
4
6
8
10
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
1.0
2.0
3.0
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
5
1990
2000
2010
2020
Year


Total biomass (mt)
0
1
2
3
4
5
6
1990
2000
2010
2020
Year


Summary biomass (mt)
0
1
2
3
4
5
6
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
6
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
2
4
6
8
10
12
14
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
1
2
3
4
1985
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
1985
1990
1995
2000
2005
2010
2015
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.4
0.2
0.0
−0.2
−0.6
1980
1990
2000
2010
2020
Year


Log recruitment deviation
−1.0
−0.5
0.0
0.5
1.0
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
1.0
0.6
0.2
−0.2
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0
1
2
3
4
5
6
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0
1
2
3
4
5
6
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Spawning biomass (mt)
1986
2021
2007
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
Log recruitment deviation
1986
2021
2007
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
Relative spawning output: B/B_MSY
2021
1987


Index
0
5
10
15
20
25
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
Observed index
0
1
2
3
4
5
6
0
2
4
6
8
10
12


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log index
3
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.0
0.5
1.0
1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
2.5
Log observed index


Effective catchability
1.975780
1.975784
1.975788
0.5
1.0
1.5
2.0
2.5
3.0
Vulnerable biomass


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
1
0
−1
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
1990
1995
2000
2005
Year


Index
0
5
10
15
1990
1995
2000
2005
Year


Expected index
Observed index
0
1
2
3
4
5
6
7


Log index
3
2
1
0
−1
−2
−3
−4
1990
1995
2000
2005
Year


Log index
3
2
1
0
−1
−2
−3
−4
1990
1995
2000
2005
Year


Log expected index
0.4
0.6
0.8
1.0
1.2
1.4
−2
−1
0
1
2
Log observed index


Residual
2
1
0
−1
−2
1990
1995
2000
2005
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


0.1
4
8
Age
11
10
9
8
7
6
5
4
3
2
1
0
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


0.1
2
Age
11
10
9
8
7
6
5
4
3
2
1
0
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


0.1
4
8
100
80
60
40
20
0
Length
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


0.1
2
100
80
60
40
20
0
Length
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


Numbers at age at equilibrium
0
1
2
3
4
0
2
4
6
8
10
Age


FISHERY
Sum of N adj.=348.8
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
20
40
60
80
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
0
10
20
30
40
50
60
70
80
2009
2012
2014
2017
Year


2007
N adj.=54.8
2017
N adj.=34.4
2009
N adj.=51.2
2012
N adj.=72.2
2013
N adj.=37
2014
N adj.=38.6
2015
N adj.=60.5
0.30
0.20
0.10
0.00
0
20
40
60
80
Proportion
Length (cm)


0.01
0.2
0.4
Length (cm)
0
10
20
30
40
50
60
70
80
2007
2009
2012
2014
2017
Year


FISHERY (whole catch)
Mean length
50
45
40
35
2008
2010
2012
2014
2016
Year


FISHERY
Sum of N adj.=348.8
Sum of N eff.=545.8
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
20
40
60
80
Length (cm)


FISHERY
−500
−300
100
300
500
Length (cm)
0
10
20
30
40
50
60
70
80
2009
2012
2014
2017
Year


2007
N adj.=54.8
N eff.=93.2
2017
N adj.=34.4
N eff.=201.2
2009
N adj.=51.2
N eff.=29.6
2012
N adj.=72.2
N eff.=26.9
2013
N adj.=37
N eff.=30.2
2014
N adj.=38.6
N eff.=124.5
2015
N adj.=60.5
N eff.=40.2
Proportion
0.00
0.10
0.20
0.30
0
20
40
60
80
Length (cm)


−500
−300
100
300
500
Length (cm)
0
10
20
30
40
50
60
70
80
2007
2009
2012
2014
2017
Year


Effective sample size
0
50
100
150
200
0
20
40
60
80
Observed sample size
2017
2014
2007
2015
2012
2013
2009
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
50
45
40
35
2008
2010
2012
2014
2016
Year


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.8
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.8
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.4
0.8
1.2
0
1
2
3
4
5
6
Total biomass (mt)


Yield per recruit (kg)
0.00
0.10
0.20
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_95%width_FISHERY(1)
0
5
10
15
20
25
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
20
30
40
50
60
Density
Parameter value