Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Thu Aug 04 13:45:57 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
100
80
60
40
20
0
Length (cm, beginning of the year)
CV of lengths
0.12
0.08
0.04
0
8
4
0
SD of lengths
Age (yr)
CV of lengths
0
0.04
0.08
0.12
0
2
4
6
8
SD of lengths
Length (cm, beginning of the year)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Mean weight (kg) in last year
0
2
4
6
8
10
12
0
20
40
60
80
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
14
0
20
40
60
80
Female length (cm)


Spawning output
0
2
4
6
8
10
12
0
20
40
60
80
Length (cm)


Spawning output
0
1
2
3
4
5
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Length (cm)
0
20
40
60
80
0
5
10
15
20
25
30
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
12
14
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.5
1.0
1.5
2.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.5
0.0
−0.5
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
−1.0
0.0
0.5
1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
0.0
0.4
0.8
1.2
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0
1
2
3
4
5
6
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
2012
2009
2002
2021
2016
2008
1967
Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0
1
2
3
4
5
6
Spawning biomass (mt)


2012
2002
2009
2021
1967
2016
2008
0.5
0.0
−0.5
0.0
0.2
0.4
0.6
0.8
1.0
Log recruitment deviation
Spawning output (relative to B_0)


Landings (mt)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1.4
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
Relative spawning output: B/B_MSY
2000
2020
2010
2021
1980
1967
1970


Index
0
5
10
15
2016
2017
2018
2019
2020
2021
Year


Index
0
5
10
15
2016
2017
2018
2019
2020
2021
Year


Expected index
0
2
4
6
8
10
Observed index
0
2
4
6
8
10


Log index
1.6
1.8
2.0
2.2
2.4
2.6
2.8
2016
2017
2018
2019
2020
2021
Year


Log index
1.6
1.8
2.0
2.2
2.4
2.6
2.8
2016
2017
2018
2019
2020
2021
Year


Log expected index
2.0
2.1
2.2
2.3
2.4
1.9
2.0
2.1
2.2
2.3
Log observed index


Residual
1.5
1.0
0.5
0.0
−0.5
−1.5
2016
2017
2018
2019
2020
2021
Year


Deviation
0.3
0.2
0.1
0.0
−0.1
−0.3
2016
2017
2018
2019
2020
2021
Year


FISHERY
Standardized index
0.0
0.2
0.4
0.6
0.8
1.0
1.2
2015
2016
2017
2018
2019
2020
2021
2022
Year


0.1
2
35
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
Age
Year
0
5
10
15
20
25
30
35
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020


0.1
2
100
80
60
40
20
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


100
300
500
700
Length
0
20
40
60
80
100
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=1455.1
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
85
75
65
55
45
35
25
15
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=69.2
2010
N adj.=29.6
2016
N adj.=163
2005
N adj.=25.2
2011
N adj.=60.5
2017
N adj.=83.2
2006
N adj.=25.2
2012
N adj.=13.6
2018
N adj.=51.2
2007
N adj.=125.2
2013
N adj.=47.4
2019
N adj.=80.6
2008
N adj.=136.2
2014
N adj.=165.1
2020
N adj.=89
2009
N adj.=152
2015
N adj.=125.5
2021
N adj.=13.4
Proportion
0.00
0.10
0.20
0.30
10
30
50
70
90
Length (cm)


0.01
0.2
0.4
Length (cm)
85
75
65
55
45
35
25
15
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
55
50
45
40
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=1455.1
Sum of N eff.=1636.7
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
−15
5
15
Length (cm)
85
75
65
55
45
35
25
15
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=69.2
N eff.=74.7
2010
N adj.=29.6
N eff.=17.4
2016
N adj.=163
N eff.=204.1
2005
N adj.=25.2
N eff.=74.6
2011
N adj.=60.5
N eff.=34.6
2017
N adj.=83.2
N eff.=106.9
2006
N adj.=25.2
N eff.=46.2
2012
N adj.=13.6
N eff.=15.8
2018
N adj.=51.2
N eff.=34.9
2007
N adj.=125.2
N eff.=115.7
2013
N adj.=47.4
N eff.=206.2
2019
N adj.=80.6
N eff.=79.9
2008
N adj.=136.2
N eff.=100
2014
N adj.=165.1
N eff.=233.8
2020
N adj.=89
N eff.=62.1
2009
N adj.=152
N eff.=56.9
2015
N adj.=125.5
N eff.=148.3
2021
N adj.=13.4
N eff.=24.8
Proportion
0.00
0.10
0.20
0.30
10
30
50
70
90
Length (cm)


15
5
15
85
75
65
55
45
35
25
15
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


Effective sample size
0
50
100
200
0
50
100
150
Observed sample size
2013
2014
2016
2015
2007
2008
2017
2019
2020
2009
2005
2004
2006
2018
2011
2021
2012
2010
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
55
50
45
40
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.5
1.0
1.5
2.0
0
5
10
15
20
25
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
Abundance indices
Length compositions
FISHERY
FISHERY
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
Size_inflection_FISHERY(1)
36
38
40
42
44
46
48
50
InitF_seas_1_flt_1FISHERY
0.0
0.1
0.2
0.3
0.4
0.5
Size_95%width_FISHERY(1)
6
8
10
12
14
LnQ_base_FISHERY(1)
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value