Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Wed Jul 27 16:09:07 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
10
20
30
40
0
1
2
3
4
5
6
Age (yr)


Age (yr)
0
1
2
3
4
5
6
Length (cm, beginning of the year)
40
30
20
10
0
CV of lengths
0.08
0.04
0
SD of lengths
2.0
1.0
0.0


Age (yr)
0
1
2
3
4
5
6
Length (cm, beginning of the year)
40
30
20
10
0
Mean weight
0
0.5
1
1.5
Mean weight
1.5
1
0.5
0
Age (yr)
Frac. mature
1.0
0.5
0.0
0.0
0.5
1.0
Frac. mature
Length (cm, beginning of the year)


Length
0
10
20
30
40
0
1
2
3
4
5
6
Age


Length
0
10
20
30
40
0
1
2
3
4
5
6
Age


Mean weight (kg) in last year
0.0
0.5
1.0
1.5
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
0.0
0.5
1.0
1.5
Female weight (kg)


2.0
1.5
1.0
0.5
0.0
Fecundity
0
10
20
30
40
Female length (cm)


Spawning output
0.0
0.5
1.0
1.5
0
10
20
30
40
Length (cm)


Spawning output
0.00
0.05
0.10
0.15
0.20
0.25
0
1
2
3
4
5
6
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
5
6
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
0
1
2
3
4
5
6
Age (yr)
0.2
0.4
0.6


Spawning biomass (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
50
100
150
200
250
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
100
300
500
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
12
14
equilibrium
without fishing
with fishing
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
0.2
0.0
−0.2
−0.4
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
0.5
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0
50
100
150
200
250
0
1
2
3
4
5
6
7
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0
50
100
150
200
250
0
1
2
3
4
5
6
7
Spawning biomass (mt)
1967
1992
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


0.4
0.2
0.0
−0.2
−0.4
Log recruitment deviation
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)
1967
2021


Landings (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
Relative spawning output: B/B_MSY
2010
2020


Index
0
20
40
60
80
100
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
20
40
60
80
100
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0
2
4
6
8
0
5
10
15
20
25
30
Observed index


Log index
−1
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log index
−1
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
Log observed index
1.6
1.7
1.8
1.9
2.0
2.1
2.2
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


50
150
250
Age
0
1
2
3
4
5
6
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


50
150
Age
0
1
2
3
4
5
6
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


50
150
250
Length
0
10
20
30
40
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


20
60
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
50
100
150
200
0
1
2
3
4
5
6
Age


FISHERY
Sum of N adj.=1533.7
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
5
10
15
20
25
30
35
40
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
40
35
30
25
20
15
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=84.3
2010
N adj.=52.6
2016
N adj.=120
2005
N adj.=53.7
2011
N adj.=91.7
2017
N adj.=57.7
2006
N adj.=42.6
2012
N adj.=19.1
2018
N adj.=55.7
2007
N adj.=93.7
2013
N adj.=58.5
2019
N adj.=75.1
2008
N adj.=101.1
2014
N adj.=187.9
2020
N adj.=56.8
2009
N adj.=160.5
2015
N adj.=191.6
2021
N adj.=31.1
Proportion
0.00
0.10
0.20
0.30
5
10
20
30
40
Length (cm)


0.01
0.2
0.4
Length (cm)
40
35
30
25
20
15
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
Mean length
23.0
22.5
22.0
21.5
21.0
20.5
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=1533.7
Sum of N eff.=1775
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
5
10
15
20
25
30
35
40
Length (cm)


FISHERY
−250
−150
50
150
250
40
35
30
25
20
15
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=84.3
N eff.=170.3
2010
N adj.=52.6
N eff.=39.5
2016
N adj.=120
N eff.=76.7
2005
N adj.=53.7
N eff.=74.4
2011
N adj.=91.7
N eff.=160.5
2017
N adj.=57.7
N eff.=34.1
2006
N adj.=42.6
N eff.=79.6
2012
N adj.=19.1
N eff.=36.6
2018
N adj.=55.7
N eff.=50.8
2007
N adj.=93.7
N eff.=168.7
2013
N adj.=58.5
N eff.=153.4
2019
N adj.=75.1
N eff.=110.1
2008
N adj.=101.1
N eff.=91.9
2014
N adj.=187.9
N eff.=138.4
2020
N adj.=56.8
N eff.=27.7
2009
N adj.=160.5
N eff.=88.8
2015
N adj.=191.6
N eff.=254
2021
N adj.=31.1
N eff.=19.5
Proportion
0.00
0.10
0.20
0.30
5
10
20
30
40
Length (cm)


−250
−150
50
150
250
40
35
30
25
20
15
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


Effective sample size
Observed sample size
0
50
150
250
0
50
100
150
200
2015
2014
2013
2004
2007
2011
2019
2008
2009
2016
2006
2005
2018
2012
2017
2020
2021
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
23.0
22.5
22.0
21.5
21.0
20.5
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0
2
4
6
8
10
12
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0
2
4
6
8
10
12
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
−2
0
2
4
6
8
0
5
10
15
20
25
Total biomass (mt)


Yield per recruit (kg)
0.00
0.02
0.04
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
Size_inflection_FISHERY(1)
18
20
22
24
26
28
InitF_seas_1_flt_1FISHERY
0
2
4
6
8
10
Size_95%width_FISHERY(1)
1
2
3
4
5
6
LnQ_base_FISHERY(1)
−10
−5
0
5
10
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value