Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Wed Jul 13 15:12:08 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
60
50
40
30
20
10
0
0
5
10
15
20
25
Age (yr)


Age (yr)
0
5
10
15
20
25
Length (cm, beginning of the year)
60
50
40
30
20
10
0
CV of lengths
0.08
0.04
0
SD of lengths
4
2
0
CV of lengths
0.04
0.08
0
1
2
3
4
SD of lengths
Length (cm, beginning of the year)
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Age (yr)
Frac. mature
Frac. mature
Length (cm, beginning of the year)


Length
60
50
40
30
20
10
0
0
5
10
15
20
25
30
Age


60
50
40
30
20
10
0
Length
0
5
10
15
20
25
30
Age


Mean weight (kg) in last year
3.0
2.5
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
50
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
10
20
30
40
50
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Female weight (kg)


3.5
3.0
2.5
2.0
1.5
1.0
0.5
0.0
Fecundity
0
10
20
30
40
50
Female length (cm)


Spawning output
3.0
2.5
2.0
1.5
1.0
0.5
0.0
0
10
20
30
40
50
Length (cm)


Spawning output
1.2
1.0
0.8
0.6
0.4
0.2
0.0
0
5
10
15
20
25
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Length (cm)
0
10
20
30
40
50
0
5
10
15
20
25
Age (yr)
0.8
0.6
0.4
0.2


Length (cm)
0
10
20
30
40
50
0
5
10
15
20
25
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
20
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
1
2
3
4
5
6
equilibrium
without fishing
with fishing
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
−1.0
−0.5
0.0
0.5
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
2
1
0
−1
−2
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
−0.5
1970
1980
1990
2000
2010
2020
Year
$1-SE(\hat{r}_y)^2/\sigma_R^2$


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
1
2
3
4
5
6
7
0.0
0.5
1.0
1.5
2.0
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
2009
1997
2000
2012
2010
1967
2021
2005
2006
2008
2007
Recruitment (1,000s)
0
1
2
3
4
5
6
7
0.0
0.5
1.0
1.5
2.0
Spawning biomass (mt)


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
2009
2000
1967
2005
2006
2008
2007
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)


Landings (mt)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.00
0.10
0.20
0.30
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Relative spawning output: B/B_MSY
1967
2021


Index
0
5
10
15
20
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Observed index


Log index
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.2
0.0
−0.2
−0.4
−0.6
−2
−1
0
1
Log observed index


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
Year


Index
0
5
10
15
20
25
30
35
1990
1995
2000
2005
Year


Expected index
0.0
0.5
1.0
1.5
2.0
2.5
0
1
2
3
4
5
Observed index


Log index
2
0
−2
−4
−6
−8
1990
1995
2000
2005
Year


Log index
2
0
−2
−4
−6
−8
1990
1995
2000
2005
Year


Log expected index
1.0
0.8
0.6
0.4
0.2
−4
−2
0
Log observed index


Residual
4
2
0
−2
−4
1990
1995
2000
2005
Year


Deviation
6
4
2
0
−2
−4
−6
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
1990
1995
2000
2005
2010
2015
2020
Year


0.1
4
8
Age
Year
0
5
10
15
20
25
30
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020


0.1
4
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
4
8
60
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
Length
Year
0
10
20
30
40
50
60
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
5
10
15
20
25
Age


FISHERY
Sum of N adj.=2610.7
Proportion
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
20
25
30
35
40
45
50
55
Length (cm)


FISHERY
0.01
0.2
0.4
55
50
45
40
35
30
25
20
Length (cm)
2013
2015
Year


2012
N adj.=1018.1
2013
N adj.=1078.6
2015
N adj.=514.1
0.0
0.1
0.2
0.3
0.4
20
30
40
50
Proportion
Length (cm)


0.01
0.2
0.4
55
50
45
40
35
30
25
20
Length (cm)
2012
2013
2015
Year


FISHERY (whole catch)
Mean length
34.0
33.5
33.0
32.5
32.0
2012.0
2012.5
2013.0
2013.5
2014.0
2014.5
2015.0
Year


FISHERY
Sum of N adj.=2610.7
Sum of N eff.=1006.1
Proportion
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
20
25
30
35
40
45
50
55
Length (cm)


FISHERY
−70
−50
−30
10
30
50
70
Length (cm)
55
50
45
40
35
30
25
20
2013
2015
Year


2012
N adj.=1018.1
N eff.=425
2013
N adj.=1078.6
N eff.=226.3
2015
N adj.=514.1
N eff.=354.8
0.0
0.1
0.2
0.3
0.4
20
30
40
50
Proportion
Length (cm)


−70
−50
−30
10
30
50
70
55
50
45
40
35
30
25
20
Length (cm)
2012
2013
2015
Year


Effective sample size
0
100
200
300
400
0
200
400
600
800
1000
1200
Observed sample size
harmonic mean
arithmetic mean
2012
2015
2013


FISHERY (whole catch)
Mean length
34.0
33.5
33.0
32.5
32.0
2012.0
2012.5
2013.0
2013.5
2014.0
2014.5
2015.0
Year


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.2
0.4
0.6
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
1.0
0.5
0.0
−0.5
0
2
4
6
8
10
Total biomass (mt)


Yield per recruit (kg)
0.00
0.10
0.20
0.30
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
Size_inflection_FISHERY(1)
15
20
25
30
35
40
45
50
InitF_seas_1_flt_1FISHERY
0.0
0.5
1.0
1.5
2.0
Size_95%width_FISHERY(1)
0
2
4
6
8
10
12
14
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
LnQ_base_SURVEY2(2)
−6
−4
−2
0
2
4
Density
Parameter value