American Samoa Model Checks

Meg Oshima

2022-08-10

This is a summary report for the LERU base model run.

Model Output

Input Data





Convergence Check

```
##   Converged     MaxGrad
## 1      TRUE 2.26599e-05
```

```
## [1] "1 NOTE:  Max data length bin: 38.5  < max pop len bins: 43; so will accumulate larger pop len bi
## [2] "2 parameter init value is greater than parameter max 0.8 > 0.6 for parm: 14 ; search for <now cl
## [3] "3  Main recdev biasadj is >2 times ratio of rmse to sigmaR"
## [4] " N parameters are on or within 1% of min-max bound: 1; check results, variance may be suspect"
## [5] "N warnings: 3"
```

Fit to Model

CPUE

```
##
##  Running Runs Test Diagnosics for Index
## Plotting Residual Runs Tests
```




```
##
## Runs Test stats by Index:
## Plotting JABBA residual plot
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : Chernobyl! trL>n 6
```

```
## Warning in simpleLoess(y, x, w, span, degree = degree, parametric = parametric, : Chernobyl! trL>n 6

## Warning in sqrt(sum.squares/one.delta): NaNs produced
```




```
##
## RMSE stats by Index:
```

Length Comp

#Factor	Fleet	New_Var_adj	Type	Name
4	1	0.338524	len	FISHERY

```
##
##  Running Runs Test Diagnosics for Mean length
## Plotting Residual Runs Tests

##
## Runs Test stats by Mean length:

##     Index runs.p   test   sigma3.lo  sigma3.hi type
## 1 FISHERY  0.314 Passed -0.04139817 0.04139817  len
```

```
## Plotting JABBA residual plot
```




```
##
## RMSE stats by Index:

##    indices RMSE.perc nobs
## 1  FISHERY       1.7   18
## 2 Combined       1.7   18
```


FISHERY (whole catch)
Mean length
29
28
27
26
25
24
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=1942.8
Sum of N eff.=1745.3
Proportion
0.0
0.1
0.2
0.3
0.4
15
20
25
30
35
40
Length (cm)



Retrospective and Hindcasting

Retrospective

```
## Plotting Retrospective pattern
```

```
##
## Mohn's Rho stats, including one step ahead forecasts:
## Plotting Retrospective pattern
```




```
##
## Mohn's Rho stats, including one step ahead forecasts:

##   type     peel        Rho ForecastRho
## 1    F     2020 -0.1887474 -0.14190907
## 2    F     2019 -0.2750072 -0.21281215
## 3    F     2018  0.1193334  0.14341093
## 4    F Combined -0.1148070 -0.07043676
```

Hindcasting

```
## Plotting Hindcast Cross-Validation (one-step-ahead)
##
##  Computing MASE with only 2 of 3  prediction residuals for Index FISHERY
##
##  Warning:  Unequal spacing of naive predictions residuals may influence the interpretation of MASE
```




```
##
## MASE stats by Index:
## Plotting Hindcast Cross-Validation (one-step-ahead)
##
##  Computing MASE with only 2 of 3  prediction residuals for Index FISHERY
##
##  Warning:  Unequal spacing of naive predictions residuals may influence the interpretation of MASE

##
## MASE stats by Index:
```


FISHERY: MASE = 1.31 (0.2
Ref
2020
2019
2018
Mean length
0
5
10
15
20
25
30
2013
2015
2017
2019
Year

Recruitment Deviations





Likelihood Profile

```
## [1] "SR_LN"
```

```
## Parameter matching profile.string=SR_LN: SR_LN(R0)
```

```
## Parameter values (after subsetting based on input 'models'): 3, 3.2, 3.4, 3.6, 3.8, 4, 4.2, 4.4, 4.6
```

```
##
## Likelihood components showing max change as fraction of total change.
## To change which components are included, change input 'minfraction'.
```

```
##                       frac_change include                            label
## TOTAL                      1.0000    TRUE                            Total
## Catch                      0.3073    TRUE                            Catch
## Equil_catch                0.0000   FALSE                Equilibrium catch
## Survey                     0.1911    TRUE                       Index data
## Length_comp                0.0943    TRUE                      Length data
## Recruitment                0.5252    TRUE                      Recruitment
## InitEQ_Regime              0.0000   FALSE Initital equilibrium recruitment
## Forecast_Recruitment       0.0000   FALSE             Forecast recruitment
## Parm_priors                0.0000   FALSE                           Priors
## Parm_softbounds            0.0000   FALSE                      Soft bounds
## Parm_devs                  0.0000   FALSE             Parameter deviations
## Crash_Pen                  0.0000   FALSE                    Crash penalty
```

```
## Parameter matching profile.string = 'SR_LN': 'SR_LN(R0)
## Parameter values (after subsetting based on input 'models'): 3, 3.2, 3.4, 3.6, 3.8, 4, 4.2, 4.4, 4.6
```

```
## Fleet-specific likelihoods showing max change as fraction of total change.
## To change which components are included, change input 'minfraction'.
##                         frac_change include
## prof.table....c.1.3..           1    TRUE
```


Change in −log−likelihood
0
5
10
15
20
25
3.0
3.5
4.0
4.5
SR_LN(R0)
Total
Recruitment
Catch
Index data
Length data



Management Quantities

```
##
##  starter.sso with Bratio: SSB/SSBMSY and F: _abs_F
##
```




```
##
## Plot Comparison of stock
```




```
##
## Plot Comparison of harvest
```




```
##
## Plot Comparison of SSB
```




```
##
## Plot Comparison of F
```


MVLN
Fishing Mortality (F)
0
2
4
6
1970
1980
1990
2000
2010
2020
Year

```
## RStudioGD
##         2
```

Jitter




model 1
model 2
model 3
Spawning biomass (t)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


model 1
model 2
model 3
abs F; with F=sum(full Fs)
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


model 1
model 2
model 3
Age−0 recruits (1,000s)
0
20
40
60
80
1970
1980
1990
2000
2010
2020
Year