Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Fri Jul 22 14:21:31 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
10
20
30
40
0
1
2
3
4
5
6
Age (yr)


Age (yr)
0
1
2
3
4
5
6
Length (cm, beginning of the year)
0
10
20
30
40
CV of lengths
0
0.04
0.08
0.12
SD of lengths
0.0
1.0
2.0
3.0
Length (cm, beginning of the year)
CV of lengths
SD of lengths
Age (yr)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
0
10
20
30
40
0
1
2
3
4
5
6
Age


Length
0
10
20
30
40
0
1
2
3
4
5
6
Age


Mean weight (kg) in last year
0.0
0.5
1.0
1.5
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.5
1.0
1.5
Female weight (kg)


Fecundity
0.0
0.5
1.0
1.5
0.0
0.5
1.0
1.5
Female weight (kg)


2.0
1.5
1.0
0.5
0.0
Fecundity
0
10
20
30
40
Female length (cm)


Spawning output
0.0
0.5
1.0
1.5
0
10
20
30
40
Length (cm)


0.25
0.20
0.15
0.10
0.05
0.00
Spawning output
0
1
2
3
4
5
6
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
5
6
Age (yr)
FISHERY
SURVEY2


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
0
1
2
3
4
5
6
Age (yr)
0.2
0.4


Length (cm)
0
10
20
30
40
0
1
2
3
4
5
6
Age (yr)
0.2
0.4


Spawning biomass (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
10
20
30
40
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
500
1000
1500
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
500
1500
2500
3500
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
1.0
0.5
−0.5
−1.5
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
2
1
0
−1
−2
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
−0.5
$1-SE(\hat{r}_y)^2/\sigma_R{}^2$
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
500
1000
1500
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
Spawning biomass (mt)
0
500
1000
1500
0.0
0.5
1.0
1.5
2.0
2.5
3.0
2019
2018
1985
2016
2017
2005
1986
1967
2021
2001
1999
2000
2009
2002
2010
2011


Log recruitment deviation
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
0.0
0.5
1.0
1.5
2.0
Spawning output (relative to B_0)
2019
2018
2016
2005
1985
2017
1986
1967
2021
2011
2010
2001
1999
2015
2002
2009
2000


Landings (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
2
4
6
8
10
12
1970
1980
1990
2000
2010
2020
Year


Continuous F
0
1
2
3
4
5
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
Relative spawning output: B/B_MSY
2000
1990
1967
1970
2020
2021


Index
0
10
20
30
40
50
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
Observed index
0
5
10
15
20
25
30


Log index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log index
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
0.5
1.0
1.5
2.0
2.5
3.0
3.5
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
Log observed index


Effective catchability
5.521020
5.521010
5.521000
1
2
3
4
5
Vulnerable biomass


Residual
15
10
5
0
−5
−10
−15
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
Year


Index
0
10
20
30
40
50
1990
1995
2000
2005
Year


Expected index
0
2
4
6
8
10
0
5
10
15
20
25
Observed index


Log index
4
2
0
−2
−4
1990
1995
2000
2005
Year


Log index
4
2
0
−2
−4
1990
1995
2000
2005
Year


Log expected index
0.0
0.5
1.0
1.5
2.0
−1
0
1
2
3
Log observed index


Effective catchability
3.796238
3.796234
3.796230
0.5
1.0
1.5
2.0
2.5
Vulnerable biomass


Residual
4
2
0
−2
−4
1990
1995
2000
2005
Year


Deviation
2
1
0
−1
−2
1990
1995
2000
2005
Year


FISHERY
SURVEY2
Standardized index
Year
0
1
2
3
4
1990
1995
2000
2005
2010
2015
2020


0.1
2
Age
0
1
2
3
4
5
6
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
2
Age
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


200
600
1000
Length
0
10
20
30
40
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


100
300
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
100
200
300
400
500
0
1
2
3
4
5
6
Age


FISHERY
Sum of N adj.=45108
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
5
10
15
20
25
30
35
40
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
40
35
30
25
20
15
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=2478
2010
N adj.=1545.6
2016
N adj.=3528
2005
N adj.=1579.2
2011
N adj.=2696.4
2017
N adj.=1696.8
2006
N adj.=1251.6
2012
N adj.=562.8
2018
N adj.=1638
2007
N adj.=2755.2
2013
N adj.=1722
2019
N adj.=2209.2
2008
N adj.=2973.6
2014
N adj.=5527.2
2020
N adj.=1671.6
2009
N adj.=4720.8
2015
N adj.=5636.4
2021
N adj.=915.6
Proportion
0.00
0.10
0.20
0.30
5
10
20
30
40
Length (cm)


0.01
0.2
0.4
Length (cm)
40
35
30
25
20
15
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
23.0
22.5
22.0
21.5
21.0
20.5
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=45108
Sum of N eff.=2086.4
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
5
10
15
20
25
30
35
40
Length (cm)


FISHERY
−1400
−1000
−600
200
600
1000
1400
40
35
30
25
20
15
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=2478
N eff.=229.5
2010
N adj.=1545.6
N eff.=41.1
2016
N adj.=3528
N eff.=98.7
2005
N adj.=1579.2
N eff.=89.6
2011
N adj.=2696.4
N eff.=235.9
2017
N adj.=1696.8
N eff.=45
2006
N adj.=1251.6
N eff.=73.3
2012
N adj.=562.8
N eff.=32.9
2018
N adj.=1638
N eff.=38.7
2007
N adj.=2755.2
N eff.=160.5
2013
N adj.=1722
N eff.=167.7
2019
N adj.=2209.2
N eff.=98
2008
N adj.=2973.6
N eff.=122.1
2014
N adj.=5527.2
N eff.=186.9
2020
N adj.=1671.6
N eff.=29.9
2009
N adj.=4720.8
N eff.=103.7
2015
N adj.=5636.4
N eff.=318.7
2021
N adj.=915.6
N eff.=14.1
Proportion
0.00
0.10
0.20
0.30
5
10
20
30
40
Length (cm)


−1400
−1000
−600
200
600
1000
1400
40
35
30
25
20
15
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


Effective sample size
Observed sample size
0
50
150
250
350
0
1000
2000
3000
4000
5000
6000
harmonic mean
arithmetic mean
2015
2011
2004
2014
2013
2007
2008
2005
2019
2016
2009
2006
2012
2021
2020


FISHERY (whole catch)
23.0
22.5
22.0
21.5
21.0
20.5
Mean length
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
10
5
0
−5
0
10
20
30
40
Total biomass (mt)


Yield per recruit (kg)
0.000
0.004
0.008
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
SURVEY2
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
5
6
7
8
Size_inflection_FISHERY(1)
18
20
22
24
26
28
InitF_seas_1_flt_1FISHERY
0
2
4
6
8
10
Size_95%width_FISHERY(1)
1.0
1.5
2.0
2.5
3.0
LnQ_base_FISHERY(1)
−1
0
1
2
3
4
LnQ_base_SURVEY2(2)
−1
0
1
2
3
4
Density
Parameter value