Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Thu Jul 14 16:17:56 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
0
10
20
30
40
0
2
4
6
8
Age (yr)


Age (yr)
0
2
4
6
8
Length (cm, beginning of the year)
0
10
20
30
40
CV of lengths
0
0.02
0.06
SD of lengths
0.0
1.0
2.0
Length (cm, beginning of the year)
CV of lengths
0
0.04
0.08
SD of lengths
0.0
1.0
2.0
Age (yr)


Age (yr)
0
2
4
6
8
Length (cm, beginning of the year)
0
10
20
30
40
Mean weight
0
0.4
0.8
1.2
Frac. mature
0.0
0.5
1.0
Age (yr)


Length
0
10
20
30
40
0
2
4
6
8
Age


Length
0
10
20
30
40
0
2
4
6
8
Age


Mean weight (kg) in last year
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0
10
20
30
40
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
10
20
30
40
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Female weight (kg)


Fecundity
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0.0
0.2
0.4
0.6
0.8
1.0
1.2
Female weight (kg)


Fecundity
1.4
1.2
1.0
0.8
0.6
0.4
0.2
0.0
0
10
20
30
40
Female length (cm)


Spawning output
1.2
1.0
0.8
0.6
0.4
0.2
0.0
0
10
20
30
40
Length (cm)


Spawning output
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0
2
4
6
8
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
2
4
6
8
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
Length (cm)


Length (cm)
0
10
20
30
40
0
2
4
6
8
Age (yr)
0.8
0.6
0.4


Spawning biomass (mt)
0
2
4
6
8
10
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0
2
4
6
8
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
20
40
60
80
120
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
50
150
250
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0
1
2
3
4
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
2
1
0
−1
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
−0.5
$1-SE(\hat{r}_y)^2/\sigma_R^2$
1970
1980
1990
2000
2010
2020
Year


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
0
20
40
60
80
100
140
0
1
2
3
4
5
Spawning biomass (mt)


Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
Spawning biomass (mt)
0
20
40
60
80
100
140
0
1
2
3
4
5
1988
2006
2007
1995
2008
2000
2021
2015
2003


1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
Log recruitment deviation
0.0
0.5
1.0
1.5
Spawning output (relative to B_0)
1987
2006
1988
2007
1995
2008
2000
2021
2018
2019
2017
2016
2003
2015


Landings (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0
1
2
3
4
5
6
7
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1.4
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
5
6
7
Relative spawning output: B/B_MSY
2021
1967


Index
0
10
20
30
40
50
60
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
10
20
30
40
50
60
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
Observed index
0
5
10
15
20
25
30
0
10
20
30
40


Log index
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log index
1
2
3
4
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
1.5
2.0
2.5
3.0
1.0
1.5
2.0
2.5
3.0
3.5
Log observed index


Residual
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
0.5
0.0
−0.5
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
0.0
0.5
1.0
1.5
2.0
2.5
1990
1995
2000
2005
2010
2015
2020
Year


20
60
100
140
Age
0
1
2
3
4
5
6
7
8
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


20
60
100
Age
8
7
6
5
4
3
2
1
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


10
30
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


5
15
25
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
10
20
30
0
2
4
6
8
Age


FISHERY
Sum of N adj.=974.1
Proportion
0.00
0.05
0.10
0.15
0.20
0.25
10
15
20
25
30
35
40
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
38
34
30
26
22
18
14
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=22.3
2010
N adj.=15.6
2016
N adj.=37.9
2005
N adj.=18.8
2011
N adj.=117.3
2017
N adj.=16.6
2006
N adj.=15.8
2012
N adj.=200
2018
N adj.=21.5
2007
N adj.=12.3
2013
N adj.=234.7
2019
N adj.=28.7
2008
N adj.=37.5
2014
N adj.=28.6
2020
N adj.=23.1
2009
N adj.=49.9
2015
N adj.=81.4
2021
N adj.=12
Proportion
0.00
0.10
0.20
0.30
10
20
30
40
Length (cm)


0.01
0.2
0.4
Length (cm)
38
34
30
26
22
18
14
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


FISHERY (whole catch)
32
30
28
26
24
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=974.1
Sum of N eff.=1213.7
Proportion
0.30
0.25
0.20
0.15
0.10
0.05
0.00
10
15
20
25
30
35
40
Length (cm)


FISHERY
−350
−250
−150
50
150
250
350
Length (cm)
38
34
30
26
22
18
14
10
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


2004
N adj.=22.3
N eff.=33.8
2010
N adj.=15.6
N eff.=23.1
2016
N adj.=37.9
N eff.=48.2
2005
N adj.=18.8
N eff.=34.8
2011
N adj.=117.3
N eff.=157.9
2017
N adj.=16.6
N eff.=37.8
2006
N adj.=15.8
N eff.=24.7
2012
N adj.=200
N eff.=137
2018
N adj.=21.5
N eff.=47.3
2007
N adj.=12.3
N eff.=20.7
2013
N adj.=234.7
N eff.=156.5
2019
N adj.=28.7
N eff.=32.6
2008
N adj.=37.5
N eff.=25.3
2014
N adj.=28.6
N eff.=127.1
2020
N adj.=23.1
N eff.=41.8
2009
N adj.=49.9
N eff.=98.1
2015
N adj.=81.4
N eff.=76.7
2021
N adj.=12
N eff.=90
Proportion
0.00
0.10
0.20
0.30
10
20
30
40
Length (cm)


−350
−250
−150
50
150
250
350
38
34
30
26
22
18
14
10
Length (cm)
2004
2006
2008
2010
2012
2014
2016
2018
2020
Year


Effective sample size
Observed sample size
0
50
100
150
200
250
2011
2013
2012
2014
2009
2021
2015
harmonic mean
arithmetic mean
2016
2008


FISHERY (whole catch)
Mean length
32
30
28
26
24
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
1.0
2.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
1.0
2.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
10
5
0
0
5
10
15
20
25
Total biomass (mt)


Yield per recruit (kg)
0.00
0.04
0.08
0.12
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1965
1975
1985
1995
2005
2015
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
5
Size_95%width_FISHERY(1)
0
1
2
3
4
5
InitF_seas_1_flt_1FISHERY
0.0
0.5
1.0
1.5
2.0
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
15
20
25
30
35
40
Density
Parameter value