Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Wed Aug 03 09:28:16 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
100
80
60
40
20
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
100
80
60
40
20
0
Length (cm, beginning of the year)
CV of lengths
0.08
0.04
0
SD of lengths
6
4
2
0
Age (yr)
CV of lengths
0
0.02
0.06
0
2
3
4
5
6
SD of lengths
Length (cm, beginning of the year)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)
Frac. mature
Length (cm, beginning of the year)


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Length
100
80
60
40
20
0
0
5
10
15
20
25
30
35
Age


Mean weight (kg) in last year
0
2
4
6
8
10
12
0
20
40
60
80
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Maturity
0
20
40
60
80
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
0
2
4
6
8
10
12
Female weight (kg)


Fecundity
0
2
4
6
8
10
12
14
0
20
40
60
80
Female length (cm)


Spawning output
0
2
4
6
8
10
12
0
20
40
60
80
Length (cm)


Spawning output
0
1
2
3
4
5
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
20
40
60
80
Length (cm)


Length (cm)
0
20
40
60
80
0
5
10
15
20
25
30
Age (yr)
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0
5
10
15
1990
2000
2010
2020
Year


Spawning biomass (mt)
0
5
10
15
20
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.2
0.4
0.6
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.2
0.4
0.6
0.8
1.0
1990
2000
2010
2020
Year


Total biomass (mt)
0
5
10
15
20
25
30
35
1990
2000
2010
2020
Year


Summary biomass (mt)
0
5
10
15
20
25
30
35
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Spawning biomass (mt)
0
5
10
15
1985
1990
1995
2000
2005
2010
2015
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.6
0.4
0.2
0.0
−0.4
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
1.0
0.5
0.0
−0.5
−1.0
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.1
0.2
0.3
0.4
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
$1-SE(\hat{r}_y)^2/\sigma_R^2$
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0.0
0.5
1.0
1.5
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Recruitment (1,000s)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Spawning biomass (mt)
0.0
0.5
1.0
1.5
2011
2004
2007
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


Log recruitment deviation
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
0.00
0.05
0.10
0.15
Spawning output (relative to B_0)
2011
2004
2021
1986
2007


Landings (mt)
0
1
2
3
4
5
6
1985
1990
1995
2000
2005
2010
2015
2020
Year


Landings (mt)
0
1
2
3
4
5
6
1985
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0
1
2
3
4
5
6
1985
1990
1995
2000
2005
2010
2015
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1985
1990
1995
2000
2005
2010
2015
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
Relative spawning output: B/B_MSY
1986
1990
2000
2010
2020
2021


Index
0
5
10
15
2016
2017
2018
2019
2020
2021
Year


Index
0
5
10
15
2016
2017
2018
2019
2020
2021
Year


Expected index
Observed index
0
2
4
6
8
10


Log index
1.6
1.8
2.0
2.2
2.4
2.6
2.8
2016
2017
2018
2019
2020
2021
Year


Log index
1.6
1.8
2.0
2.2
2.4
2.6
2.8
2016
2017
2018
2019
2020
2021
Year


Log expected index
2.00
2.05
2.10
2.15
2.20
2.25
2.30
1.9
2.0
2.1
2.2
2.3
Log observed index


Residual
0.6
0.4
0.2
−0.2
−0.6
2016
2017
2018
2019
2020
2021
Year


Deviation
0.10
0.05
0.00
−0.10
2016
2017
2018
2019
2020
2021
Year


FISHERY
Standardized index
0.0
0.2
0.4
0.6
0.8
1.0
1.2
2015
2016
2017
2018
2019
2020
2021
2022
Year


0.1
2
Age
Year
0
5
10
15
20
25
30
35
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021


0.1
2
35
30
25
20
15
10
5
0
Age
1985
1989
1993
1997
2001
2005
2009
2013
2017
2021
Year


0.1
2
100
80
60
40
20
0
Length
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


0.1
2
100
80
60
40
20
0
Length
1986
1990
1994
1998
2002
2006
2010
2014
2018
2022
Year


Numbers at age at equilibrium
0.0
0.5
1.0
1.5
2.0
2.5
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=1348.1
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
0.01
0.2
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2013
2015
2017
2019
Year


2004
N adj.=69.2
2014
N adj.=165.1
2020
N adj.=89
2007
N adj.=125.2
2015
N adj.=125.5
2008
N adj.=136.2
2016
N adj.=163
2009
N adj.=152
2017
N adj.=83.2
2011
N adj.=60.5
2018
N adj.=51.2
2013
N adj.=47.4
2019
N adj.=80.6
0.30
0.20
0.10
0.00
10
30
50
70
90
Proportion
Length (cm)


0.01
0.2
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2013
2015
2017
2019
Year


FISHERY (whole catch)
55
50
45
Mean length
2005
2010
2015
2020
Year


FISHERY
Sum of N adj.=1348.1
Sum of N eff.=1150.4
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
10
20
30
40
50
60
70
80
90
Length (cm)


FISHERY
15
5
15
Length (cm)
85
75
65
55
45
35
25
15
2004
2007
2009
2011
2013
2015
2017
2019
Year


2004
N adj.=69.2
N eff.=53.3
2014
N adj.=165.1
N eff.=90.9
2020
N adj.=89
N eff.=69.7
2007
N adj.=125.2
N eff.=150.9
2015
N adj.=125.5
N eff.=143.9
2008
N adj.=136.2
N eff.=80.6
2016
N adj.=163
N eff.=138.8
2009
N adj.=152
N eff.=68.2
2017
N adj.=83.2
N eff.=98.7
2011
N adj.=60.5
N eff.=59.8
2018
N adj.=51.2
N eff.=36.1
2013
N adj.=47.4
N eff.=88.4
2019
N adj.=80.6
N eff.=71.2
Proportion
0.00
0.10
0.20
0.30
10
30
50
70
90
Length (cm)


15
5
15
85
75
65
55
45
35
25
15
Length (cm)
2004
2007
2009
2011
2013
2015
2017
2019
Year


Effective sample size
0
50
100
150
Observed sample size
0
50
100
150
harmonic mean
arithmetic mean
2007
2015
2016
2017
2013
2014
2008
2019
2020
2009
2011
2004
2018


FISHERY (whole catch)
55
50
45
Mean length
2005
2010
2015
2020
Year


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
2.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Equilibrium yield (mt)
0.0
0.5
1.0
1.5
2.0
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished
MSY
B target
SPR target
Current


Surplus production (mt)
0.0
0.5
1.0
1.5
2.0
0
2
4
6
8
Total biomass (mt)


Yield per recruit (kg)
0.0
0.2
0.4
0.6
0.8
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


Catches
Abundance indices
Length compositions
FISHERY
FISHERY
FISHERY
1985
1990
1995
2000
2005
2010
2015
2020
Year


SR_LN(R0)
prior
max. likelihood
1
2
3
4
Size_inflection_FISHERY(1)
36
38
40
42
44
46
48
50
InitF_seas_1_flt_1FISHERY
0.0
0.1
0.2
0.3
0.4
0.5
Size_95%width_FISHERY(1)
6
8
10
12
14
LnQ_base_FISHERY(1)
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
Q_extraSD_FISHERY(1)
0.0
0.5
1.0
1.5
2.0
Density
Parameter value