Plots created using the 'r4ss' package in R
Stock Synthesis version: 3.30.19.0
StartTime: Mon Jun 20 15:23:07 2022
Data_File: data.ss
Control_File: control.ss


Length (cm, beginning of the year)
60
50
40
30
20
10
0
0
5
10
15
20
25
30
Age (yr)


Age (yr)
0
5
10
15
20
25
30
Length (cm, beginning of the year)
60
50
40
30
20
10
0
CV of lengths
0.08
0.04
0
SD of lengths
4
2
0
CV of lengths
0.04
0.08
0
1
2
3
4
SD of lengths
Length (cm, beginning of the year)


Age (yr)
Mean weight
Length (cm, beginning of the year)
Mean weight
Frac. mature
Age (yr)


Length
60
50
40
30
20
10
0
0
5
10
15
20
25
30
Age


60
50
40
30
20
10
0
Length
0
5
10
15
20
25
30
Age


Mean weight (kg) in last year
0
1
2
3
0
10
20
30
40
50
Length (cm)


Maturity
0.0
0.2
0.4
0.6
0.8
0
10
20
30
40
50
Length (cm)


Egg output proportional to spawning biomass
Eggs per kg
1.0
0.8
0.6
0.4
0.2
0.0
0
1
2
3
Female weight (kg)


Fecundity
0
1
2
3
0
1
2
3
Female weight (kg)


Fecundity
0
1
2
3
4
0
10
20
30
40
50
Female length (cm)


Spawning output
0
1
2
3
0
10
20
30
40
50
Length (cm)


1.0
0.8
0.6
0.4
0.2
0.0
Spawning output
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
5
10
15
20
25
30
Age (yr)


Selectivity
0.0
0.2
0.4
0.6
0.8
1.0
0
10
20
30
40
50
Length (cm)


Length (cm)
0
10
20
30
40
50
0
5
10
15
20
25
30
Age (yr)
1
0.8
0.6
0.4
0.2


Spawning biomass (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.4
0.8
1.2
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Relative spawning biomass: B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Total biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Summary biomass (mt)
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
1
2
3
4
5
6
1970
1980
1990
2000
2010
2020
Year


Age−0 recruits (1,000s)
0
5
10
15
20
25
1970
1980
1990
2000
2010
2020
Year


Summary Fishing Mortality
0.0
1.0
2.0
3.0
1970
1980
1990
2000
2010
2020
Year


Spawning biomass (mt)
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year
equilibrium
without fishing
with fishing


Log recruitment deviation
0.8
0.6
0.4
0.2
0.0
−0.4
1970
1980
1990
2000
2010
2020
Year


Log recruitment deviation
2
1
0
−1
1970
1980
1990
2000
2010
2020
Year


Recruitment deviation variance
Asymptotic standard error estimate
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


bias adjust in model
estimated alternative
1.0
0.5
0.0
−0.5
$1-SE(\hat{r}_y)^2/\sigma_R^2$
1970
1980
1990
2000
2010
2020
Year


Recruitment (1,000s)
0
1
2
3
4
5
6
0.0
0.1
0.2
0.3
0.4
0.5
Spawning biomass (mt)
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium


1997
1998
2008
2021
1967
Exp. recruitment
Exp. recruitment after bias adj.
Estimated recruitments
Unfished equilibrium
Recruitment (1,000s)
Spawning biomass (mt)
0
1
2
3
4
5
6
0.0
0.1
0.2
0.3
0.4
0.5


1998
1997
2008
0.5
0.0
−0.5
Log recruitment deviation
0.0
0.2
0.4
0.6
0.8
1.0
Spawning output (relative to B_0)


Landings (mt)
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


Landings (mt)
0.0
0.2
0.4
0.6
1970
1980
1990
2000
2010
2020
Year


FISHERY
FISHERY obs.
Observed and expected Landings (mt)
0.0
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Year


Continuous F
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
Year


SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.5
1.0
1.5
1970
1980
1990
2000
2010
2020
Year


Fishing intensity: 1−SPR
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Relative spawning output: B/B_MSY
1969
2021


Index
0
20
40
60
80
100
140
1990
1995
2000
2005
2010
2015
2020
Year


Index
0
20
40
60
80
100
140
1990
1995
2000
2005
2010
2015
2020
Year


Expected index
0.0
0.1
0.2
0.3
0.4
0
1
2
3
4
Observed index


Log index
4
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log index
4
2
0
−2
−4
−6
1990
1995
2000
2005
2010
2015
2020
Year


Log expected index
−1.0
−2.0
−3.0
−4.0
−6
−5
−4
−3
−2
−1
0
1
Log observed index


Effective catchability
0.292735
0.292733
0.292731
0.2
0.4
0.6
0.8
1.0
1.2
Vulnerable biomass


Residual
2
1
0
−1
−2
1990
1995
2000
2005
2010
2015
2020
Year


Deviation
3
2
1
0
−1
−2
−3
1990
1995
2000
2005
2010
2015
2020
Year


FISHERY
Standardized index
Year
0
2
4
6
8
10
1990
1995
2000
2005
2010
2015
2020


0.1
4
8
Age
30
25
20
15
10
5
0
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
4
30
25
20
15
10
5
0
Age
1966
1972
1978
1984
1990
1996
2002
2008
2014
2020
Year


0.1
4
8
60
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


0.1
2
60
50
40
30
20
10
0
Length
1967
1973
1979
1985
1991
1997
2003
2009
2015
2021
Year


Numbers at age at equilibrium
0
1
2
3
4
0
5
10
15
20
25
30
Age


FISHERY
Sum of N adj.=220.4
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
10
20
30
40
Length (cm)


FISHERY
0.01
0.2
0.4
Length (cm)
42
36
30
24
18
12
6
0
2012
2015
2017
Year


2007
N adj.=50.7
2012
N adj.=42.8
2015
N adj.=33.6
2016
N adj.=33.6
2017
N adj.=59.6
Proportion
0.00
0.10
0.20
0.30
0
10
20
30
40
Length (cm)


0.01
0.2
0.4
Length (cm)
0
6
12
18
24
30
36
42
2007
2012
2015
2017
Year


FISHERY (whole catch)
32
30
28
26
24
Mean length
2008
2010
2012
2014
2016
Year


FISHERY
Sum of N adj.=220.4
Sum of N eff.=315.1
Proportion
0.25
0.20
0.15
0.10
0.05
0.00
0
10
20
30
40
Length (cm)


FISHERY
−35
−25
−15
5
15
25
35
Length (cm)
0
6
12
18
24
30
36
42
2012
2015
2017
Year


2007
N adj.=50.7
N eff.=89.9
2012
N adj.=42.8
N eff.=69.3
2015
N adj.=33.6
N eff.=33.1
2016
N adj.=33.6
N eff.=20.7
2017
N adj.=59.6
N eff.=102.1
0.00
0.10
0.20
0.30
0
10
20
30
40
Proportion
Length (cm)


−35
−25
−15
5
15
25
35
Length (cm)
0
6
12
18
24
30
36
42
2007
2012
2015
2017
Year


Effective sample size
0
20
40
60
80
0
10
20
30
40
50
60
Observed sample size
2017
2007
2012
2015
2016
harmonic mean
arithmetic mean


FISHERY (whole catch)
Mean length
32
30
28
26
24
2008
2010
2012
2014
2016
Year


Equilibrium yield (mt)
0.00
0.05
0.10
0.15
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


MSY
B target
SPR target
Current
Equilibrium yield (mt)
0.00
0.05
0.10
0.15
0.0
0.2
0.4
0.6
0.8
1.0
Fraction unfished


Surplus production (mt)
0.00
0.10
0.20
0.30
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Total biomass (mt)


Yield per recruit (kg)
0.00
0.02
0.04
0.06
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


Catches
FISHERY
Abundance indices
FISHERY
Length compositions
FISHERY
1970
1980
1990
2000
2010
2020
Year


SR_LN(R0)
prior
max. likelihood
5
10
15
20
25
30
LnQ_base_FISHERY(1)
−6
−4
−2
0
2
4
Size_inflection_FISHERY(1)
15
20
25
30
35
40
Size_95%width_FISHERY(1)
0
1
2
3
4
5
Density
Parameter value